



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:09 PM UTC

PDB ID : 4QHL / pdb_00004qhl
Title : I3.2 (unbound) from CH103 Lineage
Authors : Fera, D.; Harrison, S.C.
Deposited on : 2014-05-28
Resolution : 3.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2361 (3.20-3.12)
Clashscore	190562	2486 (3.20-3.12)
Ramachandran outliers	187476	2405 (3.20-3.12)
Sidechain outliers	187428	2404 (3.20-3.12)
RSRZ outliers	180081	2361 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	
1	C	232	
2	B	213	
2	D	213	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called I3 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1670	1054	281	330	5			
1	C	219	Total	C	N	O	S	0	0	0
			1642	1038	276	323	5			


- Molecule 2 is a protein called UCA light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	209	Total	C	N	O	S	0	0	0
			1573	988	257	322	6			
2	D	209	Total	C	N	O	S	0	0	0
			1573	988	257	322	6			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: I3 heavy chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.08Å 99.14Å 163.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.57 – 3.15 49.57 – 3.15	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.57-3.15) 99.3 (49.57-3.15)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.62 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.237 , 0.281 0.235 , 0.279	Depositor DCC
R_{free} test set	1060 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	48.7	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	6458	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1712	0.74	1/2340 (0.0%)
1	C	0.32	0/1683	0.73	0/2300
2	B	0.33	0/1614	0.76	0/2206
2	D	0.30	0/1614	0.75	1/2206 (0.0%)
All	All	0.32	0/6623	0.75	2/9052 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	94	PHE	N-CA-C	-8.02	103.07	113.17
1	A	53	HIS	N-CA-C	-5.17	107.64	114.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1670	0	1634	27	0
1	C	1642	0	1603	16	0
2	B	1573	0	1513	12	0
2	D	1573	0	1513	14	0
All	All	6458	0	6263	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:TYR:HB2	1:A:95:LEU:HB2	1.68	0.73
2:B:92:ASP:O	2:B:94:PHE:N	2.22	0.72
1:A:7:SER:HB3	1:A:21:THR:HG22	1.71	0.71
1:A:82(A):ARG:NH1	2:D:49:TYR:OH	2.27	0.67
1:C:33:TYR:HB2	1:C:95:LEU:HB2	1.78	0.65
1:C:81:ARG:HH21	1:C:82(A):ARG:HD2	1.64	0.61
2:D:25:GLY:HA3	2:D:28:LEU:HB2	1.81	0.61
2:B:83:GLU:HB2	2:B:106:VAL:HG23	1.88	0.55
1:A:9:PRO:HD2	1:A:201:LYS:HD2	1.91	0.53
1:A:21:THR:O	1:A:21:THR:HG23	2.09	0.53
1:A:65:SER:HA	2:D:53:LYS:HA	1.89	0.53
1:C:18:LEU:HD13	1:C:109:VAL:HG11	1.90	0.53
1:A:66:ARG:NH1	1:A:82(B):SER:O	2.41	0.53
2:B:27:LYS:N	2:B:28:LEU:HA	2.26	0.51
1:C:87:THR:HG23	1:C:110:THR:HA	1.94	0.50
2:B:167:GLN:HE21	2:B:169:ASN:HB2	1.76	0.50
1:C:82:LEU:HD23	1:C:82(C):VAL:HG22	1.94	0.50
2:B:196:THR:HG23	2:B:201:THR:HG22	1.93	0.49
1:C:95:LEU:HD22	1:C:100(C):ALA:HB1	1.93	0.49
2:D:95:SER:O	2:D:95(A):THR:OG1	2.29	0.49
1:A:210:ARG:O	1:A:210:ARG:HG3	2.00	0.49
1:C:39:GLN:HB2	1:C:45:LEU:HD13	1.94	0.49
2:D:83:GLU:HB2	2:D:106:VAL:HG23	1.95	0.48
1:A:7:SER:HB3	1:A:21:THR:CG2	2.42	0.48
1:C:152:VAL:HG22	1:C:198:VAL:HG22	1.96	0.48
1:C:60:ASN:ND2	1:C:62:SER:OG	2.47	0.48
2:D:13:VAL:HG21	2:D:19:ALA:HB2	1.96	0.48
1:A:119:PRO:HB3	1:A:145:TYR:HB3	1.96	0.47
2:D:26:ASP:HA	2:D:27:LYS:HA	1.61	0.47
1:C:210:ARG:HG3	1:C:211:VAL:N	2.21	0.47
2:D:32:TYR:HB2	2:D:91:TRP:HZ3	1.79	0.47
1:A:210:ARG:O	1:A:210:ARG:CG	2.60	0.47
1:A:154:TRP:CH2	1:A:196:CYS:HB3	2.50	0.46
1:A:68:THR:HB	1:A:81:ARG:HB2	1.97	0.46
2:B:80:ALA:HA	2:B:106:VAL:HG11	1.97	0.46
1:A:7:SER:O	1:A:107:THR:HG21	2.14	0.46
1:C:47:TRP:CZ2	1:C:49:GLY:HA2	2.51	0.46
2:D:61:ARG:HB2	2:D:76:SER:O	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:LYS:HE3	1:C:85:ALA:HB1	1.98	0.46
2:B:61:ARG:HB2	2:B:76:SER:O	2.15	0.45
1:A:159:LEU:HD21	1:A:182:VAL:HG11	1.99	0.45
1:A:131:THR:C	1:A:133:GLY:HA2	2.41	0.45
1:A:38:ARG:NE	1:A:46:GLU:OE1	2.47	0.45
2:D:92:ASP:OD2	2:D:95:SER:OG	2.21	0.44
1:A:11:LEU:HB2	1:A:147:PRO:HG3	2.00	0.43
2:D:50:GLN:O	2:D:52:SER:N	2.47	0.43
2:D:28:LEU:HD12	2:D:28:LEU:HA	1.88	0.43
2:D:32:TYR:HB2	2:D:91:TRP:CZ3	2.53	0.43
2:B:4:LEU:HD22	2:B:23:CYS:SG	2.58	0.43
1:C:8:GLY:HA3	1:C:20:LEU:HD23	2.01	0.43
1:C:127:SER:OG	1:C:128:SER:N	2.52	0.43
1:A:47:TRP:CZ2	1:A:49:GLY:HA2	2.54	0.42
1:A:17:THR:HG23	1:A:82(A):ARG:HA	2.01	0.42
1:A:18:LEU:HD13	1:A:109:VAL:HG11	2.01	0.42
1:A:132:SER:N	1:A:133:GLY:HA2	2.35	0.42
2:B:26:ASP:HA	2:B:27:LYS:HA	1.73	0.42
2:B:27:LYS:HA	2:B:27:LYS:HD3	1.84	0.42
1:A:94:SER:OG	1:A:102:ASN:HB2	2.20	0.42
1:A:101:ASP:OD1	1:A:101:ASP:C	2.63	0.41
2:D:48:ILE:HD13	2:D:54:ARG:HB2	2.03	0.41
1:A:51:ILE:HG13	1:A:57:THR:HG22	2.02	0.41
1:C:95:LEU:HD23	1:C:100(D):PHE:O	2.20	0.41
1:C:64:GLU:O	1:C:65:SER:OG	2.31	0.40
2:B:15:PRO:HD3	2:B:106:VAL:O	2.22	0.40
2:B:48:ILE:HD13	2:B:54:ARG:HB2	2.02	0.40
1:A:100(E):PHE:O	1:A:103:TRP:NE1	2.48	0.40
1:A:146:PHE:HA	1:A:147:PRO:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/232 (95%)	203 (92%)	14 (6%)	4 (2%)	6	29
1	C	215/232 (93%)	204 (95%)	10 (5%)	1 (0%)	24	55
2	B	207/213 (97%)	194 (94%)	10 (5%)	3 (1%)	9	34
2	D	207/213 (97%)	185 (89%)	17 (8%)	5 (2%)	4	23
All	All	850/890 (96%)	786 (92%)	51 (6%)	13 (2%)	8	33

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	93	SER
1	A	130	SER
1	A	102	ASN
1	A	135	THR
2	B	151	ASP
1	C	102	ASN
2	D	151	ASP
2	D	26	ASP
2	D	92	ASP
2	D	95	SER
1	A	52	PHE
2	D	68	GLY
2	B	108	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/200 (96%)	187 (98%)	4 (2%)	47	67
1	C	187/200 (94%)	179 (96%)	8 (4%)	26	54
2	B	178/182 (98%)	169 (95%)	9 (5%)	21	50
2	D	178/182 (98%)	177 (99%)	1 (1%)	78	81
All	All	734/764 (96%)	712 (97%)	22 (3%)	36	62

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLU
1	A	52	PHE
1	A	169	VAL
1	A	210	ARG
2	B	11	VAL
2	B	22	THR
2	B	24	SER
2	B	42	GLN
2	B	60	GLU
2	B	123	GLU
2	B	149	LYS
2	B	170	ASN
2	B	201	THR
1	C	6	GLU
1	C	100	LEU
1	C	100(A)	VAL
1	C	100(B)	ASN
1	C	100(D)	PHE
1	C	100(E)	PHE
1	C	182	VAL
1	C	210	ARG
2	D	163	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	GLN
1	A	199	ASN
2	B	50	GLN
2	B	108	GLN
2	B	194	GLN
1	C	60	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/232 (96%)	0.82	29 (13%) 7 5	33, 53, 88, 130	0
1	C	219/232 (94%)	0.84	27 (12%) 8 5	30, 56, 89, 157	0
2	B	209/213 (98%)	0.73	24 (11%) 9 6	31, 57, 102, 130	0
2	D	209/213 (98%)	0.80	27 (12%) 7 5	35, 55, 96, 138	0
All	All	860/890 (96%)	0.80	107 (12%) 8 5	30, 55, 97, 157	0

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	94	PHE	5.8
1	C	100	LEU	5.7
2	B	95	SER	5.5
1	A	129	LYS	5.5
1	A	52	PHE	5.4
2	D	30	ASP	5.2
2	D	25	GLY	5.1
1	A	130	SER	4.9
2	D	24	SER	4.9
1	A	101	ASP	4.8
1	A	82(B)	SER	4.8
1	C	128	SER	4.5
2	D	94	PHE	4.5
2	B	95(A)	THR	4.0
1	C	134	GLY	4.0
2	D	209	THR	3.8
1	A	131	THR	3.7
2	B	169	ASN	3.6
2	D	91	TRP	3.6
2	D	93	SER	3.4
1	A	26	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	100(B)	ASN	3.4
2	D	88	CYS	3.4
1	C	97	ARG	3.4
2	B	23	CYS	3.3
2	D	95(A)	THR	3.3
1	A	75	GLU	3.3
2	D	71	ALA	3.3
1	C	164	HIS	3.3
1	A	3	GLN	3.2
1	C	100(A)	VAL	3.2
1	C	99	GLN	3.2
1	C	140	CYS	3.2
1	C	42	VAL	3.2
2	D	29	GLY	3.1
2	D	95	SER	3.1
1	A	132	SER	3.1
2	D	34	CYS	3.1
2	B	91	TRP	3.1
1	A	31	GLY	3.0
2	B	170	ASN	3.0
1	A	134	GLY	2.9
2	D	26	ASP	2.9
2	D	183	GLU	2.9
1	A	1	GLN	2.9
1	A	100(A)	VAL	2.8
2	D	168	SER	2.8
1	C	43	LYS	2.8
2	D	123	GLU	2.8
1	A	15	SER	2.8
1	C	58	ASN	2.7
2	B	2	TYR	2.7
1	A	100(B)	ASN	2.7
2	D	156	LYS	2.7
1	A	9	PRO	2.6
2	B	209	THR	2.6
1	A	186	SER	2.6
1	C	1	GLN	2.6
1	C	25	SER	2.6
2	B	93	SER	2.6
2	B	88	CYS	2.6
2	D	32	TYR	2.6
1	A	74	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	70	THR	2.5
2	B	92	ASP	2.5
1	C	113	SER	2.5
1	C	133	GLY	2.5
2	B	106	VAL	2.4
2	D	149	LYS	2.4
2	D	60	GLU	2.4
2	B	156	LYS	2.4
1	C	70	SER	2.4
2	B	187	SER	2.4
2	D	106	VAL	2.3
2	D	23	CYS	2.3
1	A	82(A)	ARG	2.3
1	C	28	SER	2.3
1	C	98	GLY	2.3
2	B	107	GLY	2.3
1	A	113	SER	2.3
1	C	82(B)	SER	2.3
2	B	183	GLU	2.3
1	C	100(C)	ALA	2.3
2	B	157	ALA	2.3
2	B	28	LEU	2.3
1	A	215	SER	2.3
2	D	187	SER	2.3
1	C	214	LYS	2.3
1	A	164	HIS	2.2
1	C	74	SER	2.2
2	B	168	SER	2.2
1	A	133	GLY	2.2
1	C	73	THR	2.2
2	B	69	ASN	2.2
2	B	41	GLY	2.2
2	D	126	GLN	2.1
1	A	27	GLY	2.1
1	A	97	ARG	2.1
1	C	105	GLN	2.1
1	A	16	GLU	2.1
1	C	100(D)	PHE	2.1
2	B	3	GLU	2.1
1	A	21	THR	2.1
2	B	165	SER	2.0
2	D	107	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	162	GLY	2.0
1	A	206	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.