



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 07:41 PM UTC

PDB ID : 6QIP / pdb_00006qip
Title : Ternary complex of FcRn ectodomain, FcRn binding optimised human serum albumin and the albumin-binding side chain of the human growth hormone derivative somapacitan
Authors : Johansson, E.
Deposited on : 2019-01-21
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	
2	B	274	
3	C	105	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serum albumin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	0	2	0
			4642	2928	784	888	42			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	418	MET	VAL	engineered mutation	UNP P02768
A	420	ALA	THR	engineered mutation	UNP P02768
A	505	GLY	GLU	engineered mutation	UNP P02768
A	547	ALA	VAL	engineered mutation	UNP P02768

- Molecule 2 is a protein called IgG receptor FcRn large subunit p51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	267	Total	C	N	O	S	0	1	0
			2106	1344	364	390	8			


- Molecule 3 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	100	Total	C	N	O	S	0	1	0
			844	537	144	160	3			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	100	HIS	-	expression tag	UNP P61769
C	101	HIS	-	expression tag	UNP P61769
C	102	HIS	-	expression tag	UNP P61769
C	103	HIS	-	expression tag	UNP P61769
C	104	HIS	-	expression tag	UNP P61769
C	105	HIS	-	expression tag	UNP P61769

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	12	6	1	4	1	0	0

- Molecule 5 is Somapacitan (CCD ID: JG5) (formula: $C_{51}H_{89}N_{11}O_{19}S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
5	A	1	82	51	11	19	1	50	0

- Molecule 6 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	4	0
			4	2	1	1		

- Molecule 7 is CYSTEINE (CCD ID: CYS) (formula: $C_3H_7NO_2S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	S	0	0
			6	3	1	1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	20	Total 20	O 20	0	0
8	B	4	Total 4	O 4	0	0
8	C	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serum albumin



- Molecule 2: IgG receptor FcRn large subunit p51



- Molecule 3: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.02Å 114.45Å 159.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.48 – 2.45 46.48 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.48-2.45) 98.2 (46.48-2.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.45Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.216 , 0.259 (Not available) , 0.247	Depositor DCC
R_{free} test set	2625 reflections (1.81%)	wwPDB-VP
Wilson B-factor (Å ²)	58.6	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 31.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7722	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: JG5, MES, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/4731	0.85	6/6377 (0.1%)
2	B	0.46	0/2172	0.82	3/2951 (0.1%)
3	C	0.46	0/868	0.80	1/1175 (0.1%)
All	All	0.48	0/7771	0.84	10/10503 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	438	CYS	CA-CB-SG	-7.92	96.17	114.40
1	A	313	LYS	CA-CB-CG	-7.35	99.40	114.10
2	B	3	SER	CA-C-N	7.13	133.26	122.93
2	B	3	SER	C-N-CA	7.13	133.26	122.93
1	A	392	CYS	CA-CB-SG	-7.03	98.23	114.40
3	C	58	LYS	CA-CB-CG	-6.07	101.97	114.10
2	B	139	GLN	CA-CB-CG	5.88	125.87	114.10
1	A	190	LYS	CA-CB-CG	5.38	124.85	114.10
1	A	313	LYS	CB-CG-CD	5.09	123.00	111.30
1	A	277	GLU	CA-CB-CG	5.06	124.22	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4642	0	4561	72	0
2	B	2106	0	2009	19	0
3	C	844	0	805	14	0
4	A	12	0	12	2	0
5	A	82	0	0	0	0
6	A	4	0	5	0	0
7	B	6	0	3	0	0
8	A	20	0	0	2	0
8	B	4	0	0	0	0
8	C	2	0	0	2	0
All	All	7722	0	7395	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:435:SER:O	1:A:439:LYS:HE3	1.08	1.21
1:A:435:SER:O	1:A:439:LYS:CE	2.03	1.06
1:A:84:TYR:HB3	1:A:87:MET:HE2	1.68	0.74
1:A:299:PRO:O	1:A:302:LEU:HD11	1.89	0.73
3:C:8:GLN:HG3	8:C:201:HOH:O	1.90	0.69
1:A:390:GLN:O	1:A:393:GLU:HG2	1.93	0.69
1:A:517:SER:HB2	1:A:519:LYS:NZ	2.10	0.67
3:C:81:ARG:HB2	3:C:92:ILE:HD13	1.76	0.67
1:A:433:VAL:HG22	1:A:452:TYR:CD1	2.30	0.67
1:A:302:LEU:HD12	1:A:302:LEU:N	2.12	0.65
2:B:100:PRO:O	2:B:102:ASN:N	2.29	0.64
1:A:156:PHE:CE2	1:A:160:ARG:HD2	2.33	0.63
1:A:519:LYS:HD3	1:A:519:LYS:H	1.67	0.60
1:A:194:ALA:HB1	1:A:455:VAL:CG1	2.33	0.58
1:A:313:LYS:HE2	1:A:367:HIS:CD2	2.38	0.58
1:A:453:LEU:HD11	4:A:601:MES:H21	1.86	0.58
1:A:31:LEU:HD22	1:A:84:TYR:CE1	2.38	0.58
3:C:33:SER:HB2	3:C:54:LEU:HD21	1.86	0.58
2:B:186:ALA:HB2	2:B:196:LEU:HD23	1.87	0.57
1:A:276:LYS:HG3	1:A:277:GLU:H	1.68	0.57
2:B:106:PRO:HG3	2:B:122:LEU:HD22	1.87	0.56
1:A:194:ALA:HB1	1:A:455:VAL:HG11	1.88	0.56
3:C:8:GLN:NE2	8:C:201:HOH:O	2.33	0.56
3:C:81:ARG:HD3	3:C:92:ILE:HD11	1.87	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:PRO:O	1:A:302:LEU:CD1	2.54	0.56
1:A:40:VAL:HA	1:A:43:VAL:HG12	1.87	0.55
1:A:479:GLU:HG3	8:A:705:HOH:O	2.05	0.55
2:B:187:ARG:NE	3:C:100:HIS:O	2.26	0.55
2:B:8:LEU:HD23	2:B:95:GLY:HA3	1.90	0.53
1:A:418:MET:HE2	2:B:59:TRP:CH2	2.43	0.53
2:B:109:LYS:HZ2	3:C:58:LYS:HE3	1.74	0.52
1:A:520:GLU:O	1:A:523:ILE:HG22	2.09	0.52
1:A:117:ARG:NH1	1:A:183:ASP:OD1	2.39	0.51
1:A:306:ALA:HA	1:A:310:VAL:HG22	1.92	0.51
1:A:475:LYS:O	1:A:479:GLU:HB2	2.09	0.51
1:A:383:GLU:HB3	1:A:384:PRO:HD3	1.93	0.51
1:A:559:CYS:O	1:A:564:LYS:HE3	2.12	0.50
1:A:67:HIS:NE2	1:A:249:ASP:OD1	2.44	0.50
1:A:331:LEU:HD13	1:A:350:ALA:HB2	1.94	0.49
1:A:475:LYS:NZ	8:A:705:HOH:O	2.40	0.49
1:A:37:GLU:HA	1:A:40:VAL:HG12	1.95	0.49
1:A:110:PRO:HB2	1:A:112:LEU:HG	1.93	0.49
1:A:342:SER:OG	1:A:344:VAL:HG12	2.13	0.49
1:A:393:GLU:O	1:A:397:GLN:HG2	2.13	0.49
3:C:49:VAL:HG12	3:C:68:THR:HB	1.95	0.49
1:A:315:VAL:HG13	1:A:316:CYS:N	2.28	0.48
1:A:367:HIS:HA	1:A:370:TYR:CZ	2.49	0.48
1:A:315:VAL:HG13	1:A:316:CYS:H	1.79	0.47
1:A:120:VAL:HG21	1:A:175:ALA:HA	1.96	0.47
2:B:126:THR:HA	2:B:142:GLN:OE1	2.14	0.47
1:A:42:LEU:HD22	1:A:73:LYS:HG3	1.96	0.47
1:A:314:ASP:O	1:A:317:LYS:N	2.46	0.47
1:A:519:LYS:HD3	1:A:519:LYS:N	2.28	0.47
2:B:196:LEU:O	2:B:239:SER:HA	2.15	0.47
3:C:48:LYS:HE2	3:C:48:LYS:HB2	1.74	0.46
1:A:310:VAL:O	1:A:370:TYR:HE1	1.98	0.46
1:A:314:ASP:O	1:A:315:VAL:C	2.58	0.46
1:A:348:ARG:HH22	1:A:450:GLU:CD	2.24	0.46
1:A:87:MET:HE2	1:A:87:MET:HB3	1.65	0.46
1:A:370:TYR:C	1:A:370:TYR:CD1	2.94	0.46
1:A:502:PHE:CD2	1:A:580:GLN:HG2	2.51	0.45
1:A:23:VAL:O	1:A:27:PHE:HD1	2.00	0.45
3:C:58:LYS:O	3:C:58:LYS:HG2	2.15	0.45
1:A:305:LEU:HD21	1:A:333:GLU:HB3	1.99	0.45
1:A:23:VAL:HG11	1:A:46:VAL:HG11	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:315:VAL:HG11	1:A:370:TYR:CZ	2.51	0.45
1:A:276:LYS:HG3	1:A:277:GLU:N	2.30	0.45
2:B:187:ARG:O	2:B:187:ARG:HG3	2.17	0.45
2:B:189:SER:HB2	2:B:195:VAL:HG23	1.99	0.44
2:B:168:GLU:HA	2:B:168:GLU:OE2	2.17	0.44
1:A:42:LEU:HD22	1:A:73:LYS:CG	2.48	0.44
1:A:56:ASP:C	1:A:58:SER:H	2.25	0.44
2:B:109:LYS:NZ	3:C:58:LYS:HE3	2.32	0.44
2:B:189:SER:OG	2:B:190:SER:N	2.45	0.44
3:C:1:ILE:HG13	3:C:2:GLN:N	2.32	0.43
1:A:31:LEU:HG	1:A:74:LEU:HD22	2.00	0.43
4:A:601:MES:H82	4:A:601:MES:H31	1.41	0.43
2:B:74:LEU:HD23	2:B:74:LEU:HA	1.86	0.43
1:A:31:LEU:CD2	1:A:87:MET:HE1	2.49	0.43
1:A:179:LEU:N	1:A:180:PRO:HD2	2.34	0.43
1:A:283:LEU:HD23	1:A:284:LEU:HD12	2.01	0.42
1:A:12:LYS:HB3	1:A:12:LYS:HE3	1.65	0.42
1:A:359:LYS:HA	1:A:359:LYS:HD2	1.41	0.42
1:A:573:LYS:NZ	3:C:69:GLU:OE2	2.53	0.42
1:A:274:LYS:NZ	1:A:294:GLU:OE2	2.50	0.42
1:A:311:GLU:O	1:A:312:SER:C	2.62	0.42
2:B:38:TYR:CD1	2:B:38:TYR:C	2.97	0.42
1:A:77:VAL:O	1:A:80:LEU:HD13	2.19	0.41
1:A:73:LYS:HA	1:A:73:LYS:HD3	1.51	0.41
1:A:517:SER:HB2	1:A:519:LYS:HZ3	1.82	0.41
2:B:247:GLU:HB3	2:B:267:LEU:HD11	2.02	0.41
3:C:1:ILE:HG13	3:C:2:GLN:H	1.85	0.41
1:A:289:CYS:O	1:A:293:VAL:HG13	2.20	0.41
1:A:20:LYS:HG2	1:A:47:THR:OG1	2.20	0.41
1:A:416:PRO:O	1:A:534:LYS:HE2	2.21	0.41
1:A:233:LYS:HD3	1:A:263:TYR:OH	2.20	0.41
1:A:274:LYS:HE3	1:A:296:ASP:HA	2.03	0.41
1:A:325:VAL:O	1:A:329:MET:HG3	2.20	0.41
2:B:167:LEU:HD23	2:B:174:LEU:HD13	2.03	0.41
1:A:40:VAL:HA	1:A:43:VAL:CG1	2.49	0.40
2:B:267:LEU:HA	2:B:267:LEU:HD23	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/585 (100%)	564 (97%)	16 (3%)	3 (0%)	24	32
2	B	266/274 (97%)	254 (96%)	7 (3%)	5 (2%)	6	5
3	C	99/105 (94%)	99 (100%)	0	0	100	100
All	All	948/964 (98%)	917 (97%)	23 (2%)	8 (1%)	16	20

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	100	PRO
2	B	101	ASP
2	B	85	LYS
1	A	57	GLU
1	A	130	ASN
1	A	315	VAL
2	B	84	GLY
2	B	83	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	509/508 (100%)	509 (100%)	0	100	100
2	B	222/226 (98%)	222 (100%)	0	100	100
3	C	96/100 (96%)	96 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	827/834 (99%)	827 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	367	HIS
1	A	397	GLN
2	B	4	HIS
2	B	124	GLN
2	B	215	ASN
3	C	83	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	CYS	B	301	2	4,5,6	1.62	1 (25%)	1,5,7	1.71	0
4	MES	A	601	-	12,12,12	2.55	1 (8%)	15,16,16	2.29	6 (40%)
5	JG5	A	602	6	82,82,82	2.30	18 (21%)	93,99,99	1.66	9 (9%)
6	BME	A	603	5	3,3,3	0.50	0	2,2,2	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	CYS	B	301	2	-	1/1/4/6	-
4	MES	A	601	-	-	5/6/14/14	0/1/1/1
5	JG5	A	602	6	-	46/94/94/94	0/1/1/1
6	BME	A	603	5	-	1/1/1/1	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	601	MES	C8-S	-8.51	1.65	1.77
5	A	602	JG5	C40-N42	6.68	1.48	1.34
5	A	602	JG5	C30-N32	6.38	1.48	1.33
5	A	602	JG5	C62-N64	5.92	1.46	1.34
5	A	602	JG5	C52-N54	5.92	1.47	1.33
5	A	602	JG5	C46-N48	5.64	1.46	1.34
5	A	602	JG5	S24-N23	5.41	1.74	1.63
5	A	602	JG5	C29-C30	4.48	1.60	1.51
5	A	602	JG5	C71-N70	4.48	1.46	1.34
5	A	602	JG5	C21-N23	4.25	1.50	1.39
5	A	602	JG5	N12-N11	4.15	1.37	1.31
5	A	602	JG5	O25-S24	3.90	1.48	1.43
5	A	602	JG5	C39-C40	3.53	1.58	1.51
5	A	602	JG5	O26-S24	3.05	1.47	1.43
7	B	301	CYS	CB-SG	2.76	1.87	1.81
5	A	602	JG5	O47-C46	-2.64	1.18	1.23
5	A	602	JG5	N10-N11	2.50	1.37	1.32
5	A	602	JG5	C51-C52	2.46	1.56	1.51
5	A	602	JG5	C27-S24	2.42	1.85	1.78
5	A	602	JG5	N12-N13	2.33	1.36	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	602	JG5	O26-S24-O25	-10.58	104.55	119.34
4	A	601	MES	C5-N4-C3	4.55	118.63	108.84
4	A	601	MES	C7-N4-C3	3.81	121.39	111.24
5	A	602	JG5	C09-N13-N12	3.41	107.50	102.45
5	A	602	JG5	N13-N12-N11	-3.38	106.46	113.52
4	A	601	MES	O1S-S-C8	3.08	111.38	106.73
4	A	601	MES	C7-N4-C5	3.01	119.27	111.24
4	A	601	MES	C2-C3-N4	-2.96	105.62	110.12
5	A	602	JG5	C45-C46-N48	2.47	120.21	115.86
5	A	602	JG5	C73-C71-N70	2.26	119.98	116.12
5	A	602	JG5	C29-C30-N32	2.26	120.46	116.34
5	A	602	JG5	C61-C62-N64	2.23	120.26	116.41
4	A	601	MES	O2S-S-O1S	-2.08	107.07	113.82
5	A	602	JG5	C51-C52-N54	2.03	120.04	116.34
5	A	602	JG5	O38-C39-C40	-2.00	108.26	112.41

There are no chirality outliers.

All (53) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	601	MES	C8-C7-N4-C3
4	A	601	MES	N4-C7-C8-S
5	A	602	JG5	C61-C62-N64-C65
5	A	602	JG5	C21-N23-S24-O25
6	A	603	BME	O1-C1-C2-S2
7	B	301	CYS	N-CA-CB-SG
5	A	602	JG5	O63-C62-N64-C65
5	A	602	JG5	N32-C33-C34-O35
5	A	602	JG5	O57-C58-C59-O60
5	A	602	JG5	N54-C55-C56-O57
5	A	602	JG5	O35-C36-C37-O38
5	A	602	JG5	C50-C51-C52-N54
5	A	602	JG5	C50-C51-C52-O53
5	A	602	JG5	N48-C49-C50-C51
5	A	602	JG5	C37-C36-O35-C34
5	A	602	JG5	C17-C18-C19-C20
5	A	602	JG5	O60-C61-C62-N64
5	A	602	JG5	C18-C19-C20-C21
5	A	602	JG5	C06-C07-C08-C09
5	A	602	JG5	C03-C04-C05-C06
5	A	602	JG5	O38-C39-C40-N42
5	A	602	JG5	C66-C67-C68-C69
5	A	602	JG5	C43-C44-C45-C46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	602	JG5	C49-C50-C51-C52
5	A	602	JG5	C04-C05-C06-C07
5	A	602	JG5	O38-C39-C40-O41
5	A	602	JG5	O60-C61-C62-O63
5	A	602	JG5	N64-C65-C78-O79
5	A	602	JG5	N64-C65-C78-O80
5	A	602	JG5	N48-C49-C81-O82
5	A	602	JG5	N48-C49-C81-O83
5	A	602	JG5	C50-C49-C81-O83
5	A	602	JG5	S24-C27-C28-C29
5	A	602	JG5	C16-C17-C18-C19
5	A	602	JG5	C21-N23-S24-O26
5	A	602	JG5	C07-C08-C09-N10
5	A	602	JG5	C50-C49-C81-O82
4	A	601	MES	C7-C8-S-O2S
5	A	602	JG5	C81-C49-C50-C51
5	A	602	JG5	C55-C56-O57-C58
5	A	602	JG5	C07-C08-C09-N13
5	A	602	JG5	C21-N23-S24-C27
5	A	602	JG5	C67-C68-C69-N70
4	A	601	MES	C7-C8-S-O3S
5	A	602	JG5	C14-C15-C16-C17
5	A	602	JG5	C58-C59-O60-C61
5	A	602	JG5	C05-C06-C07-C08
5	A	602	JG5	C33-C34-O35-C36
4	A	601	MES	C7-C8-S-O1S
5	A	602	JG5	C62-C61-O60-C59
5	A	602	JG5	C59-C58-O57-C56
5	A	602	JG5	C66-C65-C78-O79
5	A	602	JG5	C66-C65-C78-O80


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	601	MES	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	582/585 (99%)	0.49	25 (4%) 40 39	18, 72, 101, 115	2 (0%)
2	B	267/274 (97%)	0.38	12 (4%) 38 37	32, 63, 99, 120	1 (0%)
3	C	100/105 (95%)	0.44	4 (4%) 42 41	27, 67, 96, 107	1 (1%)
All	All	949/964 (98%)	0.45	41 (4%) 40 39	18, 69, 100, 120	4 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	78	ALA	4.3
1	A	80	LEU	4.0
1	A	77	VAL	3.6
1	A	70	PHE	3.6
3	C	100	HIS	3.5
1	A	115	LEU	3.5
3	C	98	ASP	3.4
1	A	84	TYR	3.4
3	C	99	MET	3.2
2	B	83	GLY	3.2
1	A	585	LEU	3.1
1	A	79	THR	3.0
2	B	5	LEU	3.0
2	B	3	SER	2.8
1	A	330	PHE	2.8
2	B	124	GLN	2.7
2	B	100	PRO	2.7
1	A	371	ALA	2.6
2	B	142	GLN	2.4
1	A	26	ALA	2.4
2	B	221	THR	2.3
1	A	439	LYS	2.3
1	A	4	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	128	GLY	2.3
1	A	12	LYS	2.2
1	A	300	ALA	2.2
1	A	185	LEU	2.2
1	A	111	ASN	2.2
2	B	85	LYS	2.2
3	C	58	LYS	2.2
1	A	74	LEU	2.1
2	B	138	SER	2.1
1	A	172	ALA	2.1
1	A	30	TYR	2.1
1	A	359	LYS	2.1
2	B	4	HIS	2.1
1	A	39	HIS	2.1
1	A	327	LEU	2.0
1	A	271	ILE	2.0
1	A	270	SER	2.0
2	B	135	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	JG5	A	602	82/82	0.86	0.16	13,30,85,113	50
7	CYS	B	301	6/7	0.93	0.13	56,65,72,72	0
6	BME	A	603	4/4	-	-	13,13,13,13	4
4	MES	A	601	12/12	0.94	0.10	68,79,82,90	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.