



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:19 PM UTC

PDB ID : 1QJ6 / pdb_00001qj6
Title : Novel Covalent Active Site Thrombin Inhibitors
Authors : Jhoti, H.; Cleasby, A.
Deposited on : 1999-06-22
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	36	58% 25% 14% .
2	B	259	61% 27% 11% .
3	I	10	60% 40%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	36	Total	C	N	O	S	0	0	0
			287	177	48	61	1			


- Molecule 2 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	259	Total	C	N	O	S	0	0	0
			2093	1334	370	375	14			

- Molecule 3 is a protein called HIRUGEN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	10	Total	C	N	O	S	0	0	0
			94	59	10	24	1			

- Molecule 4 is 6-CARBAMIMIDOYL-2-[2-HYDROXY-5-(3-METHOXY-PHENYL)-INDAN-1-YL]-HEXANOIC ACID (CCD ID: 167) (formula: C₂₃H₂₈N₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			28	23	2	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	34	Total	O	0	0
			34	34		
5	B	163	Total	O	0	0
			163	163		
5	I	6	Total	O	0	0
			6	6		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: THROMBIN

Chain A:  58% 25% 14% .



• Molecule 2: THROMBIN

Chain B:  61% 27% 11% .



• Molecule 3: HIRUGEN

Chain I:  60% 40%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.80Å 72.00Å 73.20Å 90.00° 101.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.20	Depositor
% Data completeness (in resolution range)	98.0 (15.00-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.200 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2705	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TYS, 167

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	0/290	1.84	6/384 (1.6%)
2	B	1.27	6/2148 (0.3%)	2.01	71/2903 (2.4%)
3	I	0.97	0/79	1.90	3/103 (2.9%)
All	All	1.25	6/2517 (0.2%)	1.99	80/3390 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	3
All	All	0	4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	195	SER	CB-OG	-8.83	1.24	1.42
2	B	85	LEU	CA-C	-5.98	1.45	1.52
2	B	75	ARG	NE-CZ	5.38	1.39	1.33
2	B	143	ASN	CA-C	-5.17	1.46	1.52
2	B	119	HIS	CE1-NE2	5.16	1.37	1.32
2	B	57	HIS	ND1-CE1	5.11	1.37	1.32

All (80) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	77(A)	ARG	CA-C-N	-11.12	105.00	122.67
2	B	77(A)	ARG	C-N-CA	-11.12	105.00	122.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	66	VAL	N-CA-CB	-8.86	101.07	110.72
2	B	244	GLN	N-CA-C	8.85	120.93	111.28
2	B	205	ASN	N-CA-C	7.68	122.36	113.38
3	I	56	PHE	CA-CB-CG	-7.52	106.28	113.80
2	B	195	SER	CA-CB-OG	7.47	126.04	111.10
2	B	186(A)	ASP	N-CA-C	7.41	120.41	111.82
2	B	245	PHE	N-CA-C	7.39	119.52	107.32
2	B	34	PHE	CA-C-N	-7.32	109.56	122.07
2	B	34	PHE	C-N-CA	-7.32	109.56	122.07
2	B	149(A)	ALA	N-CA-C	7.28	121.41	110.64
2	B	149(B)	ASN	N-CA-C	7.23	121.35	110.14
2	B	129(B)	SER	N-CA-C	7.04	119.99	111.82
2	B	77(A)	ARG	N-CA-C	7.03	119.54	111.11
2	B	146	GLU	N-CA-C	6.78	120.47	110.30
2	B	154	VAL	N-CA-CB	-6.73	99.48	111.93
2	B	148	TRP	CA-C-N	6.40	133.76	121.54
2	B	148	TRP	C-N-CA	6.40	133.76	121.54
2	B	232	PHE	CA-CB-CG	-6.32	107.48	113.80
2	B	126	ARG	CA-CB-CG	6.27	126.65	114.10
2	B	147	THR	N-CA-C	6.26	122.82	114.12
2	B	241	VAL	N-CA-C	6.26	118.91	111.09
2	B	242	ILE	CA-CB-CG1	6.20	120.94	110.40
2	B	60(B)	PRO	N-CA-C	6.20	118.27	110.70
2	B	74	THR	N-CA-C	6.13	121.04	113.50
2	B	70	LYS	N-CA-C	6.10	120.02	110.32
1	A	9	LYS	N-CA-C	6.04	120.11	112.87
1	A	1(C)	GLU	N-CA-C	6.00	119.68	110.20
2	B	192	GLU	N-CA-C	5.99	118.49	110.35
2	B	236	LYS	N-CA-C	5.86	117.67	111.28
2	B	146	GLU	CA-CB-CG	-5.82	102.47	114.10
2	B	170	ASP	N-CA-C	5.82	120.11	113.19
2	B	243	ASP	N-CA-C	5.81	117.69	111.36
2	B	216	GLY	CA-C-N	-5.79	115.32	122.84
2	B	216	GLY	C-N-CA	-5.79	115.32	122.84
2	B	201	MET	CA-C-N	-5.70	114.86	122.72
2	B	201	MET	C-N-CA	-5.70	114.86	122.72
3	I	61	GLU	CA-C-N	-5.67	111.48	121.70
3	I	61	GLU	C-N-CA	-5.67	111.48	121.70
2	B	75	ARG	CA-CB-CG	5.67	125.44	114.10
2	B	232	PHE	N-CA-C	5.59	118.09	111.33
2	B	169	LYS	CA-CB-CG	5.54	125.18	114.10
2	B	112	VAL	CB-CA-C	-5.52	103.45	111.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	129(A)	ALA	CA-C-O	5.52	126.81	120.90
2	B	57	HIS	N-CA-C	5.47	119.06	112.38
2	B	24	ILE	CB-CG1-CD1	-5.45	102.35	113.80
2	B	187	ARG	N-CA-C	5.44	117.53	109.69
2	B	169	LYS	CB-CA-C	5.41	119.77	110.79
2	B	60(H)	PHE	CA-C-O	5.39	126.83	120.69
2	B	79	ILE	N-CA-CB	-5.38	107.22	111.64
2	B	117	TYR	CA-C-O	5.36	124.95	119.05
2	B	75	ARG	CB-CG-CD	5.34	123.57	111.30
1	A	1(A)	ASP	CA-C-O	5.29	125.68	119.28
2	B	129(A)	ALA	N-CA-C	5.28	117.44	111.11
2	B	176	ILE	CA-CB-CG2	5.27	119.46	110.50
2	B	195	SER	CA-C-N	-5.26	114.78	122.46
2	B	195	SER	C-N-CA	-5.26	114.78	122.46
2	B	204(B)	ASN	OD1-CG-ND2	-5.26	117.34	122.60
2	B	49	ASP	N-CA-C	5.21	119.26	113.01
2	B	61	GLU	N-CA-C	5.21	117.36	111.11
2	B	141	TRP	CB-CA-C	-5.18	102.68	111.02
2	B	129(B)	SER	CB-CA-C	-5.16	100.76	110.67
1	A	14(H)	GLU	N-CA-C	5.16	118.83	112.54
1	A	1	CYS	N-CA-C	5.13	117.67	110.23
2	B	143	ASN	CA-CB-CG	-5.13	107.47	112.60
2	B	62	ASN	CA-CB-CG	-5.12	107.48	112.60
2	B	101	ARG	CA-CB-CG	-5.12	103.86	114.10
1	A	8	GLU	N-CA-C	5.11	117.25	111.11
2	B	148	TRP	CD2-CE2-CZ2	-5.11	117.29	122.40
2	B	33	LEU	CA-C-N	-5.08	115.60	122.77
2	B	33	LEU	C-N-CA	-5.08	115.60	122.77
2	B	171	SER	CA-C-O	5.08	124.94	119.15
2	B	221	ARG	CD-NE-CZ	-5.08	117.29	124.40
2	B	235	LYS	N-CA-C	5.08	117.55	111.71
2	B	116	ASP	N-CA-C	5.07	118.56	112.38
2	B	95	ASN	N-CA-C	5.05	117.39	108.24
2	B	51	TRP	CA-C-O	5.05	125.92	120.32
2	B	35	ARG	N-CA-C	5.03	118.32	110.32
2	B	39	GLU	N-CA-C	5.03	116.89	108.99

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1(E)	SER	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	B	246	GLY	Peptide
2	B	35	ARG	Sidechain
2	B	73	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	287	0	278	6	0
2	B	2093	0	2063	29	0
3	I	94	0	73	1	0
4	B	28	0	25	4	0
5	A	34	0	0	1	0
5	B	163	0	0	0	0
5	I	6	0	0	0	0
All	All	2705	0	2439	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:75:ARG:HD2	2:B:77:GLU:HG2	1.49	0.95
1:A:1(C):GLU:HA	1:A:1(B):ALA:HB3	1.66	0.76
2:B:166:PRO:HA	2:B:169:LYS:HD2	1.71	0.73
2:B:130:LEU:HD23	2:B:162:ILE:HD13	1.77	0.66
2:B:35:ARG:HG2	2:B:41:LEU:HD21	1.79	0.64
2:B:148:TRP:CZ3	2:B:220:CYS:SG	2.93	0.62
2:B:32:MET:HG3	2:B:40:LEU:HD12	1.81	0.60
2:B:147:THR:HA	2:B:149(A):ALA:HA	1.83	0.60
2:B:204(B):ASN:C	2:B:204(B):ASN:HD22	2.16	0.53
1:A:14(G):LEU:HD21	2:B:202:LYS:HD3	1.91	0.52
1:A:1(C):GLU:HG2	2:B:120:PRO:HG3	1.92	0.52
2:B:146:GLU:O	2:B:149(B):ASN:HB2	2.08	0.52
1:A:1(C):GLU:HA	1:A:1(B):ALA:CB	2.37	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:164:GLU:HB2	2:B:167:VAL:HG13	1.91	0.51
2:B:73:ARG:HD3	2:B:152:PRO:O	2.12	0.50
2:B:240:LYS:HB2	2:B:240:LYS:NZ	2.28	0.49
2:B:145:LYS:HB3	2:B:149(B):ASN:HB3	1.96	0.48
2:B:148:TRP:NE1	2:B:192:GLU:HG3	2.29	0.47
2:B:129:ALA:HA	2:B:210:MET:HE1	1.98	0.46
2:B:85:LEU:HD13	2:B:106:MET:HE2	1.97	0.46
2:B:148:TRP:CZ2	4:B:1248:167:H11	2.52	0.45
2:B:101:ARG:HG2	2:B:234:LEU:HD21	1.97	0.45
4:B:1248:167:H11	4:B:1248:167:H36	1.98	0.44
2:B:144:LEU:HD21	2:B:152:PRO:HB3	1.99	0.43
3:I:58:GLU:H	3:I:58:GLU:CD	2.26	0.43
2:B:35:ARG:O	2:B:38:GLN:HA	2.18	0.42
2:B:198:PRO:HB2	2:B:200:VAL:HG13	2.00	0.42
2:B:87:LYS:HD3	2:B:88:ILE:N	2.34	0.42
1:A:14(K):ILE:HA	5:A:2028:HOH:O	2.20	0.42
2:B:61:GLU:HG3	2:B:88:ILE:HG13	2.01	0.41
2:B:98:ASN:HA	4:B:1248:167:H533	2.03	0.41
2:B:24:ILE:HG21	2:B:24:ILE:HD13	1.89	0.41
2:B:216:GLY:O	4:B:1248:167:H37	2.21	0.41
1:A:14(J):TYR:O	1:A:14(K):ILE:HG13	2.21	0.40
2:B:85:LEU:CD1	2:B:106:MET:HE2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	34/36 (94%)	25 (74%)	5 (15%)	4 (12%)	0	0
2	B	257/259 (99%)	242 (94%)	11 (4%)	4 (2%)	7	6
3	I	7/10 (70%)	7 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	298/305 (98%)	274 (92%)	16 (5%)	8 (3%)	4	2

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14(L)	ASP
2	B	37	SER
2	B	149	THR
2	B	149(E)	LYS
1	A	1(B)	ALA
2	B	149(C)	VAL
1	A	1(G)	PHE
1	A	1(F)	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	31/31 (100%)	25 (81%)	6 (19%)	1	1
2	B	225/225 (100%)	193 (86%)	32 (14%)	3	3
3	I	9/9 (100%)	9 (100%)	0	100	100
All	All	265/265 (100%)	227 (86%)	38 (14%)	3	3

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1(H)	THR
1	A	1(G)	PHE
1	A	1(E)	SER
1	A	1(C)	GLU
1	A	1(A)	ASP
1	A	1	CYS
2	B	33	LEU
2	B	46	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	50	ARG
2	B	60	LEU
2	B	60(A)	TYR
2	B	60(B)	PRO
2	B	60(C)	PRO
2	B	60(D)	TRP
2	B	60(E)	ASP
2	B	60(F)	LYS
2	B	60(G)	ASN
2	B	60(H)	PHE
2	B	60(I)	THR
2	B	64	LEU
2	B	66	VAL
2	B	81	LYS
2	B	112	VAL
2	B	145	LYS
2	B	149	THR
2	B	149(B)	ASN
2	B	149(C)	VAL
2	B	149(E)	LYS
2	B	154	VAL
2	B	167	VAL
2	B	169	LYS
2	B	192	GLU
2	B	195	SER
2	B	236	LYS
2	B	240	LYS
2	B	241	VAL
2	B	245	PHE
2	B	247	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	143	ASN
2	B	204(B)	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TYS	I	63	3	11,15,17	0.82	1 (9%)	10,19,24	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	I	63	3	-	0/7/10/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	63	TYS	OH-CZ	2.46	1.46	1.42

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	167	B	1248	2	29,30,31	1.05	1 (3%)	34,41,43	2.23	10 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	167	B	1248	2	-	8/19/31/33	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1248	167	C16-C15	3.72	1.56	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1248	167	C12-C10-C16	6.96	127.19	112.90
4	B	1248	167	C11-C12-C10	-5.43	106.41	110.84
4	B	1248	167	C11-C14-C13	-3.13	100.57	103.43
4	B	1248	167	C39-C11-C12	3.12	123.85	120.80
4	B	1248	167	O9-C13-C10	3.03	116.95	110.48
4	B	1248	167	C38-C39-C11	-2.71	117.90	121.88
4	B	1248	167	C4-C5-N6	2.67	124.02	117.15
4	B	1248	167	C37-C36-C12	-2.57	116.87	121.12
4	B	1248	167	C53-O54-C45	-2.53	112.08	117.50
4	B	1248	167	C36-C12-C10	2.37	134.26	129.66

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1248	167	C2-C1-C16-C10
4	B	1248	167	C3-C4-C5-N6
4	B	1248	167	C3-C4-C5-N7
4	B	1248	167	C12-C10-C16-C1
4	B	1248	167	C24-C45-O54-C53
4	B	1248	167	C44-C45-O54-C53
4	B	1248	167	C13-C10-C16-C1
4	B	1248	167	O8-C15-C16-C1

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1248	167	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.