



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:23 AM UTC

PDB ID : 1QJ9 / pdb_00001qj9
Title : CRYSTAL STRUCTURE OF THE OUTER MEMBRANE PROTEIN OMPX
FROM ESCHERICHIA COLI
Authors : Vogt, J.; Schulz, G.E.
Deposited on : 1999-06-23
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	24% 65% 22% 7% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1225 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OUTER MEMBRANE PROTEIN X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	0	0	0
			1158	731	190	234	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ASN	HIS	engineered mutation	UNP P36546


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	67	Total	O	0	0
			67	67		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: OUTER MEMBRANE PROTEIN X



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	82.30Å 82.30Å 208.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.00 – 2.10 35.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (35.00-2.10) 99.6 (35.00-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.10Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.233 , 0.284 0.251 , 0.296	Depositor DCC
R_{free} test set	799 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 82.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.026 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.018 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1225	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	5/1188 (0.4%)	2.17	49/1608 (3.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	21	MET	CG-SD	8.57	2.02	1.80
1	A	21	MET	SD-CE	6.60	1.96	1.79
1	A	132	ILE	N-CA	6.42	1.54	1.46
1	A	2	THR	CB-OG1	-6.03	1.34	1.43
1	A	86	GLY	CA-C	5.47	1.57	1.52

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	ASN	CA-CB-CG	18.06	130.66	112.60
1	A	50	ARG	NE-CZ-NH2	15.62	133.25	119.20
1	A	21	MET	CB-CA-C	13.58	132.50	110.78
1	A	21	MET	O-C-N	-13.27	107.26	123.05
1	A	21	MET	CG-SD-CE	-11.76	75.03	100.90
1	A	17	GLN	N-CA-C	11.10	124.57	111.02
1	A	131	ARG	O-C-N	-9.46	109.62	121.92
1	A	19	ASN	CA-CB-CG	8.96	121.56	112.60
1	A	17	GLN	CA-C-O	-8.42	111.81	121.07
1	A	2	THR	CA-CB-OG1	8.39	122.18	109.60
1	A	1	ALA	N-CA-CB	8.02	122.43	110.40
1	A	114	GLN	CB-CG-CD	7.66	125.63	112.60
1	A	95	TYR	N-CA-C	7.52	118.25	109.60
1	A	53	SER	N-CA-C	-7.25	104.06	113.12
1	A	21	MET	CA-C-O	7.24	128.07	120.54
1	A	21	MET	CA-CB-CG	7.01	128.12	114.10
1	A	118	MET	O-C-N	-6.88	114.78	123.24
1	A	60	ASN	CA-CB-CG	6.75	119.35	112.60
1	A	50	ARG	NE-CZ-NH1	-6.64	114.86	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	TYR	CA-C-N	-6.62	117.15	121.65
1	A	80	TYR	C-N-CA	-6.62	117.15	121.65
1	A	119	GLU	N-CA-C	6.38	119.04	111.71
1	A	99	LYS	CB-CG-CD	6.31	125.82	111.30
1	A	1	ALA	CA-C-O	-6.24	110.19	120.80
1	A	147	ARG	NE-CZ-NH1	-6.20	115.30	121.50
1	A	99	LYS	CG-CD-CE	6.15	125.45	111.30
1	A	100	ASN	CA-CB-CG	6.13	118.73	112.60
1	A	133	ARG	CG-CD-NE	6.12	125.47	112.00
1	A	8	GLY	CA-C-O	6.08	127.19	121.41
1	A	125	PHE	CA-CB-CG	6.07	119.87	113.80
1	A	50	ARG	NH1-CZ-NH2	-5.94	111.58	119.30
1	A	26	LEU	CA-C-O	-5.69	114.11	120.66
1	A	129	GLN	OE1-CD-NE2	5.57	128.17	122.60
1	A	50	ARG	CA-CB-CG	5.46	125.03	114.10
1	A	7	GLY	CA-C-O	5.42	126.12	121.19
1	A	40	ILE	CA-C-N	-5.39	118.46	122.18
1	A	40	ILE	C-N-CA	-5.39	118.46	122.18
1	A	52	ALA	CA-C-O	5.35	128.16	120.51
1	A	43	PHE	CA-CB-CG	-5.25	108.55	113.80
1	A	118	MET	CA-CB-CG	5.24	124.57	114.10
1	A	131	ARG	CB-CA-C	5.19	119.04	110.83
1	A	25	ASN	OD1-CG-ND2	5.16	127.76	122.60
1	A	58	ASN	N-CA-CB	5.13	120.32	111.13
1	A	140	TRP	CA-CB-CG	5.04	123.18	113.60
1	A	43	PHE	CA-C-N	-5.04	114.75	122.62
1	A	43	PHE	C-N-CA	-5.04	114.75	122.62
1	A	75	ASP	CA-C-N	-5.02	114.64	122.37
1	A	75	ASP	C-N-CA	-5.02	114.64	122.37
1	A	99	LYS	CA-CB-CG	5.01	124.11	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1158	0	1068	43	0
2	A	67	0	0	9	0
All	All	1225	0	1068	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:MET:CG	1:A:21:MET:SD	2.02	1.48
1:A:2:THR:HB	2:A:2002:HOH:O	1.46	1.15
1:A:21:MET:CG	1:A:21:MET:CE	2.42	0.96
1:A:21:MET:CE	1:A:21:MET:CB	2.49	0.90
1:A:140:TRP:HB2	2:A:2064:HOH:O	1.74	0.87
1:A:131:ARG:HD3	2:A:2049:HOH:O	1.80	0.80
1:A:21:MET:CB	1:A:21:MET:HE3	2.13	0.79
1:A:2:THR:HG23	1:A:31:GLU:HB2	1.63	0.78
1:A:21:MET:HE3	1:A:21:MET:HB3	1.65	0.78
1:A:50:ARG:HH11	1:A:50:ARG:HG3	1.49	0.77
1:A:21:MET:HB2	1:A:21:MET:HE2	1.67	0.76
1:A:32:GLU:OE1	2:A:2026:HOH:O	2.07	0.71
1:A:21:MET:CE	1:A:21:MET:HB2	2.21	0.70
1:A:17:GLN:HA	2:A:2017:HOH:O	1.93	0.67
1:A:131:ARG:HB3	2:A:2062:HOH:O	2.01	0.60
1:A:47:GLU:HB2	2:A:2033:HOH:O	2.01	0.60
1:A:95:TYR:CG	1:A:96:PRO:HA	2.36	0.60
1:A:118:MET:HB2	1:A:121:VAL:HG22	1.85	0.58
1:A:48:LYS:HE3	1:A:50:ARG:CD	2.34	0.58
1:A:50:ARG:HG3	1:A:50:ARG:NH1	2.20	0.57
1:A:14:ALA:O	1:A:15:GLN:C	2.48	0.56
1:A:95:TYR:CD1	1:A:96:PRO:HA	2.42	0.54
1:A:131:ARG:O	1:A:132:ILE:HB	2.09	0.53
1:A:17:GLN:O	1:A:18:MET:C	2.49	0.51
1:A:131:ARG:O	1:A:135:VAL:O	2.29	0.51
1:A:48:LYS:HE3	1:A:50:ARG:NE	2.26	0.51
1:A:15:GLN:HG2	2:A:2063:HOH:O	2.11	0.50
1:A:51:THR:O	1:A:52:ALA:HB2	2.11	0.49
1:A:17:GLN:HA	1:A:17:GLN:HE21	1.77	0.49
1:A:19:ASN:HD22	1:A:20:LYS:H	1.60	0.49
1:A:113:LEU:HD12	1:A:113:LEU:O	2.13	0.48
1:A:14:ALA:O	1:A:16:GLY:N	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:LYS:HE3	1:A:50:ARG:HD2	1.99	0.45
1:A:17:GLN:HE21	1:A:17:GLN:CA	2.29	0.44
1:A:2:THR:CG2	1:A:31:GLU:HB2	2.41	0.44
1:A:132:ILE:O	1:A:133:ARG:HB2	2.16	0.44
1:A:5:VAL:O	1:A:145:GLY:HA3	2.18	0.43
1:A:17:GLN:NE2	2:A:2017:HOH:O	2.53	0.42
1:A:51:THR:HB	1:A:52:ALA:H	1.61	0.42
1:A:21:MET:CG	1:A:21:MET:HE2	2.44	0.41
1:A:18:MET:SD	1:A:21:MET:SD	3.18	0.41
1:A:131:ARG:O	1:A:132:ILE:C	2.61	0.41
1:A:133:ARG:HB3	1:A:134:SER:H	1.66	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	146/148 (99%)	135 (92%)	6 (4%)	5 (3%)	3 1

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	19	ASN
1	A	15	GLN
1	A	54	SER
1	A	16	GLY
1	A	132	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	118/118 (100%)	104 (88%)	14 (12%)	5 3

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	15	GLN
1	A	17	GLN
1	A	19	ASN
1	A	21	MET
1	A	34	ASN
1	A	50	ARG
1	A	51	THR
1	A	58	ASN
1	A	99	LYS
1	A	113	LEU
1	A	120	ASN
1	A	129	GLN
1	A	133	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	19	ASN
1	A	34	ASN
1	A	60	ASN
1	A	91	GLN
1	A	100	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	148/148 (100%)	1.32	36 (24%) 2 2	34, 48, 99, 133	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	GLY	7.1
1	A	18	MET	6.7
1	A	21	MET	5.3
1	A	52	ALA	5.0
1	A	54	SER	4.6
1	A	99	LYS	4.4
1	A	134	SER	4.3
1	A	140	TRP	4.2
1	A	118	MET	4.1
1	A	60	ASN	4.1
1	A	90	PHE	3.6
1	A	57	TYR	3.5
1	A	95	TYR	3.4
1	A	55	GLY	3.4
1	A	19	ASN	3.3
1	A	132	ILE	3.3
1	A	13	ASP	3.2
1	A	20	LYS	3.1
1	A	146	TYR	3.0
1	A	30	TYR	3.0
1	A	49	SER	2.9
1	A	97	THR	2.9
1	A	17	GLN	2.9
1	A	51	THR	2.8
1	A	135	VAL	2.8
1	A	94	GLU	2.7
1	A	133	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	98	TYR	2.5
1	A	92	THR	2.5
1	A	107	PHE	2.4
1	A	56	ASP	2.4
1	A	58	ASN	2.4
1	A	50	ARG	2.3
1	A	15	GLN	2.1
1	A	125	PHE	2.1
1	A	101	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.