



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:30 PM UTC

PDB ID : 1QJF / pdb_00001qjf
Title : ISOPENICILLIN N SYNTHASE FROM ASPERGILLUS NIDULANS (Monocyclic Sulfoxide - Fe COMPLEX)
Authors : Rutledge, P.J.; Clifton, I.J.; Burzlaff, N.I.; Roach, P.L.; Adlington, R.M.; Baldwin, J.E.
Deposited on : 1999-06-23
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ISOPENICILLIN N SYNTHASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	328	2639	1686	445	503	5	0	0	0

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is 1-[(1S)-CARBOXY-2-(METHYLSULFINYL)ETHYL]-(3R)-[(5S)-5-AMINO-5-CARBOXYPENTANAMIDO]-(4R)-SULFANYLAZETIDIN-2-ONE (CCD ID: ACS) (formula: C₁₃H₂₁N₃O₇S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			25	13	3	7	2		

- Molecule 4 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	410	Total	O	0	0
			410	410		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ISOPENICILLIN N SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.78Å 71.19Å 101.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.40 8.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	85.3 (8.00-1.40) 77.3 (8.00-1.40)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.75 (at 1.40Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.131 , 0.205 0.167 , 0.195	Depositor DCC
R_{free} test set	2770 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	10.3	Xtriage
Anisotropy	0.619	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.51 , 81.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3080	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, SO4, ACS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.51	33/2714 (1.2%)	2.04	92/3693 (2.5%)

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	324	LEU	C-N	10.93	1.47	1.33
1	A	293	VAL	C-N	-8.73	1.22	1.33
1	A	157	LEU	C-N	-8.48	1.22	1.33
1	A	331	THR	C-OXT	8.33	1.40	1.23
1	A	296	PRO	C-N	-7.33	1.23	1.33
1	A	244	ALA	C-N	-7.14	1.23	1.33
1	A	311	LEU	C-N	-7.11	1.24	1.33
1	A	321	LEU	C-N	-7.05	1.25	1.33
1	A	230	ASN	C-N	6.88	1.42	1.33
1	A	217	VAL	C-N	6.30	1.41	1.33
1	A	115	PRO	C-N	-6.28	1.25	1.34
1	A	64	SER	C-N	-6.18	1.26	1.33
1	A	136	PRO	C-N	-6.18	1.24	1.33
1	A	70	LYS	C-N	6.13	1.42	1.33
1	A	237	ALA	C-N	-6.11	1.23	1.33
1	A	247	THR	C-N	-5.97	1.27	1.33
1	A	327	LYS	C-N	5.95	1.42	1.33
1	A	280	GLN	C-N	-5.70	1.26	1.33
1	A	257	MET	C-N	-5.53	1.26	1.33
1	A	68	GLU	C-N	5.50	1.41	1.33
1	A	27	ARG	C-N	-5.49	1.26	1.33
1	A	103	PHE	C-N	-5.46	1.25	1.33
1	A	251	ILE	C-N	5.41	1.40	1.33
1	A	190	PRO	C-N	-5.41	1.25	1.33
1	A	255	SER	C-N	-5.35	1.26	1.33
1	A	4	VAL	C-N	-5.31	1.26	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	230	ASN	N-CA	5.28	1.53	1.46
1	A	48	GLY	C-N	-5.19	1.23	1.33
1	A	269	ILE	C-O	-5.18	1.18	1.24
1	A	114	HIS	CG-ND1	-5.16	1.32	1.38
1	A	21	ASP	C-N	5.14	1.41	1.34
1	A	256	TYR	N-CA	5.09	1.52	1.46
1	A	222	VAL	C-N	-5.03	1.26	1.33

All (92) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	ARG	CD-NE-CZ	25.46	160.04	124.40
1	A	315	ASP	CA-CB-CG	-17.95	94.65	112.60
1	A	20	ASP	CA-CB-CG	-11.28	101.32	112.60
1	A	68	GLU	CA-C-O	10.23	131.40	120.55
1	A	263	ASN	OD1-CG-ND2	-9.96	112.64	122.60
1	A	8	ASN	OD1-CG-ND2	9.76	132.36	122.60
1	A	213	TRP	CA-C-O	9.54	131.68	121.19
1	A	324	LEU	CA-C-O	9.40	130.52	120.55
1	A	47	HIS	ND1-CG-CD2	-9.39	96.71	106.10
1	A	47	HIS	ND1-CE1-NE2	-9.01	99.39	108.40
1	A	319	ASN	OD1-CG-ND2	-8.93	113.67	122.60
1	A	8	ASN	CA-CB-CG	-8.46	104.14	112.60
1	A	263	ASN	CB-CG-ND2	8.38	128.97	116.40
1	A	195	TYR	CA-C-O	-8.11	113.01	119.66
1	A	326	ASN	CA-CB-CG	-8.00	104.60	112.60
1	A	259	HIS	CG-CD2-NE2	7.82	115.02	107.20
1	A	68	GLU	O-C-N	-7.67	113.99	122.12
1	A	47	HIS	CG-ND1-CE1	7.31	121.73	109.30
1	A	47	HIS	CB-CG-CD2	7.29	140.68	131.20
1	A	115	PRO	CA-C-N	7.26	130.73	120.28
1	A	115	PRO	C-N-CA	7.26	130.73	120.28
1	A	193	ASP	CA-C-O	-7.15	112.19	120.34
1	A	141	PHE	CA-CB-CG	-7.07	106.73	113.80
1	A	114	HIS	ND1-CE1-NE2	-6.91	101.49	108.40
1	A	205	ASP	CA-CB-CG	6.83	119.43	112.60
1	A	324	LEU	CA-C-N	-6.83	110.72	120.42
1	A	324	LEU	C-N-CA	-6.83	110.72	120.42
1	A	40	GLY	CA-C-O	-6.79	111.46	119.06
1	A	262	ASN	CA-C-O	6.64	129.24	121.54
1	A	205	ASP	CA-C-N	-6.51	113.35	122.86
1	A	205	ASP	C-N-CA	-6.51	113.35	122.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	ALA	CA-C-O	-6.50	113.99	121.47
1	A	30	GLN	CA-C-O	-6.47	113.69	120.55
1	A	293	VAL	O-C-N	-6.47	116.21	123.20
1	A	8	ASN	O-C-N	6.41	131.42	122.77
1	A	326	ASN	CA-C-O	-6.39	113.78	120.55
1	A	318	GLN	CA-CB-CG	6.33	126.76	114.10
1	A	303	ASN	CA-C-O	6.30	126.90	119.28
1	A	241	ASP	CA-C-O	6.25	127.81	120.69
1	A	328	ASN	O-C-N	6.24	129.79	122.24
1	A	62	HIS	CA-CB-CG	6.15	119.95	113.80
1	A	37	ARG	NE-CZ-NH2	-6.14	113.67	119.20
1	A	47	HIS	CG-CD2-NE2	6.11	113.31	107.20
1	A	319	ASN	CB-CG-ND2	6.06	125.49	116.40
1	A	62	HIS	CA-C-O	5.97	126.85	120.10
1	A	209	LEU	N-CA-C	5.97	119.27	109.72
1	A	76	ARG	NE-CZ-NH2	5.95	124.56	119.20
1	A	243	GLU	CB-CG-CD	5.95	122.71	112.60
1	A	48	GLY	CA-C-N	5.94	131.41	122.91
1	A	48	GLY	C-N-CA	5.94	131.41	122.91
1	A	134	LYS	CG-CD-CE	5.94	124.96	111.30
1	A	248	GLY	O-C-N	5.74	129.46	123.09
1	A	328	ASN	CA-C-O	-5.69	112.55	119.32
1	A	90	TYR	CA-C-O	5.68	127.45	120.92
1	A	71	TRP	CD2-CE2-CZ2	5.67	128.07	122.40
1	A	308	ARG	NE-CZ-NH2	5.60	124.24	119.20
1	A	259	HIS	CB-CG-ND1	5.56	131.04	122.70
1	A	55	SER	CB-CA-C	5.56	120.30	110.85
1	A	302	PRO	O-C-N	5.54	128.62	122.24
1	A	311	LEU	CA-C-O	-5.54	114.17	120.32
1	A	113	ASP	O-C-N	-5.50	114.78	122.37
1	A	48	GLY	O-C-N	-5.50	115.45	122.43
1	A	312	SER	O-C-N	5.49	129.59	122.89
1	A	103	PHE	CA-C-O	-5.46	114.45	120.24
1	A	157	LEU	CA-C-O	-5.44	113.72	119.97
1	A	300	ARG	NE-CZ-NH1	-5.42	116.08	121.50
1	A	5	SER	CA-CB-OG	5.40	121.90	111.10
1	A	59	LYS	CA-C-O	5.37	126.46	120.82
1	A	306	SER	CA-C-O	-5.36	114.84	121.11
1	A	8	ASN	CB-CA-C	-5.35	102.53	110.67
1	A	259	HIS	ND1-CG-CD2	-5.33	100.77	106.10
1	A	114	HIS	CG-ND1-CE1	5.33	118.35	109.30
1	A	176	LYS	CG-CD-CE	5.30	123.48	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	TYR	OH-CZ-CE2	5.28	135.75	119.90
1	A	292	SER	CA-C-O	5.28	126.94	120.92
1	A	38	ASP	CA-CB-CG	5.27	117.87	112.60
1	A	313	TYR	CA-CB-CG	-5.21	104.52	113.90
1	A	261	THR	CA-C-O	5.16	125.60	119.56
1	A	243	GLU	CA-CB-CG	-5.14	103.82	114.10
1	A	309	GLU	CA-C-O	5.12	125.21	120.60
1	A	30	GLN	O-C-N	5.10	127.53	122.12
1	A	124	HIS	CG-CD2-NE2	5.10	112.30	107.20
1	A	325	ILE	CA-C-N	-5.08	113.47	120.28
1	A	325	ILE	C-N-CA	-5.08	113.47	120.28
1	A	125	GLU	CA-C-O	-5.08	115.30	120.99
1	A	122	PRO	CA-C-O	-5.06	115.64	121.36
1	A	320	GLY	O-C-N	5.06	127.10	122.19
1	A	259	HIS	CA-C-N	5.06	127.06	120.28
1	A	259	HIS	C-N-CA	5.06	127.06	120.28
1	A	134	LYS	CA-C-O	5.06	125.58	119.31
1	A	147	TRP	CZ3-CH2-CZ2	-5.05	114.94	121.50
1	A	275	VAL	CA-C-O	-5.03	115.03	120.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2639	0	2522	6	0
2	A	5	0	0	0	0
3	A	25	0	18	1	0
4	A	1	0	0	0	0
5	A	410	0	0	2	0
All	All	3080	0	2540	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:LYS:HG2	1:A:7:ALA:N	2.19	0.51
1:A:131:ASP:OD1	1:A:133:THR:OG1	2.29	0.50
3:A:1333:ACS:H291	5:A:2291:HOH:O	2.13	0.49
1:A:167:GLU:OE2	1:A:169:ASN:HB2	2.13	0.48
1:A:166:LYS:NZ	5:A:2240:HOH:O	2.50	0.44
1:A:262:ASN:O	1:A:300:ARG:NH2	2.51	0.44
1:A:314:GLY:O	1:A:318:GLN:HG2	2.19	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/331 (98%)	316 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/284 (99%)	273 (97%)	9 (3%)	34	7

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	SER
1	A	6	LYS
1	A	8	ASN
1	A	20	ASP
1	A	55	SER
1	A	91	TYR
1	A	263	ASN
1	A	306	SER
1	A	324	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	240	GLN
1	A	259	HIS
1	A	287	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACS	A	1333	4	21,25,25	1.50	2 (9%)	25,35,35	2.44	8 (32%)
2	SO4	A	1332	-	4,4,4	0.55	0	6,6,6	1.69	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACS	A	1333	4	-	1/25/42/42	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1333	ACS	C24-C22	-3.83	1.48	1.53
3	A	1333	ACS	O35-C23	-3.54	1.19	1.30

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1333	ACS	C23-C22-N21	7.22	119.10	110.26
3	A	1333	ACS	C24-C22-N21	5.61	114.76	111.43
3	A	1333	ACS	O36-C23-C22	-3.79	110.80	122.44
3	A	1333	ACS	O35-C23-O36	3.10	131.11	124.08
2	A	1332	SO4	O4-S-O2	-2.90	94.38	109.56
3	A	1333	ACS	O20-C19-N21	2.60	135.33	131.75
2	A	1332	SO4	O4-S-O1	2.40	122.12	109.56
3	A	1333	ACS	C22-N21-C19	-2.39	125.41	131.57
3	A	1333	ACS	C6-C2-N1	2.31	116.14	110.12
3	A	1333	ACS	C42-N21-C19	2.12	96.51	93.92

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1333	ACS	N21-C22-C24-S25

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1333	ACS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/331 (99%)	-0.39	1 (0%) 90 92	8, 15, 30, 40	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	1332	5/5	0.89	0.12	35,37,40,40	0
3	ACS	A	1333	25/25	0.95	0.07	12,20,28,42	0
4	FE2	A	1334	1/1	1.00	0.03	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.