



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:26 PM UTC

PDB ID : 1QJP / pdb_00001qjp
Title : HIGH RESOLUTION STRUCTURE OF THE OUTER MEMBRANE PROTEIN A (OMPA) TRANSMEMBRANE DOMAIN
Authors : Pautsch, A.; Schulz, G.E.
Deposited on : 1999-06-29
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	171	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C8E	A	1174	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called OUTER MEMBRANE PROTEIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	137	Total	C	N	O	S	0	6	0
			1101	715	178	204	4			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	PHE	LEU	engineered mutation	UNP P02934
A	34	LYS	GLN	engineered mutation	UNP P02934
A	107	TYR	LYS	engineered mutation	UNP P02934

- Molecule 2 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (CCD ID: C8E) (formula: C₁₆H₃₄O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	16	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	5	0
			21	16	5		
2	A	1	Total	C	O	6	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	2	0
			21	16	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	64	Total	O	0	0
			64	64		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: OUTER MEMBRANE PROTEIN A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	65.10Å 79.70Å 50.20Å 90.00° 94.30° 90.00°	Depositor
Resolution (Å)	12.00 – 1.65 12.00 – 1.65	Depositor EDS
% Data completeness (in resolution range)	95.4 (12.00-1.65) 94.9 (12.00-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 1.65Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.155 , 0.198 0.218 , 0.248	Depositor DCC
R_{free} test set	1479 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 83.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1291	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: C8E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	1/1165 (0.1%)	1.71	17/1582 (1.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	71	ALA	C-N	5.36	1.40	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	ASP	CA-CB-CG	13.50	126.10	112.60
1	A	76	GLY	O-C-N	8.56	130.16	123.23
1	A	32	GLU	N-CA-C	7.25	119.34	107.23
1	A	87	ILE	N-CA-C	-7.08	103.28	111.00
1	A	32	GLU	CB-CA-C	-7.00	102.25	111.42
1	A	17	GLN	OE1-CD-NE2	-6.55	116.05	122.60
1	A	114	ASN	CA-C-O	-6.32	114.50	121.33
1	A	120	SER	O-C-N	6.17	126.17	121.85
1	A	72	TYR	N-CA-CB	5.93	119.91	110.65
1	A	114	ASN	CA-CB-CG	-5.88	106.72	112.60
1	A	87	ILE	CB-CA-C	-5.71	104.58	112.24
1	A	32	GLU	O-C-N	-5.69	116.97	123.40
1	A	116	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	5	ASN	CA-CB-CG	5.43	118.03	112.60
1	A	71	ALA	N-CA-CB	5.14	118.10	110.40
1	A	139	LEU	CD1-CG-CD2	5.13	122.08	110.80
1	A	72	TYR	N-CA-C	-5.00	100.25	108.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1101	0	1040	53	0
2	A	126	0	200	21	0
3	A	64	0	0	5	0
All	All	1291	0	1240	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93[A]:ILE:CD1	1:A:127[A]:VAL:HG12	1.76	1.15
1:A:93[B]:ILE:HD12	2:A:1174:C8E:H21	1.26	1.13
1:A:93[A]:ILE:HD13	1:A:127[A]:VAL:HG12	1.09	1.06
1:A:93[A]:ILE:HD13	1:A:127[A]:VAL:CG1	1.89	1.01
1:A:93[B]:ILE:CD1	2:A:1174:C8E:H21	1.90	1.01
1:A:93[A]:ILE:CD1	1:A:127[A]:VAL:CG1	2.40	0.98
1:A:93[B]:ILE:CD1	2:A:1174:C8E:C1	2.46	0.94
1:A:93[B]:ILE:HD12	2:A:1174:C8E:C2	1.99	0.92
1:A:35:LEU:HB2	1:A:61:MET:HE3	1.50	0.91
1:A:35:LEU:HB2	1:A:61:MET:CE	2.05	0.86
1:A:93[B]:ILE:CD1	2:A:1174:C8E:C2	2.54	0.85
1:A:101:VAL:HG13	2:A:1175:C8E:H32	1.57	0.85
1:A:139:LEU:HD21	2:A:1173:C8E:H62	1.61	0.80
1:A:60:ARG:NH2	1:A:72:TYR:CD1	2.56	0.73
1:A:93[B]:ILE:HD13	2:A:1174:C8E:C1	2.22	0.67
1:A:17:GLN:HE22	2:A:1173:C8E:H202	1.59	0.67
1:A:93[B]:ILE:CD1	2:A:1174:C8E:H11	2.22	0.67
1:A:5:ASN:HB2	3:A:2010:HOH:O	1.94	0.66
1:A:1:ALA:N	3:A:2001:HOH:O	2.22	0.66
1:A:93[B]:ILE:HD13	2:A:1174:C8E:H12	1.78	0.65
1:A:84:GLY:HA3	3:A:2043:HOH:O	2.00	0.62
1:A:93[A]:ILE:HD11	1:A:127[A]:VAL:CG1	2.29	0.60
1:A:101:VAL:HG12	1:A:119:VAL:HG22	1.83	0.59
1:A:60:ARG:NH2	1:A:72:TYR:CE1	2.71	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93[B]:ILE:HD11	2:A:1174:C8E:H11	1.85	0.58
1:A:91:LEU:HD22	1:A:127[B]:VAL:CG1	2.34	0.57
1:A:35:LEU:HD12	1:A:61:MET:HE3	1.88	0.56
1:A:101:VAL:CG1	2:A:1175:C8E:H32	2.34	0.56
1:A:93[B]:ILE:HD13	2:A:1174:C8E:C2	2.36	0.55
1:A:94:TYR:HB3	3:A:2043:HOH:O	2.09	0.52
1:A:91:LEU:HD22	1:A:127[B]:VAL:HG12	1.92	0.51
1:A:83:LEU:HD12	1:A:83:LEU:N	2.26	0.51
1:A:77:VAL:HG21	2:A:1175:C8E:H141	1.92	0.51
1:A:87:ILE:HG12	1:A:93[A]:ILE:HG12	1.93	0.50
1:A:35:LEU:CB	1:A:61:MET:HE3	2.33	0.50
1:A:91:LEU:CD2	1:A:127[B]:VAL:CG1	2.89	0.49
1:A:35:LEU:HB2	1:A:61:MET:HE2	1.89	0.47
1:A:92:ASP:HB3	3:A:2043:HOH:O	2.14	0.47
1:A:93[A]:ILE:HD11	1:A:127[A]:VAL:HG12	1.81	0.47
1:A:93[B]:ILE:HG23	2:A:1174:C8E:H21	1.96	0.47
1:A:93[A]:ILE:CD1	1:A:127[A]:VAL:HG13	2.40	0.46
1:A:60:ARG:NH2	1:A:72:TYR:CG	2.84	0.46
1:A:17:GLN:HE22	2:A:1173:C8E:C20	2.29	0.45
1:A:53:MET:HE3	1:A:53:MET:HB2	1.74	0.44
1:A:71:ALA:HB3	1:A:107:TYR:HD2	1.82	0.44
1:A:51:PHE:CG	2:A:1172:C8E:H51	2.53	0.43
1:A:60:ARG:CZ	1:A:72:TYR:CG	3.02	0.43
1:A:91:LEU:CD2	1:A:127[B]:VAL:HG12	2.47	0.43
1:A:87:ILE:CG1	1:A:93[A]:ILE:HG12	2.49	0.42
1:A:93[B]:ILE:HD13	2:A:1174:C8E:H21	1.86	0.42
1:A:60:ARG:NH1	1:A:72:TYR:CD2	2.89	0.41
1:A:139:LEU:CD2	2:A:1173:C8E:H62	2.43	0.41
1:A:143:TRP:HE1	2:A:1177:C8E:H131	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/171 (79%)	134 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/133 (84%)	106 (95%)	6 (5%)	20	4

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	GLU
1	A	91	LEU
1	A	107	TYR
1	A	111	TYR
1	A	139	LEU
1	A	145	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	75	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	C8E	A	1177	-	20,20,20	2.33	1 (5%)	19,19,19	1.65	3 (15%)
2	C8E	A	1173	-	20,20,20	0.89	0	19,19,19	1.69	5 (26%)
2	C8E	A	1174	-	20,20,20	1.76	1 (5%)	19,19,19	3.02	8 (42%)
2	C8E	A	1176	-	20,20,20	0.89	0	19,19,19	1.96	6 (31%)
2	C8E	A	1172	-	20,20,20	0.92	0	19,19,19	2.01	6 (31%)
2	C8E	A	1175	-	20,20,20	0.95	0	19,19,19	1.97	7 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C8E	A	1177	-	-	12/18/18/18	-
2	C8E	A	1173	-	-	11/18/18/18	-
2	C8E	A	1174	-	-	12/18/18/18	-
2	C8E	A	1176	-	-	11/18/18/18	-
2	C8E	A	1172	-	-	10/18/18/18	-
2	C8E	A	1175	-	-	13/18/18/18	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1177	C8E	C19-C20	9.80	2.01	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1174	C8E	C17-C16	-6.81	1.14	1.49

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1174	C8E	O15-C16-C17	7.90	146.34	110.35
2	A	1174	C8E	O18-C17-C16	6.37	139.38	110.35
2	A	1174	C8E	O15-C14-C13	5.39	134.92	110.35
2	A	1175	C8E	O15-C14-C13	4.66	131.59	110.35
2	A	1172	C8E	O15-C14-C13	4.55	131.09	110.35
2	A	1176	C8E	O15-C14-C13	4.29	129.90	110.35
2	A	1177	C8E	O15-C14-C13	4.15	129.27	110.35
2	A	1173	C8E	O15-C14-C13	4.04	128.76	110.35
2	A	1172	C8E	O12-C13-C14	3.40	125.86	110.35
2	A	1174	C8E	O18-C19-C20	3.38	124.99	110.11
2	A	1175	C8E	O18-C17-C16	3.35	125.62	110.35
2	A	1176	C8E	O18-C19-C20	3.18	124.12	110.11
2	A	1176	C8E	O12-C13-C14	3.15	124.71	110.35
2	A	1172	C8E	O18-C19-C20	3.06	123.59	110.11
2	A	1175	C8E	C7-C6-C5	-3.06	98.92	114.37
2	A	1177	C8E	O18-C17-C16	3.04	124.19	110.35
2	A	1172	C8E	O9-C8-C7	3.04	126.26	110.32
2	A	1176	C8E	O18-C17-C16	2.99	124.00	110.35
2	A	1174	C8E	O12-C13-C14	2.95	123.81	110.35
2	A	1172	C8E	C7-C6-C5	-2.93	99.55	114.37
2	A	1175	C8E	O18-C19-C20	2.92	122.99	110.11
2	A	1173	C8E	O18-C17-C16	2.85	123.32	110.35
2	A	1172	C8E	O18-C17-C16	2.77	122.95	110.35
2	A	1174	C8E	C5-C4-C3	-2.73	100.57	114.37
2	A	1173	C8E	C7-C6-C5	-2.57	101.38	114.37
2	A	1173	C8E	O18-C19-C20	2.40	120.71	110.11
2	A	1175	C8E	C5-C4-C3	-2.36	102.44	114.37
2	A	1176	C8E	C5-C4-C3	-2.36	102.45	114.37
2	A	1174	C8E	O9-C8-C7	2.32	122.52	110.32
2	A	1173	C8E	O12-C13-C14	2.29	120.80	110.35
2	A	1175	C8E	O12-C13-C14	2.27	120.70	110.35
2	A	1176	C8E	C7-C6-C5	-2.27	102.91	114.37
2	A	1177	C8E	C7-C6-C5	-2.26	102.95	114.37
2	A	1174	C8E	C7-C6-C5	-2.10	103.73	114.37
2	A	1175	C8E	O9-C8-C7	2.05	121.07	110.32

There are no chirality outliers.

All (69) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1176	C8E	O12-C13-C14-O15
2	A	1175	C8E	O12-C13-C14-O15
2	A	1176	C8E	O15-C16-C17-O18
2	A	1173	C8E	O12-C13-C14-O15
2	A	1174	C8E	O12-C13-C14-O15
2	A	1172	C8E	O15-C16-C17-O18
2	A	1173	C8E	C6-C7-C8-O9
2	A	1172	C8E	C6-C7-C8-O9
2	A	1174	C8E	C6-C7-C8-O9
2	A	1175	C8E	C6-C7-C8-O9
2	A	1174	C8E	O15-C16-C17-O18
2	A	1177	C8E	O12-C13-C14-O15
2	A	1174	C8E	C16-C17-O18-C19
2	A	1172	C8E	O12-C13-C14-O15
2	A	1172	C8E	C2-C3-C4-C5
2	A	1174	C8E	C5-C6-C7-C8
2	A	1173	C8E	C3-C4-C5-C6
2	A	1176	C8E	C2-C3-C4-C5
2	A	1177	C8E	C3-C4-C5-C6
2	A	1173	C8E	O15-C16-C17-O18
2	A	1174	C8E	C3-C4-C5-C6
2	A	1176	C8E	C4-C5-C6-C7
2	A	1173	C8E	C4-C5-C6-C7
2	A	1177	C8E	C2-C3-C4-C5
2	A	1172	C8E	C3-C4-C5-C6
2	A	1175	C8E	C3-C4-C5-C6
2	A	1175	C8E	C2-C3-C4-C5
2	A	1172	C8E	C7-C8-O9-C10
2	A	1173	C8E	C5-C6-C7-C8
2	A	1177	C8E	C4-C5-C6-C7
2	A	1174	C8E	C1-C2-C3-C4
2	A	1175	C8E	O15-C16-C17-O18
2	A	1176	C8E	C1-C2-C3-C4
2	A	1174	C8E	C4-C5-C6-C7
2	A	1177	C8E	C5-C6-C7-C8
2	A	1177	C8E	C16-C17-O18-C19
2	A	1177	C8E	O15-C16-C17-O18
2	A	1175	C8E	C14-C13-O12-C11
2	A	1174	C8E	C14-C13-O12-C11
2	A	1177	C8E	C14-C13-O12-C11
2	A	1177	C8E	C1-C2-C3-C4
2	A	1172	C8E	C20-C19-O18-C17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1174	C8E	C10-C11-O12-C13
2	A	1175	C8E	C20-C19-O18-C17
2	A	1177	C8E	C10-C11-O12-C13
2	A	1176	C8E	C20-C19-O18-C17
2	A	1172	C8E	C14-C13-O12-C11
2	A	1173	C8E	C2-C3-C4-C5
2	A	1176	C8E	C13-C14-O15-C16
2	A	1175	C8E	C16-C17-O18-C19
2	A	1176	C8E	C5-C6-C7-C8
2	A	1172	C8E	C10-C11-O12-C13
2	A	1174	C8E	C20-C19-O18-C17
2	A	1175	C8E	C4-C5-C6-C7
2	A	1176	C8E	C6-C7-C8-O9
2	A	1175	C8E	C10-C11-O12-C13
2	A	1177	C8E	C20-C19-O18-C17
2	A	1175	C8E	C1-C2-C3-C4
2	A	1174	C8E	C7-C8-O9-C10
2	A	1172	C8E	O18-C19-C20-O21
2	A	1177	C8E	C13-C14-O15-C16
2	A	1176	C8E	C11-C10-O9-C8
2	A	1173	C8E	C10-C11-O12-C13
2	A	1175	C8E	C13-C14-O15-C16
2	A	1176	C8E	C16-C17-O18-C19
2	A	1175	C8E	C7-C8-O9-C10
2	A	1173	C8E	C20-C19-O18-C17
2	A	1173	C8E	C16-C17-O18-C19
2	A	1173	C8E	C13-C14-O15-C16


There are no ring outliers.


5 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1177	C8E	1	0
2	A	1173	C8E	4	0
2	A	1174	C8E	12	0
2	A	1172	C8E	1	0
2	A	1175	C8E	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	137/171 (80%)	0.94	18 (13%) 7 8	15, 33, 88, 120	6 (4%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	72	TYR	7.8
1	A	63	TYR	4.6
1	A	71	ALA	4.4
1	A	129[A]	TYR	4.4
1	A	107	TYR	4.1
1	A	31	HIS	4.0
1	A	109	ASN	3.8
1	A	108	SER	3.7
1	A	110	VAL	3.6
1	A	145	ASN	3.5
1	A	106	THR	2.8
1	A	85[A]	TYR	2.5
1	A	105	ASP	2.3
1	A	3	LYS	2.1
1	A	5	ASN	2.1
1	A	111	TYR	2.1
1	A	62	PRO	2.0
1	A	116	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	C8E	A	1177	21/21	0.72	0.32	82,86,117,121	21
2	C8E	A	1176	21/21	0.76	0.33	34,83,113,115	21
2	C8E	A	1174	21/21	0.79	0.27	35,75,129,131	21
2	C8E	A	1175	21/21	0.81	0.30	45,94,126,126	21
2	C8E	A	1173	21/21	0.83	0.24	21,71,97,97	21
2	C8E	A	1172	21/21	0.88	0.19	30,46,78,80	21

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around C8E A 1176:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C8E A 1174:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around C8E A 1175:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around C8E A 1173:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.