



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:44 PM UTC

PDB ID : 3QJ4 / pdb_00003qj4
Title : Crystal structure of Human Renalase (isoform 1)
Authors : Milani, M.; Ciriello, F.; Baroni, S.; Pandini, V.; Aliverti, A.; Canevari, G.; Bolognesi, M.
Deposited on : 2011-01-28
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	10% 77% 20% ..
1	B	342	10% 75% 20% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Renalase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	1	0
			2604	1648	448	489	19			
1	B	333	Total	C	N	O	S	0	1	0
			2593	1645	447	484	17			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37	ASP	GLU	SEE REMARK 999	UNP Q5VYX0
B	37	ASP	GLU	SEE REMARK 999	UNP Q5VYX0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	67	Total	O	0	0
			67	67		
4	B	61	Total	O	0	0
			61	61		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Renalase



• Molecule 1: Renalase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.84Å 86.59Å 93.18Å 90.00° 95.30° 90.00°	Depositor
Resolution (Å)	28.94 – 2.50 28.94 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (28.94-2.50) 84.4 (28.94-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, R_{free}	0.217 , 0.260 0.243 , 0.296	Depositor DCC
R_{free} test set	1288 reflections (4.34%)	wwPDB-VP
Wilson B-factor (Å ²)	37.3	Xtriage
Anisotropy	0.621	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 67.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5446	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.32 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.5878e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/2663	1.27	10/3614 (0.3%)
1	B	0.79	0/2650	1.28	8/3591 (0.2%)
All	All	0.80	0/5313	1.28	18/7205 (0.2%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	316	GLY	CA-C-N	7.40	130.20	120.28
1	A	316	GLY	C-N-CA	7.40	130.20	120.28
1	B	227	ASP	CA-C-N	6.76	130.19	120.79
1	B	227	ASP	C-N-CA	6.76	130.19	120.79
1	A	201	GLU	N-CA-C	-5.89	105.69	114.64
1	A	316	GLY	N-CA-C	5.58	119.74	111.76
1	A	150	SER	N-CA-C	5.47	116.59	109.64
1	B	83	TYR	N-CA-C	-5.43	103.53	110.53
1	B	295	ASN	N-CA-C	5.40	117.32	108.52
1	B	298	ALA	CB-CA-C	-5.31	110.47	116.63
1	A	123	GLY	CA-C-N	5.23	128.51	120.82
1	A	123	GLY	C-N-CA	5.23	128.51	120.82
1	B	83	TYR	CA-C-O	-5.20	115.96	121.94
1	B	337	ALA	CA-C-N	5.16	127.65	120.63
1	B	337	ALA	C-N-CA	5.16	127.65	120.63
1	A	84	GLY	CA-C-N	5.16	127.80	120.53
1	A	84	GLY	C-N-CA	5.16	127.80	120.53
1	A	93	ILE	CB-CA-C	5.13	116.79	111.09

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2573	34	0
1	B	2593	0	2571	34	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	10	0	0	0	0
3	B	5	0	0	1	0
4	A	67	0	0	0	0
4	B	61	0	0	0	0
All	All	5446	0	5206	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ASN:O	1:A:300:CYS:HB2	1.66	0.95
1:B:41:GLY:O	1:B:44:THR:HG23	1.76	0.84
1:A:299:ASN:O	1:A:300:CYS:CB	2.30	0.80
1:A:300:CYS:N	1:A:301:PRO:HD3	1.99	0.78
1:A:297:ALA:HB1	1:A:300:CYS:H	1.49	0.78
1:A:189:SER:H	1:A:296:ALA:HB2	1.47	0.77
1:B:45:THR:HG22	1:B:112:SER:HB3	1.71	0.73
1:B:185:LEU:HD11	1:B:304:MET:HE1	1.71	0.72
1:B:45:THR:HG22	1:B:112:SER:CB	2.26	0.66
1:A:218:ASN:HD22	1:A:221:ILE:H	1.41	0.66
1:A:299:ASN:C	1:A:301:PRO:HD3	2.20	0.66
1:B:149:GLY:O	1:B:151:PRO:HD3	1.96	0.66
1:A:188:VAL:HA	1:A:296:ALA:HB1	1.76	0.65
1:B:218:ASN:HD22	1:B:221:ILE:H	1.43	0.65
1:B:220:CYS:HB2	1:B:269:GLN:HG2	1.77	0.65
1:A:35:LYS:HD2	1:A:131[B]:ARG:HD2	1.79	0.64
1:B:11:MET:HE1	1:B:114:ILE:HG21	1.78	0.64
1:A:220:CYS:HB2	1:A:269:GLN:HG2	1.80	0.64
1:B:335:LEU:HA	1:B:338:LEU:HD12	1.79	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:ALA:HB3	1:A:298:ALA:C	2.23	0.64
1:A:41:GLY:O	1:A:44:THR:HG23	1.98	0.63
1:A:293:VAL:CG1	1:A:296:ALA:HB3	2.29	0.63
1:A:188:VAL:HA	1:A:296:ALA:CB	2.30	0.61
1:A:300:CYS:N	1:A:301:PRO:CD	2.64	0.60
1:A:61:GLN:HG3	1:A:223:PHE:CE1	2.38	0.59
1:B:11:MET:HE3	1:B:324:PHE:HD2	1.69	0.58
1:B:11:MET:HE2	1:B:43:MET:SD	2.46	0.56
1:B:188:VAL:HG22	1:B:297:ALA:HB2	1.88	0.55
1:A:218:ASN:ND2	1:A:221:ILE:H	2.04	0.54
1:A:297:ALA:HB3	1:A:298:ALA:CA	2.37	0.54
1:B:218:ASN:ND2	1:B:221:ILE:H	2.05	0.53
1:B:188:VAL:CG2	1:B:297:ALA:HB2	2.40	0.52
1:B:45:THR:HG22	1:B:112:SER:OG	2.10	0.51
1:B:98:MET:HG2	1:B:102:ASP:HB2	1.92	0.51
1:B:21:ARG:HG3	1:B:123:GLY:H	1.76	0.50
1:A:40:GLY:HA2	1:A:115:ILE:HD13	1.93	0.50
1:A:178:SER:H	1:A:181:GLN:HE21	1.60	0.50
1:A:67:PRO:HG2	1:B:260:GLU:HG2	1.93	0.50
1:B:178:SER:H	1:B:181:GLN:HE21	1.60	0.49
1:B:40:GLY:HA2	1:B:115:ILE:HD13	1.94	0.49
1:A:297:ALA:HB3	1:A:298:ALA:HA	1.94	0.49
1:B:11:MET:HE1	1:B:114:ILE:CG2	2.43	0.48
1:B:200:TYR:HB2	1:B:240:PRO:HG2	1.96	0.48
1:A:200:TYR:HB2	1:A:240:PRO:HG2	1.95	0.47
1:A:188:VAL:HG13	1:A:296:ALA:HB1	1.96	0.46
1:B:218:ASN:ND2	1:B:220:CYS:H	2.14	0.46
1:A:218:ASN:ND2	1:A:220:CYS:H	2.15	0.45
1:B:63:ILE:HD11	1:B:111:ILE:HD11	2.00	0.44
1:B:218:ASN:HD21	1:B:269:GLN:HG3	1.83	0.43
1:A:223:PHE:CZ	1:A:225:SER:HB2	2.54	0.43
1:B:196:LEU:HD23	1:B:198:LEU:HD21	2.01	0.43
1:A:293:VAL:HG13	1:A:296:ALA:HB3	1.99	0.43
1:A:148:THR:HA	1:A:149:GLY:HA2	1.72	0.42
1:B:92:PRO:HG2	1:B:210:TRP:HA	2.00	0.42
1:A:297:ALA:HB1	1:A:300:CYS:N	2.27	0.42
1:B:149:GLY:O	1:B:151:PRO:CD	2.66	0.42
1:A:92:PRO:HG2	1:A:210:TRP:HA	2.01	0.41
1:A:48:SER:HB3	1:A:54:CYS:HB3	2.02	0.41
1:A:26:GLY:HA3	1:A:27:PRO:HA	1.95	0.41
1:B:193:ARG:HD2	1:B:292:GLN:HE22	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:PRO:HD2	1:B:165:GLU:OE1	2.20	0.41
1:B:273:ILE:HG22	1:B:274:LEU:HD23	2.03	0.41
1:A:297:ALA:CB	1:A:298:ALA:CA	2.98	0.41
1:B:193:ARG:NH2	3:B:343:SO4:O4	2.54	0.40
1:A:247:THR:HG23	1:A:250:PHE:H	1.86	0.40
1:B:11:MET:HE3	1:B:324:PHE:CD2	2.53	0.40
1:B:218:ASN:HD21	1:B:269:GLN:CG	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/342 (95%)	310 (95%)	15 (5%)	1 (0%)	36	55
1	B	320/342 (94%)	307 (96%)	12 (4%)	1 (0%)	36	55
All	All	646/684 (94%)	617 (96%)	27 (4%)	2 (0%)	36	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	300	CYS
1	B	339	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/293 (98%)	254 (88%)	33 (12%)	5	11
1	B	285/293 (97%)	252 (88%)	33 (12%)	5	11
All	All	572/586 (98%)	506 (88%)	66 (12%)	5	11

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LEU
1	A	47	CYS
1	A	50	HIS
1	A	55	THR
1	A	72	LYS
1	A	75	ARG
1	A	91	SER
1	A	93	ILE
1	A	111	ILE
1	A	120	LYS
1	A	126	VAL
1	A	130	HIS
1	A	139	ASP
1	A	161	MET
1	A	196	LEU
1	A	201	GLU
1	A	208	VAL
1	A	227	ASP
1	A	234	GLU
1	A	238	ILE
1	A	242	LEU
1	A	256	GLU
1	A	258	SER
1	A	265	LEU
1	A	292	GLN
1	A	295	ASN
1	A	300	CYS
1	A	305	THR
1	A	317	ASP
1	A	321	GLN
1	A	322	SER
1	A	329	THR
1	A	339	LYS
1	B	20	LEU
1	B	23	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	47	CYS
1	B	55	THR
1	B	75	ARG
1	B	79	GLU
1	B	87	ARG
1	B	91	SER
1	B	109	GLN
1	B	111	ILE
1	B	139	ASP
1	B	146	LYS
1	B	161	MET
1	B	170	GLN
1	B	173	ILE
1	B	196	LEU
1	B	204	THR
1	B	208	VAL
1	B	227	ASP
1	B	229	LYS
1	B	242	LEU
1	B	258	SER
1	B	265	LEU
1	B	267	PHE
1	B	269	GLN
1	B	292	GLN
1	B	303	GLN
1	B	305	THR
1	B	309	LYS
1	B	322	SER
1	B	323	ASN
1	B	329	THR
1	B	339	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	ASN
1	A	117	HIS
1	A	136	ASN
1	A	181	GLN
1	A	183	GLN
1	A	218	ASN
1	A	292	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	303	GLN
1	A	308	HIS
1	A	323	ASN
1	B	136	ASN
1	B	181	GLN
1	B	218	ASN
1	B	232	ASN
1	B	292	GLN
1	B	307	HIS
1	B	323	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	401	-	58,58,58	1.16	8 (13%)	85,89,89	1.29	13 (15%)
3	SO4	B	343	-	4,4,4	0.18	0	6,6,6	0.14	0
3	SO4	A	344	-	4,4,4	0.26	0	6,6,6	0.14	0
3	SO4	A	343	-	4,4,4	0.26	0	6,6,6	0.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	401	-	58,58,58	1.17	8 (13%)	85,89,89	1.29	10 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	401	-	-	7/34/50/50	0/6/6/6
2	FAD	B	401	-	-	7/34/50/50	0/6/6/6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	FAD	C10-N10	2.83	1.43	1.37
2	A	401	FAD	C9-C9A	2.75	1.44	1.39
2	A	401	FAD	C4X-C10	2.70	1.52	1.44
2	A	401	FAD	C4X-C4	2.58	1.54	1.44
2	B	401	FAD	C4X-C10	2.54	1.51	1.44
2	A	401	FAD	C2A-N3A	2.53	1.38	1.33
2	B	401	FAD	C4X-C4	2.47	1.53	1.44
2	B	401	FAD	C8-C7	2.44	1.46	1.40
2	A	401	FAD	C8-C7	2.38	1.46	1.40
2	B	401	FAD	P-O3P	-2.21	1.57	1.59
2	B	401	FAD	C5X-N5	2.20	1.43	1.39
2	A	401	FAD	C10-N10	2.08	1.41	1.37
2	B	401	FAD	C4A-N3A	2.07	1.38	1.34
2	A	401	FAD	C5'-C4'	2.05	1.54	1.51
2	B	401	FAD	C2A-N3A	2.02	1.37	1.33
2	A	401	FAD	C1'-C2'	-2.02	1.49	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FAD	C10-N1-C2	3.68	124.83	116.85
2	B	401	FAD	C10-N1-C2	3.63	124.70	116.85
2	B	401	FAD	C1'-C2'-C3'	-3.22	100.92	109.66
2	A	401	FAD	C1'-C2'-C3'	-2.95	101.67	109.66
2	A	401	FAD	C4-C4X-N5	2.85	122.15	118.21
2	B	401	FAD	C4-C4X-N5	2.69	121.93	118.21
2	A	401	FAD	O4'-C4'-C3'	-2.49	103.42	109.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	FAD	C4X-C10-N1	-2.47	118.53	124.59
2	B	401	FAD	O4'-C4'-C3'	-2.43	103.55	109.25
2	A	401	FAD	C4X-C10-N1	-2.38	118.76	124.59
2	A	401	FAD	N3A-C2A-N1A	-2.38	124.98	128.58
2	A	401	FAD	O2A-PA-O3P	-2.24	101.22	107.27
2	B	401	FAD	N10-C10-N1	2.23	125.06	118.51
2	A	401	FAD	C1'-N10-C9A	-2.19	116.38	120.63
2	B	401	FAD	N3A-C2A-N1A	-2.19	125.27	128.58
2	A	401	FAD	N10-C10-N1	2.17	124.90	118.51
2	A	401	FAD	O4B-C1B-C2B	-2.11	102.10	106.62
2	B	401	FAD	O4B-C1B-N9A	-2.10	104.05	108.09
2	A	401	FAD	O4B-C1B-N9A	-2.09	104.08	108.09
2	A	401	FAD	C5X-N5-C4X	2.08	121.46	118.09
2	B	401	FAD	C1'-N10-C9A	-2.08	116.59	120.63
2	A	401	FAD	O2A-PA-O1A	2.05	122.00	112.44
2	B	401	FAD	O2P-P-O3P	2.02	112.74	107.27

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	FAD	C5B-O5B-PA-O2A
2	A	401	FAD	C5'-O5'-P-O2P
2	A	401	FAD	PA-O3P-P-O5'
2	B	401	FAD	C5B-O5B-PA-O2A
2	B	401	FAD	PA-O3P-P-O5'
2	A	401	FAD	C2'-C1'-N10-C10
2	A	401	FAD	C5'-O5'-P-O3P
2	B	401	FAD	C2'-C1'-N10-C10
2	B	401	FAD	C5'-O5'-P-O2P
2	B	401	FAD	C5'-O5'-P-O3P
2	B	401	FAD	O4B-C4B-C5B-O5B
2	A	401	FAD	O4B-C4B-C5B-O5B
2	A	401	FAD	C3'-C4'-C5'-O5'
2	B	401	FAD	C3'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	343	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	335/342 (97%)	0.73	35 (10%)	11 10	20, 46, 79, 107	16 (4%)
1	B	333/342 (97%)	0.85	34 (10%)	12 10	18, 51, 84, 112	16 (4%)
All	All	668/684 (97%)	0.79	69 (10%)	12 10	18, 49, 82, 112	32 (4%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	341	TYR	12.9
1	B	123	GLY	9.1
1	A	1	MET	8.5
1	A	71	LYS	7.8
1	B	207	ASP	6.8
1	B	205	LYS	6.4
1	B	299	ASN	6.3
1	B	172	ASP	6.2
1	A	268	GLN	6.0
1	B	71	LYS	5.9
1	B	72	LYS	5.8
1	B	238	ILE	5.6
1	A	179	GLU	5.6
1	A	236	SER	5.4
1	A	23	GLN	5.3
1	A	300	CYS	5.3
1	B	203	GLY	5.2
1	A	259	ILE	5.2
1	A	204	THR	4.7
1	A	140	ASP	4.6
1	B	149	GLY	4.4
1	B	141	LYS	4.4
1	A	53	GLN	4.3
1	B	53	GLN	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	207	ASP	3.9
1	A	295	ASN	3.8
1	B	259	ILE	3.8
1	B	139	ASP	3.6
1	B	202	ALA	3.5
1	B	120	LYS	3.4
1	B	100	GLU	3.4
1	A	141	LYS	3.3
1	A	301	PRO	3.1
1	B	323	ASN	3.0
1	B	268	GLN	3.0
1	A	273	ILE	2.9
1	B	98	MET	2.9
1	A	202	ALA	2.9
1	B	310	PRO	2.7
1	A	24	THR	2.7
1	B	67	PRO	2.7
1	A	280	PRO	2.6
1	B	206	ILE	2.6
1	B	234	GLU	2.5
1	A	329	THR	2.5
1	B	179	GLU	2.5
1	A	208	VAL	2.5
1	A	98	MET	2.5
1	A	54	CYS	2.4
1	B	21	ARG	2.4
1	A	281	ILE	2.4
1	B	298	ALA	2.4
1	A	148	THR	2.3
1	A	341	TYR	2.3
1	A	238	ILE	2.3
1	B	208	VAL	2.3
1	A	150	SER	2.3
1	A	67	PRO	2.3
1	B	209	PRO	2.3
1	A	70	ALA	2.2
1	A	213	GLN	2.2
1	B	267	PHE	2.2
1	A	298	ALA	2.2
1	B	93	ILE	2.1
1	A	50	HIS	2.1
1	A	97	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	340	ASN	2.1
1	B	96	MET	2.1
1	A	297	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	343	5/5	0.83	0.12	83,88,88,88	0
2	FAD	A	401	53/53	0.96	0.08	22,25,29,31	0
3	SO4	B	343	5/5	0.96	0.07	42,46,47,47	0
3	SO4	A	344	5/5	0.97	0.07	53,57,58,58	0
2	FAD	B	401	53/53	0.97	0.08	30,31,36,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.