



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:51 AM UTC

PDB ID : 3QJ5 / pdb_00003qj5
Title : S-nitrosogluthione reductase (GSNOR) in complex with N6022
Authors : Chun, L.; Kim, H.; Sun, X.
Deposited on : 2011-01-28
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	93% 6%
1	B	374	93% 6%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6536 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase class-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	1	0
			2770	1764	464	520	22			
1	B	373	Total	C	N	O	S	0	4	0
			2793	1782	466	522	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P11766
B	0	SER	-	expression tag	UNP P11766

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		
3	B	1	Total	K	0	0
			1	1		

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
4	B	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 5 is 3-{1-(4-carbamoyl-2-methylphenyl)-5-[4-(1H-imidazol-1-yl)phenyl]-1H-pyrrol-2-yl}propanoic acid (CCD ID: 022) (formula: C₂₄H₂₂N₄O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			31	24	4	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			31	24	4	3		

- Molecule 6 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			16	10	6		
6	A	1	Total	C	O	0	0
			16	10	6		
6	B	1	Total	C	O	0	0
			16	10	6		

- Molecule 7 is PENTANE-1,5-DIAMINE (CCD ID: N2P) (formula: $C_5H_{14}N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	N	0	0
			7	5	2		
7	B	1	Total	C	N	0	0
			7	5	2		
7	B	1	Total	C	N	0	0
			7	5	2		

- Molecule 8 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	343	Total	O	0	0
			343	343		
9	B	397	Total	O	0	0
			397	397		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alcohol dehydrogenase class-3



- Molecule 1: Alcohol dehydrogenase class-3



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.86Å 78.86Å 310.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.09 – 1.90 49.09 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.09-1.90) 99.6 (49.09-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.184 , 0.210 0.183 , 0.209	Depositor DCC
R_{free} test set	3927 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6536	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 022, ZN, TRS, 1PE, NAD, N2P, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/2824	0.73	1/3818 (0.0%)
1	B	0.55	0/2856	0.72	0/3860
All	All	0.55	0/5680	0.73	1/7678 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	323	SER	N-CA-C	5.46	117.31	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2770	0	2824	15	0
1	B	2793	0	2864	30	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	44	0	26	2	0
4	B	44	0	26	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	31	0	21	1	0
5	B	31	0	21	6	0
6	A	32	0	44	0	0
6	B	16	0	22	1	0
7	A	7	0	14	3	0
7	B	14	0	28	4	0
8	A	8	0	12	2	0
9	A	343	0	0	1	0
9	B	397	0	0	3	0
All	All	6536	0	5902	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:HIS:HE1	7:B:379:N2P:H3C1	1.18	1.07
1:B:45:HIS:CE1	7:B:379:N2P:H3C1	1.96	0.99
1:A:45:HIS:HE1	7:A:379:N2P:H3C2	1.27	0.97
1:B:1:ALA:HA	1:B:74:SER:HB2	1.60	0.83
1:B:1:ALA:HB3	9:B:629:HOH:O	1.79	0.80
1:A:366:SER:HB2	9:A:686:HOH:O	1.81	0.80
1:B:1:ALA:CB	1:B:29:PRO:HD2	2.12	0.80
1:A:45:HIS:CE1	7:A:379:N2P:H3C2	2.16	0.78
1:B:1:ALA:HB1	1:B:29:PRO:HD2	1.69	0.73
1:B:1:ALA:HB2	1:B:29:PRO:HG2	1.79	0.63
1:A:49:TYR:CD1	7:A:379:N2P:H3C1	2.38	0.59
1:A:101:PHE:CE1	1:A:111:GLN:HG3	2.38	0.59
5:A:374:022:H27	1:B:309:THR:HG22	1.85	0.58
1:B:93:ILE:CD1	5:B:377:022:C64	2.82	0.57
1:A:309:THR:HG22	5:B:377:022:H27	1.89	0.54
1:A:281:CYS:HB3	1:A:286:GLY:HA3	1.89	0.54
1:B:2:ASN:ND2	1:B:3:GLU:H	2.07	0.53
1:B:111:GLN:OE1	1:B:114:ARG:CD	2.58	0.51
1:B:93:ILE:HD11	5:B:377:022:C64	2.41	0.51
1:B:281:CYS:HB3	1:B:286:GLY:HA3	1.93	0.51
1:B:93:ILE:HD13	5:B:377:022:C64	2.39	0.51
1:B:325:GLU:HB3	6:B:378:1PE:H231	1.92	0.51
1:A:67:GLU:OE1	1:A:173:CYS:HB3	2.10	0.51
1:B:2:ASN:HD22	1:B:3:GLU:H	1.59	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:MET:HA	1:A:122:MET:HE2	1.94	0.50
1:A:24:ILE:HG22	1:A:131:CYS:HB2	1.96	0.48
1:B:1:ALA:HB2	1:B:29:PRO:CG	2.43	0.48
1:B:49:TYR:CD1	7:B:379:N2P:H3C2	2.49	0.47
1:B:93:ILE:HD13	5:B:377:022:H64	1.95	0.47
1:A:48:ALA:HB2	1:A:361:MET:HE1	1.99	0.45
1:A:314:LYS:NZ	8:A:380:TRS:H11	2.31	0.45
1:B:1:ALA:HB2	1:B:29:PRO:HD2	1.95	0.44
1:B:45:HIS:HD2	9:B:677:HOH:O	1.98	0.44
1:B:268:ILE:HD12	1:B:273:VAL:HG11	2.00	0.44
1:B:1:ALA:HA	1:B:74:SER:CB	2.39	0.43
8:A:380:TRS:H12	1:B:314:LYS:HE2	2.00	0.43
1:B:93:ILE:CD1	5:B:377:022:H64B	2.48	0.43
1:B:111:GLN:OE1	1:B:114:ARG:HD2	2.19	0.43
1:B:49:TYR:CG	7:B:379:N2P:H3C2	2.53	0.43
1:A:177:THR:HG21	4:A:500:NAD:C4N	2.49	0.42
1:B:240:ILE:HD11	1:B:253[B]:LEU:HD11	2.01	0.42
1:B:291:VAL:O	4:B:374:NAD:H2N	2.20	0.42
1:A:291:VAL:O	4:A:500:NAD:H2N	2.20	0.42
1:A:195:ALA:HB2	1:A:261:VAL:HG11	2.03	0.41
1:B:67:GLU:CG	1:B:173:CYS:HB3	2.51	0.41
4:B:374:NAD:H6N	9:B:498:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/374 (99%)	362 (98%)	9 (2%)	0	100	100
1	B	375/374 (100%)	366 (98%)	8 (2%)	1 (0%)	36	29
All	All	746/748 (100%)	728 (98%)	17 (2%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/300 (100%)	298 (99%)	2 (1%)	76	78
1	B	303/300 (101%)	301 (99%)	2 (1%)	76	78
All	All	603/600 (100%)	599 (99%)	4 (1%)	76	78

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	CYS
1	A	188	LEU
1	B	188	LEU
1	B	313	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	HIS
1	A	184	ASN
1	A	243	GLN
1	A	306	GLN
1	B	2	ASN
1	B	45	HIS
1	B	306	GLN
1	B	362	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAD	A	500	-	46,48,48	1.71	6 (13%)	64,73,73	1.55	9 (14%)
6	1PE	A	377	-	15,15,15	0.47	0	14,14,14	0.27	0
5	022	B	377	2	34,34,34	2.03	5 (14%)	45,48,48	2.14	12 (26%)
7	N2P	B	380	-	6,6,6	0.19	0	5,5,5	0.41	0
4	NAD	B	374	-	46,48,48	1.66	7 (15%)	64,73,73	1.63	11 (17%)
6	1PE	A	378	-	15,15,15	0.47	0	14,14,14	0.30	0
7	N2P	A	379	-	6,6,6	0.20	0	5,5,5	0.39	0
6	1PE	B	378	-	15,15,15	0.46	0	14,14,14	0.30	0
7	N2P	B	379	-	6,6,6	0.23	0	5,5,5	0.43	0
5	022	A	374	2	34,34,34	2.00	5 (14%)	45,48,48	2.07	14 (31%)
8	TRS	A	380	-	7,7,7	0.31	0	9,9,9	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAD	A	500	-	-	4/30/62/62	0/5/5/5
6	1PE	A	377	-	-	8/13/13/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	022	B	377	2	-	5/21/21/21	0/4/4/4
7	N2P	B	380	-	-	3/4/4/4	-
4	NAD	B	374	-	-	4/30/62/62	0/5/5/5
6	1PE	A	378	-	-	2/13/13/13	-
7	N2P	A	379	-	-	4/4/4/4	-
6	1PE	B	378	-	-	6/13/13/13	-
7	N2P	B	379	-	-	2/4/4/4	-
5	022	A	374	2	-	3/21/21/21	0/4/4/4
8	TRS	A	380	-	-	3/9/9/9	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	500	NAD	O7N-C7N	8.84	1.40	1.24
4	B	374	NAD	O7N-C7N	8.17	1.39	1.24
5	A	374	022	C45-N29	-6.70	1.35	1.44
5	B	377	022	C45-N29	-6.66	1.35	1.44
5	A	374	022	C11-N2	-5.55	1.34	1.43
5	B	377	022	C11-N2	-5.33	1.34	1.43
5	B	377	022	C24-N29	-5.10	1.33	1.40
5	A	374	022	C24-N29	-4.50	1.34	1.40
5	B	377	022	C28-N29	-3.81	1.34	1.40
5	A	374	022	C28-N29	-3.61	1.34	1.40
4	A	500	NAD	C2A-N1A	2.98	1.39	1.33
4	B	374	NAD	C2A-N3A	2.90	1.39	1.33
4	B	374	NAD	C2A-N1A	2.83	1.39	1.33
5	A	374	022	C3-N2	-2.75	1.32	1.38
4	A	500	NAD	C2A-N3A	2.67	1.38	1.33
5	B	377	022	C3-N2	-2.51	1.33	1.38
4	B	374	NAD	PA-O3	2.45	1.62	1.59
4	A	500	NAD	PN-O3	2.15	1.61	1.59
4	A	500	NAD	C8A-N7A	2.13	1.35	1.31
4	B	374	NAD	C8A-N7A	2.08	1.35	1.31
4	B	374	NAD	C2N-N1N	2.05	1.37	1.35
4	A	500	NAD	C2N-N1N	2.04	1.37	1.35
4	B	374	NAD	PN-O3	2.01	1.61	1.59

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	374	NAD	N3A-C2A-N1A	-6.11	119.33	128.58
5	A	374	022	C3-N2-C1	6.02	115.33	106.40
5	B	377	022	C3-N2-C1	5.93	115.20	106.40
4	A	500	NAD	N3A-C2A-N1A	-5.91	119.63	128.58
5	B	377	022	C48-C47-C45	5.25	123.18	117.17
5	A	374	022	C48-C47-C45	4.99	122.88	117.17
5	B	377	022	C45-N29-C28	-4.76	118.65	124.57
4	B	374	NAD	C5A-C4A-N3A	-4.10	121.06	126.72
4	A	500	NAD	C5A-C4A-N3A	-4.09	121.09	126.72
4	A	500	NAD	N9A-C8A-N7A	-3.99	108.28	113.94
4	B	374	NAD	N9A-C8A-N7A	-3.97	108.30	113.94
5	B	377	022	C11-N2-C1	-3.63	121.26	127.20
4	B	374	NAD	C2A-N3A-C4A	3.41	120.16	111.83
4	B	374	NAD	C5A-N7A-C8A	3.37	108.74	103.45
4	A	500	NAD	C5A-N7A-C8A	3.31	108.66	103.45
4	A	500	NAD	C2A-N3A-C4A	3.28	119.83	111.83
5	B	377	022	C51-C45-C47	-3.27	116.14	121.16
5	A	374	022	C51-C45-C47	-3.23	116.20	121.16
5	A	374	022	C11-N2-C1	-3.13	122.08	127.20
4	B	374	NAD	C3N-C7N-N7N	3.06	121.51	117.74
5	A	374	022	C11-N2-C3	-2.91	122.58	126.84
4	A	500	NAD	N3A-C4A-N9A	2.88	132.07	127.17
5	B	377	022	C27-C28-N29	-2.87	105.69	107.48
4	B	374	NAD	N3A-C4A-N9A	2.86	132.03	127.17
5	A	374	022	N2-C1-N5	-2.85	103.81	111.13
5	A	374	022	C27-C28-N29	-2.77	105.75	107.48
5	A	374	022	C35-C28-C27	-2.74	125.61	130.13
5	B	377	022	N2-C1-N5	-2.67	104.28	111.13
5	B	377	022	C47-C45-N29	2.66	122.04	118.69
4	B	374	NAD	O7N-C7N-C3N	-2.55	116.47	119.60
5	A	374	022	C45-N29-C28	-2.53	121.42	124.57
5	B	377	022	C17-C11-N2	2.45	123.19	119.80
5	A	374	022	C49-C58-N62	2.41	120.71	117.74
5	B	377	022	C49-C58-N62	2.38	120.67	117.74
5	B	377	022	C35-C37-C38	-2.35	107.42	113.67
4	A	500	NAD	O2A-PA-O3	2.30	113.50	107.27
5	B	377	022	C11-N2-C3	-2.25	123.54	126.84
4	A	500	NAD	C4A-N9A-C8A	2.24	108.09	105.74
5	A	374	022	C14-C13-C11	2.23	123.13	120.30
5	A	374	022	C17-C11-N2	2.19	122.84	119.80
4	B	374	NAD	C4A-C5A-N7A	-2.16	108.11	110.58
5	A	374	022	C47-C45-N29	2.14	121.39	118.69
4	A	500	NAD	C4A-C5A-N7A	-2.13	108.14	110.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	374	NAD	C4A-N9A-C8A	2.09	107.94	105.74
4	B	374	NAD	C4D-O4D-C1D	-2.04	108.06	109.92
5	A	374	022	C15-C24-N29	2.00	128.03	124.15

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	500	NAD	O4D-C1D-N1N-C2N
4	A	500	NAD	O4D-C1D-N1N-C6N
4	A	500	NAD	C2D-C1D-N1N-C2N
4	B	374	NAD	O4D-C1D-N1N-C2N
4	B	374	NAD	O4D-C1D-N1N-C6N
4	B	374	NAD	C2D-C1D-N1N-C2N
4	B	374	NAD	C2D-C1D-N1N-C6N
6	A	377	1PE	OH6-C15-C25-OH5
6	A	377	1PE	OH2-C12-C22-OH3
6	A	378	1PE	OH2-C12-C22-OH3
6	A	377	1PE	OH7-C16-C26-OH6
6	B	378	1PE	OH7-C16-C26-OH6
7	B	380	N2P	C1-C2-C3-C4
5	B	377	022	C28-C35-C37-C38
5	B	377	022	C17-C11-N2-C3
5	B	377	022	C13-C11-N2-C3
6	A	377	1PE	C25-C15-OH6-C26
7	B	379	N2P	C2-C3-C4-C5
6	B	378	1PE	C13-C23-OH3-C22
6	A	377	1PE	C23-C13-OH4-C24
6	B	378	1PE	C12-C22-OH3-C23
6	A	377	1PE	C15-C25-OH5-C14
7	B	380	N2P	C2-C3-C4-C5
6	B	378	1PE	OH6-C15-C25-OH5
8	A	380	TRS	C1-C-C3-O3
5	B	377	022	C17-C11-N2-C1
6	A	377	1PE	C24-C14-OH5-C25
7	A	379	N2P	NE2-C1-C2-C3
7	B	379	N2P	C3-C4-C5-N1
5	B	377	022	C13-C11-N2-C1
7	A	379	N2P	C2-C3-C4-C5
6	A	377	1PE	OH5-C14-C24-OH4
4	A	500	NAD	C2D-C1D-N1N-C6N
6	A	378	1PE	C15-C25-OH5-C14

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
6	B	378	1PE	C14-C24-OH4-C13
7	A	379	N2P	C1-C2-C3-C4
8	A	380	TRS	C2-C-C3-O3
7	A	379	N2P	C3-C4-C5-N1
7	B	380	N2P	NE2-C1-C2-C3
5	A	374	022	C13-C11-N2-C3
5	A	374	022	C17-C11-N2-C3
5	A	374	022	C35-C37-C38-O43
8	A	380	TRS	N-C-C3-O3
6	B	378	1PE	C15-C25-OH5-C14



There are no ring outliers.

8 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	500	NAD	2	0
5	B	377	022	6	0
4	B	374	NAD	2	0
7	A	379	N2P	3	0
6	B	378	1PE	1	0
7	B	379	N2P	4	0
5	A	374	022	1	0
8	A	380	TRS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	372/374 (99%)	-0.01	5 (1%) 75 78	13, 21, 34, 44	1 (0%)
1	B	373/374 (99%)	-0.24	5 (1%) 75 78	9, 18, 27, 35	4 (1%)
All	All	745/748 (99%)	-0.12	10 (1%) 75 78	9, 19, 31, 44	5 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	8.4
1	A	57	GLU	3.1
1	A	56	PRO	2.8
1	B	362	HIS	2.8
1	A	362	HIS	2.8
1	B	2	ASN	2.5
1	A	23	GLU	2.5
1	B	246	SER	2.4
1	B	3	GLU	2.3
1	A	3	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	1PE	B	378	16/16	0.79	0.18	45,47,52,52	0
8	TRS	A	380	8/8	0.79	0.18	49,49,49,49	0
7	N2P	A	379	7/7	0.84	0.17	43,44,44,44	0
6	1PE	A	378	16/16	0.86	0.14	34,38,44,44	0
7	N2P	B	379	7/7	0.86	0.14	27,29,30,31	0
6	1PE	A	377	16/16	0.86	0.15	54,55,55,55	0
5	022	B	377	31/31	0.91	0.09	16,24,35,37	0
7	N2P	B	380	7/7	0.92	0.13	40,41,42,42	0
5	022	A	374	31/31	0.93	0.08	22,25,33,34	0
4	NAD	B	374	44/44	0.98	0.05	12,17,18,19	0
2	ZN	A	376	1/1	0.98	0.03	27,27,27,27	0
3	K	A	602	1/1	0.98	0.07	26,26,26,26	0
3	K	B	501	1/1	0.98	0.07	27,27,27,27	0
4	NAD	A	500	44/44	0.98	0.05	15,18,20,20	0
2	ZN	B	376	1/1	0.99	0.01	19,19,19,19	0
2	ZN	A	375	1/1	0.99	0.02	21,21,21,21	0
2	ZN	B	375	1/1	1.00	0.01	17,17,17,17	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 022 B 377:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 022 A 374:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD B 374:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.