



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:12 AM UTC

PDB ID : 5QJ5 / pdb_00005qj5
Title : PanDDA analysis group deposition – Crystal Structure of NUDT5 in complex with Z44592329
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	18% 81%10%7%
1	B	209	22% 82%10%7%
1	C	209	28% 78%12%8%
1	D	209	33% 81%10%9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	B	301	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1458	922	243	285	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1402	884	234	277	7			
1	D	190	Total	C	N	O	S	0	1	0
			1410	890	238	274	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is N-phenyl-N'-pyridin-3-ylurea (CCD ID: K0G) (formula: C₁₂H₁₁N₃O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 16 12 3 1	0	0
4	B	1	Total C N O 16 12 3 1	0	0
4	C	1	Total C N O 16 12 3 1	0	0
4	D	1	Total C N O 16 12 3 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	67	Total	O	0	0
			67	67		
6	B	73	Total	O	0	0
			73	73		
6	C	70	Total	O	0	0
			70	70		
6	D	39	Total	O	0	0
			39	39		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.15Å 59.77Å 80.09Å 79.47° 81.89° 75.78°	Depositor
Resolution (Å)	78.34 – 1.72 78.33 – 1.72	Depositor EDS
% Data completeness (in resolution range)	96.4 (78.34-1.72) 96.4 (78.33-1.72)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.215 , 0.253 0.242 , 0.272	Depositor DCC
R_{free} test set	4621 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6064	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K0G, CL, EDO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.37	6/1486 (0.4%)	1.12	0/2028
1	B	1.28	2/1492 (0.1%)	1.13	1/2035 (0.0%)
1	C	1.31	1/1429 (0.1%)	1.21	3/1958 (0.2%)
1	D	1.28	3/1436 (0.2%)	1.19	6/1961 (0.3%)
All	All	1.31	12/5843 (0.2%)	1.16	10/7982 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	198	TYR	C-O	-6.65	1.16	1.24
1	D	67	VAL	CA-C	6.62	1.60	1.52
1	B	120	LYS	CA-C	-6.41	1.45	1.52
1	A	124	ALA	CA-C	-6.20	1.46	1.52
1	B	127	SER	C-O	-5.80	1.16	1.23
1	D	175	LYS	CA-C	5.71	1.60	1.52
1	A	196	ARG	N-CA	5.65	1.53	1.46
1	A	75	GLU	N-CA	5.65	1.52	1.46
1	A	174	PRO	CA-C	-5.57	1.46	1.52
1	D	201	ALA	CA-C	5.40	1.59	1.52
1	A	127	SER	C-O	-5.33	1.17	1.23
1	A	141	ILE	CA-C	-5.26	1.46	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	85	PRO	N-CA-C	7.28	119.58	110.70
1	D	85	PRO	CA-C-N	-7.21	112.28	119.56
1	D	85	PRO	C-N-CA	-7.21	112.28	119.56
1	C	190	HIS	N-CA-CB	6.62	120.21	110.35
1	D	107	ALA	N-CA-C	-5.64	105.28	111.82
1	D	193	VAL	N-CA-C	-5.40	100.60	108.97
1	D	195	ALA	N-CA-C	5.27	117.24	111.50
1	C	206	HIS	N-CA-C	5.27	118.97	112.54
1	B	23	ILE	N-CA-C	-5.19	107.35	111.81
1	D	45	THR	N-CA-C	5.15	116.49	109.18

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	160	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1458	0	1420	18	0
1	B	1464	0	1414	16	0
1	C	1402	0	1298	19	0
1	D	1410	0	1349	12	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
4	A	16	0	0	0	0
4	B	16	0	0	0	0
4	C	16	0	0	0	0
4	D	16	0	0	0	0
5	B	4	0	6	4	0
5	C	4	0	6	0	0
6	A	67	0	0	1	0
6	B	73	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	70	0	0	4	0
6	D	39	0	0	1	0
All	All	6064	0	5493	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125:GLU:OE1	6:C:401:HOH:O	1.90	0.89
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.53	0.89
1:D:120:LYS:H	1:D:155:ASN:HD21	1.27	0.82
1:A:104:THR:HG23	1:A:107:ALA:H	1.46	0.79
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.69	0.74
5:B:301:EDO:H22	6:B:409:HOH:O	1.88	0.73
1:B:196:ARG:NH2	6:B:401:HOH:O	2.23	0.70
1:A:92:ILE:HD11	1:A:191:LEU:HD11	1.72	0.70
5:B:301:EDO:C2	6:B:409:HOH:O	2.41	0.68
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.80	0.63
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.33	0.62
1:A:179:LEU:HD23	1:A:205:LYS:HD2	1.83	0.60
1:C:104:THR:HG23	1:C:107:ALA:H	1.67	0.60
6:C:401:HOH:O	1:D:206:HIS:HE1	1.86	0.58
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.40	0.56
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.40	0.56
1:B:104:THR:HG23	1:B:107:ALA:H	1.70	0.56
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.90	0.55
1:C:132:MET:O	1:D:196[B]:ARG:NH1	2.41	0.54
5:B:301:EDO:H21	6:B:452:HOH:O	2.09	0.53
1:A:71:THR:HG23	1:A:151:ASP:OD2	2.08	0.53
1:A:206:HIS:HE1	1:B:125:GLU:OE1	1.92	0.53
1:D:112:GLU:OE2	6:D:401:HOH:O	2.19	0.51
1:C:200:TYR:OH	1:D:206:HIS:HD2	1.94	0.50
1:C:62:VAL:HG12	1:C:142:HIS:HB2	1.93	0.49
1:C:191:LEU:HD12	1:C:191:LEU:C	2.37	0.49
1:B:62:VAL:HG12	1:B:142:HIS:HB2	1.94	0.49
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.96	0.49
1:C:206:HIS:HE1	1:D:125:GLU:OE1	1.96	0.49
1:C:125:GLU:HG2	6:C:405:HOH:O	2.12	0.48
5:B:301:EDO:C2	6:B:452:HOH:O	2.61	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:VAL:HA	1:B:142:HIS:O	2.16	0.46
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.51	0.45
1:C:92:ILE:HD12	1:C:191:LEU:HD11	1.99	0.45
1:A:203:ALA:HB2	1:B:200:TYR:CD1	2.52	0.45
1:A:38:ASP:OD1	1:A:38:ASP:C	2.60	0.45
1:D:17:ILE:HD12	1:D:34:THR:CG2	2.48	0.44
1:A:92:ILE:HD12	1:A:191:LEU:HD21	1.99	0.44
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.48	0.43
1:C:206:HIS:HD2	1:D:200:TYR:OH	2.01	0.43
1:A:92:ILE:CD1	1:A:191:LEU:HD11	2.46	0.43
1:B:38:ASP:OD1	1:B:38:ASP:C	2.62	0.43
1:A:70:ARG:HD3	6:A:433:HOH:O	2.18	0.42
1:B:189:GLU:OE2	1:B:191:LEU:HD11	2.18	0.42
1:A:200:TYR:OH	1:B:206:HIS:HD2	2.00	0.42
1:A:191:LEU:HD23	1:A:192:THR:N	2.34	0.42
1:C:31:LEU:HD23	1:C:32:GLU:H	1.85	0.41
1:C:165:GLY:HA2	1:C:167:PHE:CE1	2.55	0.41
1:C:92:ILE:HD12	1:C:191:LEU:CD1	2.51	0.41
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.51	0.41
1:B:158:PRO:O	1:B:160:PRO:HD3	2.21	0.41
1:A:71:THR:O	1:A:73:HIS:HD2	2.04	0.40
1:D:38:ASP:OD1	1:D:38:ASP:C	2.63	0.40
1:B:92:ILE:CD1	1:B:191:LEU:HD13	2.37	0.40
1:C:74:TYR:HB3	6:C:424:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	192/209 (92%)	186 (97%)	5 (3%)	1 (0%)	24 11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	190/209 (91%)	185 (97%)	4 (2%)	1 (0%)	24	11
1	C	188/209 (90%)	181 (96%)	5 (3%)	2 (1%)	11	3
1	D	187/209 (90%)	180 (96%)	7 (4%)	0	100	100
All	All	757/836 (91%)	732 (97%)	21 (3%)	4 (0%)	24	11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	53	THR
1	A	66	PRO
1	C	66	PRO
1	B	66	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/179 (86%)	148 (97%)	5 (3%)	33	10
1	B	154/179 (86%)	150 (97%)	4 (3%)	40	18
1	C	139/179 (78%)	134 (96%)	5 (4%)	31	8
1	D	144/179 (80%)	142 (99%)	2 (1%)	59	40
All	All	590/716 (82%)	574 (97%)	16 (3%)	39	17

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	38	ASP
1	A	70	ARG
1	A	125	GLU
1	A	191	LEU
1	B	20	GLU
1	B	31	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	38	ASP
1	B	208	ASN
1	C	29	VAL
1	C	35	THR
1	C	51	ARG
1	C	71	THR
1	C	189	GLU
1	D	45	THR
1	D	72	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	73	HIS
1	A	180	GLN
1	A	206	HIS
1	B	15	GLN
1	B	73	HIS
1	B	206	HIS
1	C	138	ASN
1	C	206	HIS
1	D	15	GLN
1	D	149	ASN
1	D	155	ASN
1	D	206	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 15 ligands modelled in this entry, 9 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	K0G	A	304	-	17,17,17	1.18	2 (11%)	21,21,21	0.96	1 (4%)
5	EDO	C	301	-	3,3,3	0.78	0	2,2,2	0.07	0
4	K0G	B	304	-	17,17,17	1.27	1 (5%)	21,21,21	0.60	0
4	K0G	D	303	-	17,17,17	1.44	3 (17%)	21,21,21	0.99	0
5	EDO	B	301	-	3,3,3	0.60	0	2,2,2	1.05	0
4	K0G	C	304	-	17,17,17	1.13	1 (5%)	21,21,21	1.09	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	K0G	A	304	-	-	1/8/8/8	0/2/2/2
5	EDO	C	301	-	-	1/1/1/1	-
4	K0G	B	304	-	-	2/8/8/8	0/2/2/2
4	K0G	D	303	-	-	1/8/8/8	0/2/2/2
5	EDO	B	301	-	-	1/1/1/1	-
4	K0G	C	304	-	-	2/8/8/8	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	303	K0G	C2-C1	2.73	1.43	1.39
4	B	304	K0G	C2-C1	2.70	1.43	1.39
4	D	303	K0G	C-N1	2.42	1.42	1.37
4	C	304	K0G	C8-C7	2.37	1.43	1.39
4	A	304	K0G	C8-C7	2.25	1.43	1.39
4	A	304	K0G	C6-C1	2.22	1.43	1.39
4	D	303	K0G	C5-C6	2.15	1.42	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	304	K0G	C7-N1-C	-3.36	119.76	126.61
4	A	304	K0G	C7-N1-C	-2.17	122.19	126.61

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	304	K0G	C11-C7-N1-C
4	B	304	K0G	C11-C7-N1-C
4	D	303	K0G	C11-C7-N1-C
5	B	301	EDO	O1-C1-C2-O2
4	A	304	K0G	C11-C7-N1-C
4	C	304	K0G	C8-C7-N1-C
5	C	301	EDO	O1-C1-C2-O2
4	B	304	K0G	C8-C7-N1-C

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	301	EDO	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	1.38	37 (19%) 3 5	12, 30, 55, 67	26 (13%)
1	B	194/209 (92%)	1.57	46 (23%) 2 4	13, 35, 62, 75	21 (10%)
1	C	192/209 (91%)	1.63	58 (30%) 1 2	21, 37, 67, 93	21 (10%)
1	D	190/209 (90%)	2.06	69 (36%) 1 1	10, 40, 69, 78	26 (13%)
All	All	770/836 (92%)	1.66	210 (27%) 1 3	10, 36, 65, 93	94 (12%)

All (210) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	53	THR	14.3
1	B	52	THR	11.5
1	D	29	VAL	10.8
1	D	31	LEU	10.4
1	A	28	TRP	9.9
1	D	153	ALA	9.8
1	A	191	LEU	9.1
1	B	62	VAL	8.8
1	C	23	ILE	8.8
1	D	74	TYR	8.7
1	B	55	LYS	8.7
1	B	54	ARG	8.7
1	B	59	ALA	8.7
1	A	59	ALA	8.6
1	C	22	LEU	8.6
1	B	63	ALA	8.6
1	D	28	TRP	8.3
1	B	135	GLY	8.2
1	D	53	THR	8.0
1	A	54	ARG	7.9
1	A	53	THR	7.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	LYS	7.7
1	B	50	LYS	7.5
1	A	57	GLN	7.5
1	C	52	THR	7.5
1	A	139	CYS	7.4
1	A	51	ARG	7.3
1	B	49	VAL	7.1
1	B	134	PRO	7.1
1	A	52	THR	7.1
1	A	136	LEU	6.9
1	C	49	VAL	6.9
1	D	49	VAL	6.8
1	A	135	GLY	6.8
1	C	26	GLY	6.8
1	A	190	HIS	6.8
1	B	208	ASN	6.6
1	B	61	GLY	6.6
1	A	27	LYS	6.5
1	D	73	HIS	6.5
1	A	29	VAL	6.5
1	A	58	THR	6.4
1	C	137	SER	6.4
1	B	51	ARG	6.3
1	D	134	PRO	6.3
1	A	61	GLY	6.3
1	B	60	ASP	6.3
1	C	24	SER	6.1
1	C	139	CYS	6.1
1	D	136	LEU	6.1
1	B	136	LEU	6.0
1	A	50	LYS	6.0
1	D	30	LYS	6.0
1	D	52	THR	6.0
1	A	56	GLU	6.0
1	C	25	GLU	6.0
1	B	133	ASP	5.9
1	B	139	CYS	5.9
1	D	51	ARG	5.9
1	D	196[A]	ARG	5.9
1	C	141	ILE	5.9
1	A	26	GLY	5.9
1	C	50	LYS	5.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	28	TRP	5.8
1	C	136	LEU	5.8
1	C	51	ARG	5.8
1	D	154	GLU	5.5
1	D	139	CYS	5.4
1	C	138	ASN	5.3
1	B	132	MET	5.3
1	D	50	LYS	5.2
1	A	189	GLU	5.2
1	D	99	ILE	5.2
1	A	137	SER	5.2
1	D	135	GLY	5.1
1	B	28	TRP	5.0
1	D	23	ILE	4.9
1	B	137	SER	4.9
1	C	140	THR	4.9
1	A	138	ASN	4.7
1	A	188	GLU	4.7
1	D	16	TYR	4.7
1	B	196	ARG	4.7
1	B	58	THR	4.7
1	C	53	THR	4.6
1	D	48	SER	4.5
1	C	55	LYS	4.5
1	D	137	SER	4.4
1	C	54	ARG	4.4
1	C	16	TYR	4.4
1	C	165	GLY	4.4
1	A	72	LEU	4.3
1	C	58	THR	4.2
1	A	60	ASP	4.2
1	D	97	GLY	4.1
1	C	40	THR	4.1
1	C	60	ASP	4.1
1	C	142	HIS	4.1
1	D	32	GLU	4.0
1	D	131	CYS	3.9
1	D	167	PHE	3.9
1	C	41	GLY	3.9
1	D	43	THR	3.9
1	A	33	LYS	3.8
1	D	34	THR	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	152	ASP	3.7
1	D	138	ASN	3.7
1	D	47	GLU	3.7
1	C	59	ALA	3.6
1	D	18	ILE	3.6
1	B	165	GLY	3.5
1	A	92	ILE	3.5
1	D	88	GLY	3.5
1	C	57	GLN	3.4
1	C	39	PRO	3.4
1	D	162	PRO	3.4
1	C	153	ALA	3.4
1	C	43	THR	3.3
1	D	90	TYR	3.3
1	D	158	PRO	3.3
1	C	29	VAL	3.3
1	D	100	ASP	3.2
1	B	40	THR	3.2
1	D	133	ASP	3.1
1	A	16	TYR	3.1
1	D	40	THR	3.1
1	B	167	PHE	3.0
1	D	36	TYR	3.0
1	D	35	THR	3.0
1	B	18	ILE	3.0
1	B	191	LEU	3.0
1	C	37	MET	3.0
1	C	184	ALA	2.9
1	A	167	PHE	2.9
1	A	14	LYS	2.9
1	C	48	SER	2.9
1	B	39	PRO	2.9
1	B	163	GLY	2.8
1	D	59	ALA	2.8
1	D	160	PRO	2.8
1	C	35	THR	2.8
1	D	14	LYS	2.8
1	B	14	LYS	2.8
1	C	34	THR	2.8
1	C	17	ILE	2.7
1	D	161	LYS	2.7
1	C	190	HIS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	98	LEU	2.7
1	B	104	THR	2.6
1	D	132	MET	2.6
1	B	160	PRO	2.6
1	D	41	GLY	2.6
1	C	74	TYR	2.6
1	C	45	THR	2.6
1	B	151	ASP	2.6
1	D	78	VAL	2.6
1	C	18	ILE	2.6
1	C	99	ILE	2.6
1	D	155	ASN	2.5
1	D	102	GLY	2.5
1	B	102	GLY	2.4
1	D	163	GLY	2.4
1	D	164	ASP	2.4
1	C	150	GLY	2.4
1	B	99	ILE	2.4
1	B	164	ASP	2.4
1	C	102	GLY	2.4
1	A	168	VAL	2.4
1	C	104	THR	2.3
1	D	83	PHE	2.3
1	D	87	MET	2.3
1	C	46	TRP	2.3
1	D	119	TYR	2.3
1	A	40	THR	2.3
1	C	98	LEU	2.3
1	A	157	ARG	2.3
1	D	70	ARG	2.3
1	C	152	ASP	2.3
1	A	22	LEU	2.3
1	B	23	ILE	2.3
1	B	190	HIS	2.2
1	C	15	GLN	2.2
1	D	22	LEU	2.2
1	C	101	ASP	2.2
1	B	150	GLY	2.2
1	C	83	PHE	2.2
1	D	46	TRP	2.2
1	D	208	ASN	2.2
1	C	31	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	17	ILE	2.2
1	D	39	PRO	2.2
1	C	19	SER	2.2
1	B	108	ALA	2.2
1	B	156	ALA	2.2
1	D	166	GLU	2.2
1	C	185	LEU	2.2
1	C	162	PRO	2.2
1	D	151	ASP	2.1
1	B	186	VAL	2.1
1	D	113	LEU	2.1
1	C	100	ASP	2.1
1	C	168	VAL	2.1
1	D	96	ALA	2.1
1	D	42	LYS	2.1
1	D	101	ASP	2.1
1	A	41	GLY	2.1
1	B	24	SER	2.1
1	B	22	LEU	2.0
1	D	123	ILE	2.0
1	B	153	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	K0G	A	304	16/16	0.65	0.36	32,39,44,44	16
4	K0G	B	304	16/16	0.70	0.28	33,36,40,41	16

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	K0G	D	303	16/16	0.73	0.28	39,48,52,53	16
5	EDO	C	301	4/4	0.75	0.17	39,52,52,58	0
4	K0G	C	304	16/16	0.80	0.22	35,42,49,50	16
5	EDO	B	301	4/4	0.84	0.13	38,40,41,42	0
2	MG	D	301	1/1	0.92	0.07	43,43,43,43	0
3	CL	A	303	1/1	0.92	0.16	70,70,70,70	0
2	MG	C	302	1/1	0.95	0.05	62,62,62,62	0
2	MG	B	303	1/1	0.96	0.08	58,58,58,58	0
2	MG	A	302	1/1	0.97	0.04	52,52,52,52	0
2	MG	D	302	1/1	0.97	0.11	41,41,41,41	0
2	MG	B	302	1/1	0.97	0.07	35,35,35,35	0
2	MG	C	303	1/1	0.98	0.05	42,42,42,42	0
2	MG	A	301	1/1	0.99	0.05	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.