



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:02 AM UTC

PDB ID : 5QJ8 / pdb_00005qj8
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z2856434829
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	11% 82% 10% 7%
1	B	209	10% 78% 14% 7%
1	C	209	28% 74% 15% 8%
1	D	209	47% 70% 19% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5997 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1462	925	244	285	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1402	884	234	277	7			
1	D	190	Total	C	N	O	S	0	1	0
			1404	887	235	274	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0

- Molecule 5 is [3,4-bis(fluoranyl)phenyl]-(4-methylpiperazin-1-yl)methanone (CCD ID: ELQ) (formula: $C_{12}H_{14}F_2N_2O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	F	N	O	0	0
			17	12	2	2	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	71	Total	O	0	0
			71	71		
6	B	71	Total	O	0	0
			71	71		
6	C	62	Total	O	0	0
			62	62		
6	D	27	Total	O	0	0
			27	27		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.53Å 59.99Å 79.65Å 79.56° 81.09° 75.49°	Depositor
Resolution (Å)	77.81 – 1.76 77.81 – 1.76	Depositor EDS
% Data completeness (in resolution range)	96.8 (77.81-1.76) 96.8 (77.81-1.76)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.234 , 0.278 (Not available) , 0.301	Depositor DCC
R_{free} test set	4344 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5997	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG, ELQ, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	2/1490 (0.1%)	1.13	0/2032
1	B	1.22	2/1492 (0.1%)	1.17	6/2035 (0.3%)
1	C	1.29	3/1429 (0.2%)	1.25	5/1958 (0.3%)
1	D	1.27	4/1430 (0.3%)	1.22	5/1954 (0.3%)
All	All	1.26	11/5841 (0.2%)	1.19	16/7979 (0.2%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	78	VAL	C-O	-6.98	1.16	1.24
1	D	84	ARG	C-O	-6.13	1.19	1.25
1	A	100	ASP	N-CA	5.80	1.53	1.45
1	D	197	VAL	C-O	-5.67	1.17	1.24
1	C	81	LYS	N-CA	-5.45	1.39	1.46
1	C	77	ILE	CA-CB	-5.43	1.48	1.54
1	A	204	LEU	CA-C	5.36	1.59	1.52
1	D	173	LEU	CA-CB	-5.27	1.46	1.53
1	C	104	THR	CA-C	5.19	1.59	1.53
1	B	63	ALA	C-O	-5.06	1.17	1.23
1	D	201	ALA	CA-C	5.00	1.59	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	38	ASP	CA-C-N	9.61	129.84	119.28
1	C	38	ASP	C-N-CA	9.61	129.84	119.28
1	D	94	PHE	N-CA-C	-7.10	100.82	109.83
1	C	85	PRO	N-CA-C	6.99	119.23	110.70
1	D	85	PRO	N-CA-C	6.89	119.11	110.70
1	D	193	VAL	N-CA-C	-5.86	99.88	108.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	83	PHE	N-CA-C	-5.57	100.90	109.76
1	D	176	ASN	N-CA-C	5.42	118.99	112.38
1	C	132	MET	N-CA-C	5.31	117.88	111.40
1	B	23	ILE	N-CA-C	-5.24	107.31	111.81
1	D	138	ASN	N-CA-C	-5.18	106.64	113.17
1	B	65	ILE	CA-C-N	-5.10	113.46	119.84
1	B	65	ILE	C-N-CA	-5.10	113.46	119.84
1	C	140	THR	N-CA-C	5.08	117.53	109.50
1	B	94	PHE	CA-C-N	-5.03	114.77	119.90
1	B	94	PHE	C-N-CA	-5.03	114.77	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1462	0	1431	18	0
1	B	1464	0	1414	22	0
1	C	1402	0	1298	22	0
1	D	1404	0	1338	29	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	B	1	0	0	0	0
4	B	4	0	6	1	0
4	C	4	0	6	0	0
5	B	17	0	0	0	0
6	A	71	0	0	1	0
6	B	71	0	0	1	0
6	C	62	0	0	4	0
6	D	27	0	0	0	0
All	All	5997	0	5493	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:CG	1.50	1.21
1:D:196[B]:ARG:HG3	1:D:196[B]:ARG:NH2	1.48	1.00
1:C:125:GLU:OE1	6:C:401:HOH:O	1.87	0.91
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.62	0.80
1:C:104:THR:HG23	1:C:107:ALA:H	1.52	0.75
1:A:175:LYS:NZ	1:B:206:HIS:O	2.20	0.72
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:N	2.38	0.71
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:HG3	0.64	0.70
1:D:120:LYS:H	1:D:155:ASN:HD21	1.40	0.69
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.30	0.67
1:D:196[B]:ARG:CG	1:D:196[B]:ARG:NH2	2.23	0.64
1:B:104:THR:HG23	1:B:107:ALA:H	1.64	0.63
1:A:104:THR:HG23	1:A:107:ALA:H	1.64	0.62
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.80	0.62
1:C:191:LEU:HD12	1:C:191:LEU:C	2.26	0.60
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.83	0.60
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.32	0.59
1:A:70:ARG:NH1	6:A:401:HOH:O	2.36	0.59
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:H	2.02	0.57
1:D:35:THR:HG23	1:D:43:THR:HG23	1.86	0.57
1:C:38:ASP:HB2	1:D:167:PHE:CE2	2.41	0.56
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.41	0.56
1:B:17:ILE:HD12	1:B:34:THR:CG2	2.38	0.54
1:D:81:LYS:HA	1:D:91:CYS:O	2.07	0.53
1:D:196[B]:ARG:H	1:D:196[B]:ARG:CZ	2.22	0.53
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.91	0.53
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.44	0.52
1:A:104:THR:CG2	1:A:107:ALA:H	2.22	0.52
1:D:195:ALA:HB3	1:D:196[B]:ARG:HH12	1.75	0.51
1:C:20:GLU:HA	1:C:33:LYS:O	2.10	0.51
1:D:177:ASP:OD1	1:D:180:GLN:CB	2.58	0.51
6:C:401:HOH:O	1:D:206:HIS:HE1	1.92	0.51
1:C:200:TYR:OH	1:D:206:HIS:HD2	1.94	0.50
1:D:38:ASP:OD1	1:D:40:THR:N	2.39	0.49
1:B:189:GLU:HB3	1:B:191:LEU:HD12	1.95	0.48
1:C:206:HIS:HD2	1:D:200:TYR:OH	1.95	0.48
1:C:60:ASP:OD2	1:C:140:THR:N	2.30	0.48
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.48	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:15:GLN:N	6:C:405:HOH:O	2.46	0.47
1:A:49:VAL:HG21	1:B:49:VAL:HG11	1.97	0.47
1:A:203:ALA:HB2	1:B:200:TYR:CD1	2.50	0.46
1:C:92:ILE:HG13	1:C:191:LEU:HD13	1.98	0.46
1:B:92:ILE:CD1	1:B:191:LEU:HD13	2.39	0.46
1:B:158:PRO:O	1:B:160:PRO:HD3	2.16	0.46
1:C:138:ASN:HB3	1:D:86:PRO:O	2.16	0.45
1:D:121:GLY:HA3	1:D:146:VAL:CG1	2.47	0.45
1:A:104:THR:HG22	1:A:107:ALA:CB	2.47	0.45
1:C:131:CYS:SG	1:D:194:ASP:HA	2.57	0.45
1:B:38:ASP:OD1	1:B:38:ASP:C	2.60	0.44
1:C:74:TYR:HB3	6:C:426:HOH:O	2.18	0.44
1:D:182:LEU:O	1:D:186:VAL:HG23	2.17	0.44
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.99	0.44
1:B:17:ILE:HD12	1:B:34:THR:HG21	1.97	0.44
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.47	0.44
1:D:104:THR:HG23	1:D:107:ALA:H	1.82	0.44
1:A:200:TYR:OH	1:B:206:HIS:HD2	2.00	0.43
1:B:94:PHE:O	1:B:95:PRO:C	2.60	0.43
1:A:86:PRO:HD3	1:B:46:TRP:CD1	2.54	0.43
1:A:138:ASN:HB3	1:B:86:PRO:O	2.18	0.42
1:C:127:SER:HB2	1:C:128:PRO:HD2	2.01	0.42
1:A:39:PRO:HG2	1:B:167:PHE:CD2	2.54	0.42
4:B:302:EDO:C2	6:B:411:HOH:O	2.68	0.42
1:A:167:PHE:CZ	1:B:39:PRO:HD2	2.55	0.42
1:B:58:THR:CB	1:B:142:HIS:NE2	2.83	0.42
1:D:64:VAL:O	1:D:94:PHE:HB3	2.19	0.42
1:D:97:GLY:HA3	1:D:112:GLU:CD	2.45	0.41
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:CA	2.83	0.41
1:C:21:GLU:O	1:C:32:GLU:HA	2.20	0.41
1:A:70:ARG:HB3	1:A:73:HIS:HB2	2.02	0.41
1:C:87:MET:HE2	1:C:192:THR:HG21	2.03	0.41
1:C:132:MET:O	1:D:196[B]:ARG:NH1	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	187 (97%)	4 (2%)	1 (0%)	24	11
1	B	190/209 (91%)	183 (96%)	6 (3%)	1 (0%)	24	11
1	C	188/209 (90%)	176 (94%)	9 (5%)	3 (2%)	7	1
1	D	187/209 (90%)	174 (93%)	13 (7%)	0	100	100
All	All	757/836 (91%)	720 (95%)	32 (4%)	5 (1%)	18	7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	24	SER
1	C	22	LEU
1	A	66	PRO
1	B	66	PRO
1	C	66	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/179 (86%)	150 (97%)	4 (3%)	40	20
1	B	154/179 (86%)	149 (97%)	5 (3%)	34	14
1	C	139/179 (78%)	133 (96%)	6 (4%)	26	8
1	D	143/179 (80%)	132 (92%)	11 (8%)	12	2
All	All	590/716 (82%)	564 (96%)	26 (4%)	25	7

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	33	LYS
1	A	70	ARG
1	A	191	LEU
1	B	14	LYS
1	B	31	LEU
1	B	43	THR
1	B	189	GLU
1	B	205	LYS
1	C	22	LEU
1	C	29	VAL
1	C	31	LEU
1	C	35	THR
1	C	164	ASP
1	C	191	LEU
1	D	19	SER
1	D	24	SER
1	D	38	ASP
1	D	42	LYS
1	D	43	THR
1	D	48	SER
1	D	72	LEU
1	D	76	CYS
1	D	112	GLU
1	D	117	THR
1	D	139	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	206	HIS
1	B	206	HIS
1	C	73	HIS
1	C	206	HIS
1	D	82	GLN
1	D	155	ASN
1	D	206	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ELQ	B	305	-	18,18,18	2.03	5 (27%)	25,25,25	1.46	4 (16%)
4	EDO	C	301	-	3,3,3	0.62	0	2,2,2	0.13	0
4	EDO	B	302	-	3,3,3	0.47	0	2,2,2	0.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ELQ	B	305	-	-	0/8/18/18	1/2/2/2
4	EDO	C	301	-	-	0/1/1/1	-
4	EDO	B	302	-	-	1/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	305	ELQ	C08-N05	5.07	1.45	1.34
5	B	305	ELQ	C04-N05	2.88	1.52	1.47
5	B	305	ELQ	C07-N02	2.76	1.53	1.46
5	B	305	ELQ	C03-N02	2.29	1.51	1.46
5	B	305	ELQ	C11-C10	2.06	1.42	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	305	ELQ	C10-C08-N05	4.71	124.54	118.66
5	B	305	ELQ	C07-N02-C03	2.38	113.33	109.54
5	B	305	ELQ	O09-C08-C10	-2.35	115.63	120.29
5	B	305	ELQ	C15-C10-C08	-2.23	115.18	120.13

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	302	EDO	O1-C1-C2-O2

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	305	ELQ	C03-C04-C06-C07-N02-N05

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	302	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	0.83	23 (11%)	9 10	16, 31, 51, 64	13 (6%)
1	B	194/209 (92%)	0.85	20 (10%)	12 13	23, 36, 61, 76	3 (1%)
1	C	192/209 (91%)	1.51	59 (30%)	1 1	21, 37, 67, 79	7 (3%)
1	D	190/209 (90%)	2.54	98 (51%)	0 0	9, 45, 67, 81	17 (8%)
All	All	770/836 (92%)	1.43	200 (25%)	1 2	9, 37, 64, 81	40 (5%)

All (200) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	28	TRP	15.6
1	D	28	TRP	13.6
1	C	23	ILE	13.2
1	D	196[A]	ARG	12.7
1	D	160	PRO	11.7
1	C	26	GLY	11.6
1	D	191	LEU	11.3
1	C	22	LEU	11.0
1	D	26	GLY	11.0
1	D	156	ALA	10.3
1	C	25	GLU	10.0
1	B	190	HIS	9.9
1	A	191	LEU	9.8
1	A	187	ALA	9.6
1	D	158	PRO	9.3
1	C	27	LYS	9.2
1	D	27	LYS	9.2
1	C	24	SER	8.7
1	D	159	LYS	7.9
1	A	190	HIS	7.6
1	D	208	ASN	7.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	157	ARG	7.4
1	A	188	GLU	7.2
1	B	189	GLU	7.0
1	B	188	GLU	6.6
1	A	28	TRP	6.4
1	D	153	ALA	6.3
1	A	33	LYS	6.2
1	D	190	HIS	6.0
1	D	155	ASN	5.8
1	D	90	TYR	5.8
1	C	18	ILE	5.7
1	A	189	GLU	5.7
1	D	188	GLU	5.7
1	B	28	TRP	5.0
1	D	154	GLU	4.9
1	D	168	VAL	4.9
1	D	99	ILE	4.8
1	D	151	ASP	4.8
1	D	167	PHE	4.8
1	C	39	PRO	4.8
1	D	74	TYR	4.7
1	D	189	GLU	4.7
1	D	97	GLY	4.6
1	D	31	LEU	4.5
1	C	16	TYR	4.4
1	D	119	TYR	4.2
1	A	57	GLN	4.2
1	A	56	GLU	4.2
1	D	29	VAL	4.1
1	D	23	ILE	4.1
1	A	27	LYS	4.1
1	C	35	THR	4.0
1	D	146	VAL	4.0
1	D	171	ILE	4.0
1	C	53	THR	4.0
1	D	152	ASP	4.0
1	B	58	THR	3.9
1	C	17	ILE	3.9
1	D	100	ASP	3.8
1	C	40	THR	3.8
1	C	139	CYS	3.8
1	D	40	THR	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	29	VAL	3.8
1	D	51	ARG	3.8
1	D	78	VAL	3.7
1	C	43	THR	3.7
1	A	54	ARG	3.6
1	C	136	LEU	3.5
1	D	67	VAL	3.5
1	A	55	LYS	3.5
1	B	14	LYS	3.5
1	D	96	ALA	3.4
1	C	36	TYR	3.4
1	B	165	GLY	3.4
1	D	87	MET	3.4
1	D	173	LEU	3.4
1	C	49	VAL	3.4
1	B	162	PRO	3.4
1	C	31	LEU	3.3
1	D	98	LEU	3.3
1	A	72	LEU	3.3
1	D	59	ALA	3.3
1	D	85	PRO	3.2
1	C	135	GLY	3.1
1	C	32	GLU	3.1
1	D	101	ASP	3.1
1	D	121	GLY	3.1
1	C	19	SER	3.1
1	C	99	ILE	3.1
1	A	167	PHE	3.0
1	D	70	ARG	3.0
1	D	18	ILE	3.0
1	D	148	ILE	3.0
1	B	191	LEU	2.9
1	D	76	CYS	2.9
1	D	14	LYS	2.9
1	D	46	TRP	2.9
1	C	190	HIS	2.9
1	C	37	MET	2.8
1	A	14	LYS	2.8
1	B	18	ILE	2.8
1	D	73	HIS	2.7
1	A	164	ASP	2.7
1	D	16	TYR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	22	LEU	2.7
1	D	68	LEU	2.7
1	D	110	LEU	2.7
1	C	58	THR	2.7
1	C	15	GLN	2.7
1	C	129	ALA	2.7
1	C	46	TRP	2.7
1	D	49	VAL	2.7
1	D	113	LEU	2.7
1	D	178	LEU	2.7
1	A	163	GLY	2.7
1	D	107	ALA	2.7
1	B	22	LEU	2.6
1	C	74	TYR	2.6
1	C	20	GLU	2.6
1	A	53	THR	2.6
1	B	163	GLY	2.6
1	C	165	GLY	2.6
1	D	136	LEU	2.6
1	C	134	PRO	2.6
1	D	33	LYS	2.6
1	D	21	GLU	2.6
1	D	88	GLY	2.6
1	C	34	THR	2.5
1	D	149	ASN	2.5
1	C	187	ALA	2.5
1	D	108	ALA	2.5
1	D	102	GLY	2.5
1	C	48	SER	2.5
1	D	72	LEU	2.5
1	C	184	ALA	2.5
1	D	86	PRO	2.5
1	C	41	GLY	2.5
1	A	92	ILE	2.4
1	D	117	THR	2.4
1	D	164	ASP	2.4
1	C	47	GLU	2.4
1	D	161	LYS	2.4
1	D	83	PHE	2.4
1	B	176	ASN	2.4
1	B	29	VAL	2.4
1	D	169	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	118	GLY	2.4
1	D	122	ASP	2.4
1	C	130	VAL	2.4
1	D	174	PRO	2.3
1	B	186	VAL	2.3
1	D	166	GLU	2.3
1	B	101	ASP	2.3
1	C	30	LYS	2.3
1	D	94	PHE	2.3
1	C	21	GLU	2.3
1	D	62	VAL	2.3
1	A	39	PRO	2.3
1	D	162	PRO	2.3
1	C	207	ALA	2.3
1	D	124	ALA	2.3
1	C	51	ARG	2.3
1	A	58	THR	2.2
1	D	172	SER	2.2
1	A	41	GLY	2.2
1	C	92	ILE	2.2
1	D	34	THR	2.2
1	A	22	LEU	2.2
1	D	165	GLY	2.2
1	C	167	PHE	2.2
1	B	23	ILE	2.2
1	C	42	LYS	2.2
1	C	161	LYS	2.2
1	D	139	CYS	2.2
1	B	159	LYS	2.2
1	C	104	THR	2.2
1	D	104	THR	2.2
1	C	163	GLY	2.1
1	C	45	THR	2.1
1	D	52	THR	2.1
1	D	84	ARG	2.1
1	B	187	ALA	2.1
1	D	80	VAL	2.1
1	D	197	VAL	2.1
1	C	138	ASN	2.1
1	D	60	ASP	2.1
1	D	35	THR	2.1
1	D	53	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	192	THR	2.1
1	D	65	ILE	2.1
1	C	88	GLY	2.1
1	C	98	LEU	2.1
1	C	141	ILE	2.1
1	D	63	ALA	2.0
1	D	109	ALA	2.0
1	B	104	THR	2.0
1	D	45	THR	2.0
1	C	55	LYS	2.0
1	C	100	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ELQ	B	305	17/17	0.76	0.30	56,62,68,68	17
4	EDO	C	301	4/4	0.87	0.12	36,47,49,57	0
3	CL	B	301	1/1	0.88	0.15	69,69,69,69	0
4	EDO	B	302	4/4	0.89	0.11	46,46,47,50	0
2	MG	B	304	1/1	0.90	0.09	71,71,71,71	0
2	MG	C	302	1/1	0.90	0.07	70,70,70,70	0
2	MG	A	302	1/1	0.92	0.07	65,65,65,65	0
2	MG	D	301	1/1	0.94	0.06	40,40,40,40	0
2	MG	D	302	1/1	0.95	0.08	46,46,46,46	0
2	MG	B	303	1/1	0.98	0.03	34,34,34,34	0
2	MG	A	301	1/1	0.99	0.03	28,28,28,28	0
2	MG	C	303	1/1	0.99	0.02	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.