



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:25 AM UTC

PDB ID : 5QJA / pdb_00005qja
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z1497321453
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	11% 82%10%7%
1	B	209	12% 85%8%7%
1	C	209	21% 82%9%8%
1	D	209	32% 78%13%9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1454	920	243	283	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1416	892	238	279	7			
1	D	190	Total	C	N	O	S	0	1	0
			1425	903	240	274	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is (2,5-dimethyl-1,3-thiazol-4-yl)(pyrrolidin-1-yl)methanone (CCD ID: K0A) (formula: C₁₀H₁₄N₂OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			14	10	2	1	1		
4	B	1	Total	C	N	O	S	0	0
			14	10	2	1	1		
4	D	1	Total	C	N	O	S	0	0
			14	10	2	1	1		
4	D	1	Total	C	N	O	S	0	0
			14	10	2	1	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	77	Total	O	0	0
			77	77		
6	B	77	Total	O	0	0
			77	77		
6	C	74	Total	O	0	0
			74	74		
6	D	47	Total	O	0	0
			47	47		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.44Å 59.84Å 80.19Å 79.33° 81.51° 75.50°	Depositor
Resolution (Å)	78.35 – 1.64 78.35 – 1.64	Depositor EDS
% Data completeness (in resolution range)	95.8 (78.35-1.64) 95.8 (78.35-1.64)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.225 , 0.257 (Not available) , 0.262	Depositor DCC
R_{free} test set	5375 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.143	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6119	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL, K0A, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.30	3/1482 (0.2%)	1.13	1/2023 (0.0%)
1	B	1.16	0/1492	1.12	1/2035 (0.0%)
1	C	1.20	2/1443 (0.1%)	1.17	6/1974 (0.3%)
1	D	1.18	4/1453 (0.3%)	1.15	4/1984 (0.2%)
All	All	1.21	9/5870 (0.2%)	1.14	12/8016 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	174	PRO	CA-C	-8.23	1.43	1.52
1	D	84	ARG	C-O	-7.32	1.18	1.24
1	A	181	ARG	N-CA	-6.25	1.38	1.46
1	A	104	THR	CA-C	5.69	1.60	1.52
1	C	92	ILE	CA-CB	5.53	1.60	1.54
1	C	198	TYR	C-O	-5.46	1.17	1.24
1	D	144	VAL	N-CA	-5.41	1.39	1.46
1	D	144	VAL	C-O	-5.29	1.18	1.24
1	D	201	ALA	CA-C	5.06	1.59	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	38	ASP	CA-C-N	6.42	125.85	118.85
1	C	38	ASP	C-N-CA	6.42	125.85	118.85
1	C	127	SER	CA-C-N	-6.29	111.97	119.84
1	C	127	SER	C-N-CA	-6.29	111.97	119.84
1	A	188	GLU	N-CA-C	6.21	120.86	113.28
1	D	94	PHE	N-CA-C	-5.93	102.30	109.83
1	B	190	HIS	N-CA-C	5.70	118.15	111.02
1	D	138	ASN	N-CA-C	-5.55	106.10	112.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	45	THR	N-CA-C	5.22	116.92	109.14
1	C	190	HIS	N-CA-CB	5.20	117.78	110.24
1	D	132	MET	N-CA-C	5.11	116.66	111.14
1	D	18	ILE	N-CA-C	-5.09	106.72	111.45

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1454	0	1416	13	0
1	B	1464	0	1414	12	0
1	C	1416	0	1324	11	0
1	D	1425	0	1375	13	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
4	A	14	0	0	0	0
4	B	14	0	0	0	0
4	D	28	0	0	0	0
5	A	4	0	6	0	0
5	B	4	0	6	1	0
5	C	8	0	12	0	0
5	D	4	0	6	1	0
6	A	77	0	0	0	0
6	B	77	0	0	1	0
6	C	74	0	0	2	0
6	D	47	0	0	3	0
All	All	6119	0	5559	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:104:THR:HG23	1:C:107:ALA:H	1.52	0.75
1:C:125:GLU:OE1	6:C:401:HOH:O	2.04	0.74
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.71	0.72
1:B:104:THR:HG23	1:B:107:ALA:H	1.58	0.69
1:D:188:GLU:HG3	1:D:189:GLU:HG3	1.75	0.68
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.78	0.66
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.77	0.66
1:A:104:THR:HG23	1:A:107:ALA:H	1.60	0.66
1:D:112:GLU:OE2	6:D:401:HOH:O	2.14	0.65
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.28	0.64
6:C:401:HOH:O	1:D:206:HIS:HE1	1.80	0.64
5:D:305:EDO:O2	6:D:402:HOH:O	2.15	0.63
1:D:120:LYS:H	1:D:155:ASN:HD21	1.50	0.59
1:A:71:THR:HG23	1:A:151:ASP:OD2	2.02	0.59
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.40	0.57
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.88	0.55
1:C:132:MET:O	1:D:196[B]:ARG:NH1	2.41	0.53
1:B:158:PRO:O	1:B:160:PRO:HD3	2.10	0.52
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.46	0.51
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.41	0.50
1:C:206:HIS:HE1	1:D:125:GLU:OE1	1.94	0.50
1:D:191:LEU:C	1:D:191:LEU:HD12	2.36	0.50
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.94	0.50
1:A:104:THR:CG2	1:A:107:ALA:H	2.23	0.50
1:A:200:TYR:OH	1:B:206:HIS:HD2	1.94	0.50
1:B:58:THR:CB	1:B:142:HIS:NE2	2.75	0.49
1:D:177:ASP:OD1	1:D:180:GLN:CB	2.60	0.49
1:C:136:LEU:HG	6:D:402:HOH:O	2.13	0.48
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.52	0.45
1:C:206:HIS:HD2	1:D:200:TYR:OH	2.00	0.43
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.52	0.43
5:B:301:EDO:C2	6:B:405:HOH:O	2.66	0.43
1:A:54:ARG:HD3	1:A:60:ASP:OD1	2.20	0.41
1:B:54:ARG:HG2	1:B:55:LYS:N	2.35	0.41
1:A:206:HIS:CD2	1:B:200:TYR:OH	2.67	0.41
1:B:190:HIS:O	1:B:190:HIS:CG	2.74	0.41
1:D:110:LEU:HG	1:D:123:ILE:HD11	2.01	0.41
1:A:104:THR:HG22	1:A:107:ALA:CB	2.50	0.41
1:A:104:THR:HG22	1:A:107:ALA:HB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	188 (98%)	3 (2%)	1 (0%)	24	9
1	B	190/209 (91%)	184 (97%)	5 (3%)	1 (0%)	24	9
1	C	188/209 (90%)	185 (98%)	2 (1%)	1 (0%)	24	9
1	D	187/209 (90%)	180 (96%)	7 (4%)	0	100	100
All	All	757/836 (91%)	737 (97%)	17 (2%)	3 (0%)	30	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	66	PRO
1	A	66	PRO
1	B	66	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/179 (85%)	145 (95%)	7 (5%)	24	4
1	B	154/179 (86%)	152 (99%)	2 (1%)	61	38
1	C	142/179 (79%)	137 (96%)	5 (4%)	32	6
1	D	147/179 (82%)	141 (96%)	6 (4%)	27	4
All	All	595/716 (83%)	575 (97%)	20 (3%)	32	7

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	38	ASP
1	A	43	THR
1	A	55	LYS
1	A	70	ARG
1	A	125	GLU
1	A	191	LEU
1	B	38	ASP
1	B	188	GLU
1	C	31	LEU
1	C	45	THR
1	C	71	THR
1	C	164	ASP
1	C	189	GLU
1	D	28	TRP
1	D	43	THR
1	D	45	THR
1	D	70	ARG
1	D	72	LEU
1	D	208	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	73	HIS
1	A	206	HIS
1	B	15	GLN
1	B	69	GLN
1	B	73	HIS
1	B	82	GLN
1	B	206	HIS
1	B	208	ASN
1	C	15	GLN
1	C	206	HIS
1	D	15	GLN
1	D	149	ASN
1	D	155	ASN
1	D	190	HIS
1	D	206	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 9 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	EDO	C	301	-	3,3,3	0.82	0	2,2,2	0.58	0
4	K0A	B	304	-	15,15,15	1.07	0	19,21,21	1.16	2 (10%)
5	EDO	B	301	-	3,3,3	0.67	0	2,2,2	0.43	0
4	K0A	A	304	-	15,15,15	1.56	2 (13%)	19,21,21	1.55	4 (21%)
4	K0A	D	304	-	15,15,15	1.40	1 (6%)	19,21,21	1.67	5 (26%)
5	EDO	D	305	-	3,3,3	0.37	0	2,2,2	0.50	0
5	EDO	A	305	-	3,3,3	0.50	0	2,2,2	0.53	0
5	EDO	C	304	-	3,3,3	0.48	0	2,2,2	0.48	0
4	K0A	D	303	-	15,15,15	1.39	1 (6%)	19,21,21	1.21	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	301	-	-	0/1/1/1	-
4	K0A	B	304	-	-	3/8/15/15	0/2/2/2
5	EDO	B	301	-	-	0/1/1/1	-
4	K0A	A	304	-	-	2/8/15/15	0/2/2/2
4	K0A	D	304	-	-	2/8/15/15	0/2/2/2
5	EDO	D	305	-	-	0/1/1/1	-
5	EDO	A	305	-	-	0/1/1/1	-
5	EDO	C	304	-	-	0/1/1/1	-
4	K0A	D	303	-	-	2/8/15/15	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	304	K0A	C3-N1	4.32	1.42	1.34
4	A	304	K0A	C3-N1	4.17	1.41	1.34
4	D	303	K0A	C3-N1	3.99	1.41	1.34
4	A	304	K0A	C-C1	-2.02	1.44	1.49

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	304	K0A	C9-C8-C2	-3.47	126.80	130.10
4	A	304	K0A	C2-C8-S	3.40	111.30	108.89
4	A	304	K0A	S-C1-N	3.03	116.73	114.64
4	D	304	K0A	C2-C3-N1	3.01	123.88	118.97
4	A	304	K0A	C2-N-C1	-2.91	107.68	110.55
4	B	304	K0A	C3-C2-N	-2.85	116.18	122.48
4	D	304	K0A	O-C3-N1	-2.65	118.48	122.67
4	D	304	K0A	C2-C8-S	2.57	110.70	108.89
4	A	304	K0A	C3-C2-N	-2.46	117.04	122.48
4	D	304	K0A	S-C1-N	2.16	116.13	114.64
4	B	304	K0A	S-C1-N	2.14	116.11	114.64
4	D	303	K0A	C2-C8-S	2.08	110.36	108.89
4	D	303	K0A	C3-C2-N	-2.03	117.99	122.48

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	304	K0A	N-C2-C3-O
4	B	304	K0A	N-C2-C3-O

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	D	303	K0A	N-C2-C3-O
4	D	304	K0A	N-C2-C3-O
4	D	303	K0A	N-C2-C3-N1
4	B	304	K0A	O-C3-N1-C7
4	A	304	K0A	N-C2-C3-N1
4	B	304	K0A	N-C2-C3-N1
4	D	304	K0A	N-C2-C3-N1

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	301	EDO	1	0
5	D	305	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	0.65	24 (12%) 8 9	17, 26, 47, 54	11 (5%)
1	B	194/209 (92%)	0.75	26 (13%) 7 8	19, 30, 51, 67	3 (1%)
1	C	192/209 (91%)	1.14	43 (22%) 2 2	19, 32, 53, 63	4 (2%)
1	D	190/209 (90%)	1.83	67 (35%) 1 1	10, 34, 57, 73	17 (8%)
All	All	770/836 (92%)	1.09	160 (20%) 2 3	10, 31, 53, 73	35 (4%)

All (160) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	207	ALA	12.8
1	D	208	ASN	9.0
1	A	190	HIS	8.1
1	D	155	ASN	8.1
1	D	53	THR	7.8
1	A	189	GLU	7.6
1	D	73	HIS	6.6
1	D	154	GLU	6.2
1	A	55	LYS	5.8
1	D	31	LEU	5.6
1	C	28	TRP	5.4
1	C	16	TYR	5.4
1	C	58	THR	5.2
1	D	52	THR	5.2
1	D	29	VAL	5.1
1	D	23	ILE	4.9
1	D	40	THR	4.6
1	B	28	TRP	4.5
1	D	28	TRP	4.5
1	D	26	GLY	4.4
1	A	56	GLU	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	18	ILE	4.3
1	B	208	ASN	4.2
1	C	22	LEU	4.1
1	A	28	TRP	4.1
1	D	167	PHE	4.1
1	C	17	ILE	4.1
1	D	41	GLY	4.1
1	D	16	TYR	4.0
1	D	74	TYR	4.0
1	C	102	GLY	3.9
1	D	196[A]	ARG	3.8
1	C	207	ALA	3.7
1	D	161	LYS	3.6
1	D	153	ALA	3.6
1	C	136	LEU	3.6
1	C	43	THR	3.5
1	B	165	GLY	3.5
1	D	88	GLY	3.5
1	C	23	ILE	3.5
1	C	190	HIS	3.4
1	D	158	PRO	3.3
1	A	54	ARG	3.3
1	D	156	ALA	3.3
1	C	40	THR	3.2
1	D	152	ASP	3.2
1	D	173	LEU	3.2
1	D	100	ASP	3.2
1	D	43	THR	3.2
1	D	160	PRO	3.2
1	C	57	GLN	3.2
1	C	74	TYR	3.1
1	D	90	TYR	3.1
1	D	101	ASP	3.1
1	B	40	THR	3.1
1	D	148	ILE	3.1
1	B	58	THR	3.1
1	D	162	PRO	3.0
1	B	31	LEU	3.0
1	D	22	LEU	3.0
1	D	25	GLU	3.0
1	A	167	PHE	3.0
1	C	184	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	102	GLY	2.9
1	A	72	LEU	2.9
1	C	29	VAL	2.9
1	B	162	PRO	2.9
1	D	164	ASP	2.9
1	C	31	LEU	2.8
1	C	15	GLN	2.8
1	D	18	ILE	2.8
1	A	160	PRO	2.7
1	A	58	THR	2.7
1	C	104	THR	2.7
1	D	149	ASN	2.7
1	D	59	ALA	2.7
1	D	27	LYS	2.7
1	D	119	TYR	2.7
1	A	42	LYS	2.7
1	A	157	ARG	2.7
1	D	168	VAL	2.7
1	A	14	LYS	2.7
1	D	30	LYS	2.7
1	C	39	PRO	2.6
1	B	102	GLY	2.6
1	B	163	GLY	2.6
1	D	97	GLY	2.6
1	D	104	THR	2.6
1	D	159	LYS	2.6
1	B	176	ASN	2.6
1	C	53	THR	2.6
1	D	151	ASP	2.6
1	B	22	LEU	2.6
1	B	190	HIS	2.6
1	C	191	LEU	2.6
1	A	23	ILE	2.6
1	B	18	ILE	2.6
1	C	55	LYS	2.6
1	B	26	GLY	2.5
1	B	191	LEU	2.5
1	C	36	TYR	2.5
1	A	29	VAL	2.5
1	D	107	ALA	2.5
1	D	123	ILE	2.5
1	C	135	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	19	SER	2.5
1	B	153	ALA	2.5
1	D	163	GLY	2.5
1	C	54	ARG	2.5
1	C	46	TRP	2.4
1	B	14	LYS	2.4
1	B	104	THR	2.4
1	A	41	GLY	2.4
1	D	39	PRO	2.4
1	B	27	LYS	2.4
1	C	168	VAL	2.4
1	A	40	THR	2.3
1	D	96	ALA	2.3
1	C	161	LYS	2.3
1	C	172	SER	2.3
1	C	165	GLY	2.3
1	C	35	THR	2.3
1	D	172	SER	2.3
1	B	168	VAL	2.3
1	A	159	LYS	2.2
1	A	162	PRO	2.2
1	D	105	PRO	2.2
1	C	34	THR	2.2
1	C	52	THR	2.2
1	D	45	THR	2.2
1	A	104	THR	2.2
1	B	160	PRO	2.2
1	D	14	LYS	2.2
1	D	118	GLY	2.2
1	D	70	ARG	2.2
1	A	27	LYS	2.2
1	B	39	PRO	2.2
1	B	152	ASP	2.1
1	C	49	VAL	2.1
1	D	50	LYS	2.1
1	C	45	THR	2.1
1	C	21	GLU	2.1
1	C	160	PRO	2.1
1	D	83	PHE	2.1
1	D	36	TYR	2.1
1	D	49	VAL	2.1
1	B	41	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	55	LYS	2.1
1	B	161	LYS	2.1
1	D	99	ILE	2.1
1	C	41	GLY	2.1
1	C	50	LYS	2.1
1	A	158	PRO	2.1
1	D	46	TRP	2.1
1	A	156	ALA	2.0
1	D	78	VAL	2.0
1	A	46	TRP	2.0
1	D	171	ILE	2.0
1	D	122	ASP	2.0
1	C	42	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	A	303	1/1	0.79	0.25	72,72,72,72	0
5	EDO	C	301	4/4	0.79	0.16	30,38,40,45	0
5	EDO	B	301	4/4	0.83	0.14	38,40,41,47	0
4	K0A	D	303	14/14	0.85	0.22	41,47,56,56	14
4	K0A	D	304	14/14	0.85	0.18	32,38,42,45	14
2	MG	B	303	1/1	0.87	0.13	52,52,52,52	0
5	EDO	C	304	4/4	0.88	0.14	34,34,36,36	4
4	K0A	B	304	14/14	0.89	0.17	31,37,38,38	14
2	MG	C	302	1/1	0.89	0.10	41,41,41,41	0
2	MG	A	302	1/1	0.91	0.14	41,41,41,41	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	K0A	A	304	14/14	0.91	0.14	29,34,40,40	14
5	EDO	A	305	4/4	0.93	0.10	23,23,25,25	0
2	MG	D	301	1/1	0.93	0.06	31,31,31,31	0
2	MG	D	302	1/1	0.95	0.07	38,38,38,38	0
5	EDO	D	305	4/4	0.95	0.10	24,24,26,26	4
2	MG	C	303	1/1	0.97	0.05	33,33,33,33	0
2	MG	B	302	1/1	0.99	0.03	30,30,30,30	0
2	MG	A	301	1/1	0.99	0.03	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.