



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:09 AM UTC

PDB ID : 5QJO / pdb_00005qjo
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z57292369
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	11% 86% 6% • 7%
1	B	209	8% 83% 10% 7%
1	C	209	13% 83% 9% 8%
1	D	209	17% 81% 10% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1462	925	245	284	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1419	896	236	280	7			
1	D	190	Total	C	N	O	S	0	1	0
			1433	908	241	276	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is 2-methyl-N-(2-methyl-2H-tetrazol-5-yl)propanamide (CCD ID: K41) (formula: $C_6H_{11}N_5O$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 12 6 5 1	0	0
4	B	1	Total C N O 12 6 5 1	0	0
4	C	1	Total C N O 12 6 5 1	0	0
4	D	1	Total C N O 12 6 5 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	76	Total	O	0	0
			76	76		
6	B	80	Total	O	0	0
			80	80		
6	C	71	Total	O	0	0
			71	71		
6	D	36	Total	O	0	0
			36	36		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.06Å 59.57Å 79.93Å 79.55° 81.90° 75.89°	Depositor
Resolution (Å)	78.20 – 1.98 78.20 – 1.98	Depositor EDS
% Data completeness (in resolution range)	96.4 (78.20-1.98) 96.4 (78.20-1.98)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.210 , 0.249 (Not available) , 0.267	Depositor DCC
R_{free} test set	3007 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6110	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MG, CL, K41

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	1/1490 (0.1%)	1.10	5/2032 (0.2%)
1	B	1.10	1/1492 (0.1%)	1.07	3/2035 (0.1%)
1	C	1.18	1/1446 (0.1%)	1.08	4/1978 (0.2%)
1	D	1.13	1/1461 (0.1%)	1.09	1/1993 (0.1%)
All	All	1.15	4/5889 (0.1%)	1.08	13/8038 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	139	CYS	C-O	5.52	1.30	1.23
1	C	199	SER	CA-C	-5.37	1.45	1.52
1	D	84	ARG	C-O	-5.24	1.20	1.25
1	A	104	THR	CA-C	5.22	1.59	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	85	PRO	N-CA-C	6.71	118.89	110.70
1	D	138	ASN	N-CA-C	-6.32	105.20	113.17
1	A	161	LYS	CA-C-N	-6.00	113.78	119.85
1	A	161	LYS	C-N-CA	-6.00	113.78	119.85
1	A	53	THR	N-CA-C	5.76	120.15	113.18
1	A	84	ARG	CA-C-N	-5.34	114.88	120.38
1	A	84	ARG	C-N-CA	-5.34	114.88	120.38
1	B	38	ASP	CB-CA-C	5.27	118.98	109.45
1	C	107	ALA	N-CA-C	-5.15	105.35	111.69
1	B	43	THR	CB-CA-C	5.09	118.15	109.80
1	C	127	SER	CA-C-N	-5.07	115.34	120.21
1	C	127	SER	C-N-CA	-5.07	115.34	120.21
1	B	152	ASP	N-CA-C	-5.07	102.26	110.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1462	0	1433	12	0
1	B	1464	0	1414	12	0
1	C	1419	0	1338	13	0
1	D	1433	0	1390	14	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
4	A	12	0	0	0	0
4	B	12	0	0	0	0
4	C	12	0	0	0	0
4	D	12	0	0	0	0
5	B	8	0	12	0	0
5	C	4	0	6	0	0
6	A	76	0	0	0	0
6	B	80	0	0	0	0
6	C	71	0	0	3	0
6	D	36	0	0	0	0
All	All	6110	0	5593	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.78	0.65
1:C:104:THR:HG23	1:C:107:ALA:H	1.61	0.65
1:A:104:THR:HG23	1:A:107:ALA:H	1.62	0.65
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.80	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.84	0.60
1:C:206:HIS:HD2	1:D:200:TYR:OH	1.84	0.60
1:D:120:LYS:H	1:D:155:ASN:HD21	1.51	0.59
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.39	0.58
1:B:104:THR:HG23	1:B:107:ALA:H	1.71	0.56
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.36	0.55
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.89	0.54
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.44	0.53
1:D:104:THR:HG23	1:D:107:ALA:H	1.74	0.51
1:A:203:ALA:HB2	1:B:200:TYR:CD1	2.46	0.51
1:A:104:THR:HG22	1:A:107:ALA:CB	2.41	0.51
1:B:189:GLU:HB3	1:B:191:LEU:HD12	1.93	0.50
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.48	0.49
1:C:22:LEU:C	1:C:22:LEU:HD13	2.39	0.48
1:D:188:GLU:HG3	1:D:189:GLU:HG3	1.95	0.48
1:C:45:THR:HG23	6:C:460:HOH:O	2.13	0.47
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.96	0.47
1:A:104:THR:CG2	1:A:107:ALA:H	2.27	0.46
1:B:17:ILE:HD12	1:B:34:THR:CG2	2.45	0.46
1:B:179:LEU:CD2	1:B:205:LYS:HE2	2.45	0.45
1:D:149:ASN:C	1:D:149:ASN:HD22	2.24	0.45
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.48	0.44
1:A:206:HIS:O	1:B:175:LYS:NZ	2.47	0.44
1:C:74:TYR:HB3	6:C:431:HOH:O	2.18	0.44
1:C:15:GLN:N	6:C:405:HOH:O	2.51	0.43
1:D:149:ASN:HD21	1:D:151:ASP:HB2	1.83	0.43
1:C:206:HIS:HE1	1:D:125:GLU:OE1	2.02	0.43
1:A:64:VAL:O	1:A:66:PRO:HD3	2.18	0.43
1:D:37:MET:SD	1:D:41:GLY:O	2.77	0.43
1:D:149:ASN:C	1:D:149:ASN:ND2	2.75	0.43
1:A:134:PRO:HB2	1:B:84:ARG:HG3	2.01	0.43
1:C:71:THR:OG1	1:C:151:ASP:OD2	2.36	0.43
1:B:14:LYS:HE3	1:B:39:PRO:HB2	2.01	0.42
1:A:38:ASP:OD1	1:A:38:ASP:C	2.62	0.42
1:D:92:ILE:HG13	1:D:191:LEU:HD13	2.01	0.41
1:D:49:VAL:O	1:D:137:SER:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	188 (98%)	3 (2%)	1 (0%)	24	14
1	B	190/209 (91%)	186 (98%)	3 (2%)	1 (0%)	24	14
1	C	188/209 (90%)	184 (98%)	3 (2%)	1 (0%)	24	14
1	D	187/209 (90%)	183 (98%)	3 (2%)	1 (0%)	24	14
All	All	757/836 (91%)	741 (98%)	12 (2%)	4 (0%)	24	14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	66	PRO
1	A	66	PRO
1	D	66	PRO
1	B	66	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	5/179 (3%)	5 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 9 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	EDO	B	305	-	3,3,3	0.47	0	2,2,2	0.41	0
4	K41	C	304	-	12,12,12	2.31	5 (41%)	14,16,16	2.28	4 (28%)
5	EDO	C	301	-	3,3,3	0.64	0	2,2,2	0.17	0
5	EDO	B	301	-	3,3,3	0.68	0	2,2,2	0.38	0
4	K41	A	304	-	12,12,12	2.74	6 (50%)	14,16,16	2.11	4 (28%)
4	K41	B	304	-	12,12,12	2.49	5 (41%)	14,16,16	2.63	4 (28%)
4	K41	D	303	-	12,12,12	2.67	5 (41%)	14,16,16	1.77	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	305	-	-	0/1/1/1	-
4	K41	C	304	-	-	0/8/8/8	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	301	-	-	1/1/1/1	-
5	EDO	B	301	-	-	0/1/1/1	-
4	K41	A	304	-	-	0/8/8/8	0/1/1/1
4	K41	B	304	-	-	0/8/8/8	0/1/1/1
4	K41	D	303	-	-	0/8/8/8	0/1/1/1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	304	K41	N3-N2	5.48	1.38	1.31
4	B	304	K41	C4-N4	5.04	1.39	1.33
4	D	303	K41	N3-N4	4.48	1.39	1.33
4	D	303	K41	C4-N4	4.28	1.38	1.33
4	D	303	K41	N3-N2	4.22	1.37	1.31
4	C	304	K41	N3-N2	3.99	1.36	1.31
4	A	304	K41	C3-N	3.89	1.43	1.36
4	B	304	K41	N3-N2	3.84	1.36	1.31
4	C	304	K41	C3-N	3.77	1.43	1.36
4	B	304	K41	C3-N	3.76	1.43	1.36
4	A	304	K41	C4-N4	3.67	1.37	1.33
4	A	304	K41	C4-N1	3.66	1.45	1.34
4	C	304	K41	C4-N4	3.53	1.37	1.33
4	D	303	K41	C4-N1	3.48	1.44	1.34
4	D	303	K41	C3-N	3.23	1.42	1.36
4	C	304	K41	C4-N1	2.99	1.43	1.34
4	B	304	K41	C4-N1	2.62	1.42	1.34
4	A	304	K41	N3-N4	2.58	1.36	1.33
4	C	304	K41	N1-N2	-2.56	1.27	1.32
4	B	304	K41	N1-N2	-2.54	1.27	1.32
4	A	304	K41	C1-C3	2.42	1.58	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	K41	C5-N3-N2	6.44	127.06	122.51
4	A	304	K41	C5-N3-N2	5.97	126.73	122.51
4	B	304	K41	N3-N2-N1	5.97	109.24	106.10
4	C	304	K41	C5-N3-N2	5.64	126.50	122.51
4	C	304	K41	N3-N2-N1	4.73	108.59	106.10
4	D	303	K41	C5-N3-N2	3.99	125.33	122.51
4	D	303	K41	N3-N2-N1	3.82	108.11	106.10
4	A	304	K41	N3-N2-N1	3.23	107.80	106.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	304	K41	N4-N3-N2	-2.83	112.69	114.35
4	B	304	K41	N4-N3-N2	-2.67	112.78	114.35
4	A	304	K41	N4-N3-N2	-2.30	113.00	114.35
4	C	304	K41	C4-N4-N3	2.14	102.83	100.56
4	B	304	K41	C5-N3-N4	-2.12	120.10	122.38
4	A	304	K41	C5-N3-N4	-2.01	120.23	122.38

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	0.54	22 (11%) 10 13	16, 26, 51, 77	14 (7%)
1	B	194/209 (92%)	0.46	17 (8%) 15 22	18, 31, 61, 82	1 (0%)
1	C	192/209 (91%)	0.71	27 (14%) 6 9	17, 32, 58, 67	1 (0%)
1	D	190/209 (90%)	1.21	35 (18%) 3 4	11, 33, 60, 82	7 (3%)
All	All	770/836 (92%)	0.73	101 (13%) 7 10	11, 31, 59, 82	23 (2%)

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	207	ALA	12.2
1	C	17	ILE	12.1
1	D	208	ASN	10.2
1	A	191	LEU	10.0
1	D	156	ALA	9.1
1	D	206	HIS	8.3
1	A	190	HIS	8.2
1	D	155	ASN	8.0
1	A	188	GLU	7.9
1	A	189	GLU	7.6
1	D	154	GLU	5.5
1	B	163	GLY	4.5
1	D	40	THR	4.3
1	C	16	TYR	4.3
1	D	74	TYR	4.0
1	C	190	HIS	4.0
1	D	167	PHE	3.9
1	A	162	PRO	3.9
1	B	165	GLY	3.8
1	D	23	ILE	3.5
1	B	208	ASN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	23	ILE	3.3
1	B	14	LYS	3.3
1	D	16	TYR	3.3
1	B	160	PRO	3.2
1	D	160	PRO	3.2
1	A	157	ARG	3.1
1	A	164	ASP	3.1
1	D	153	ALA	3.1
1	C	22	LEU	3.1
1	C	18	ILE	3.0
1	D	166	GLU	3.0
1	A	163	GLY	3.0
1	D	152	ASP	3.0
1	D	161	LYS	3.0
1	B	162	PRO	2.9
1	B	190	HIS	2.9
1	D	73	HIS	2.9
1	C	55	LYS	2.9
1	C	58	THR	2.8
1	B	40	THR	2.8
1	D	162	PRO	2.8
1	D	100	ASP	2.8
1	D	53	THR	2.8
1	B	58	THR	2.8
1	B	191	LEU	2.8
1	D	150	GLY	2.7
1	D	43	THR	2.7
1	A	56	GLU	2.7
1	A	54	ARG	2.7
1	D	31	LEU	2.7
1	A	167	PHE	2.6
1	D	158	PRO	2.6
1	D	70	ARG	2.6
1	D	14	LYS	2.6
1	B	158	PRO	2.6
1	B	22	LEU	2.5
1	B	176	ASN	2.5
1	C	184	ALA	2.5
1	C	40	THR	2.5
1	D	119	TYR	2.5
1	A	160	PRO	2.5
1	C	34	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	165	GLY	2.4
1	C	74	TYR	2.4
1	C	43	THR	2.4
1	D	28	TRP	2.4
1	C	57	GLN	2.4
1	D	41	GLY	2.4
1	B	159	LYS	2.4
1	D	151	ASP	2.3
1	C	39	PRO	2.3
1	A	14	LYS	2.3
1	A	45	THR	2.3
1	A	187	ALA	2.3
1	D	168	VAL	2.3
1	B	153	ALA	2.2
1	C	100	ASP	2.2
1	D	29	VAL	2.2
1	C	153	ALA	2.2
1	C	187	ALA	2.2
1	C	207	ALA	2.2
1	B	28	TRP	2.2
1	A	55	LYS	2.2
1	C	37	MET	2.2
1	A	92	ILE	2.2
1	D	101	ASP	2.2
1	A	104	THR	2.1
1	C	35	THR	2.1
1	C	53	THR	2.1
1	A	72	LEU	2.1
1	D	107	ALA	2.1
1	B	157	ARG	2.1
1	A	40	THR	2.0
1	C	104	THR	2.0
1	C	102	GLY	2.0
1	C	101	ASP	2.0
1	A	166	GLU	2.0
1	C	30	LYS	2.0
1	C	165	GLY	2.0
1	D	165	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	C	301	4/4	0.73	0.19	35,50,52,57	0
5	EDO	B	301	4/4	0.77	0.17	42,46,49,50	0
4	K41	A	304	12/12	0.86	0.16	31,33,40,42	12
2	MG	A	302	1/1	0.87	0.11	68,68,68,68	0
3	CL	A	303	1/1	0.89	0.15	63,63,63,63	0
2	MG	B	303	1/1	0.90	0.11	70,70,70,70	0
2	MG	D	301	1/1	0.90	0.07	36,36,36,36	0
4	K41	C	304	12/12	0.91	0.14	30,34,44,46	12
4	K41	D	303	12/12	0.91	0.10	39,41,46,47	0
5	EDO	B	305	4/4	0.92	0.09	27,27,28,28	4
2	MG	C	302	1/1	0.92	0.08	42,42,42,42	0
4	K41	B	304	12/12	0.94	0.10	26,29,34,35	12
2	MG	A	301	1/1	0.98	0.04	26,26,26,26	0
2	MG	B	302	1/1	0.98	0.02	27,27,27,27	0
2	MG	C	303	1/1	0.98	0.03	28,28,28,28	0
2	MG	D	302	1/1	0.99	0.07	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.