



Full wwPDB X-ray Structure Validation Report ⓘ

May 7, 2026 – 08:38 AM EDT

PDB ID : 5QJR / pdb_00005qjr
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z220816104
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	 11% 82% 10% 7%
1	B	209	 13% 83% 10% 7%
1	C	209	 30% 81% 11% 8%
1	D	209	 52% 79% 11% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1462	925	245	284	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1402	884	234	277	7			
1	D	190	Total	C	N	O	S	0	1	0
			1404	887	235	274	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is (3R)-1-acetylpiperidine-3-carboxamide (CCD ID: K1V) (formula: C₈H₁₄N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			12	8	2	2		
3	A	1	Total	C	N	O	0	0
			12	8	2	2		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	74	Total	O	0	0
			74	74		
6	B	72	Total	O	0	0
			72	72		
6	C	69	Total	O	0	0
			69	69		
6	D	29	Total	O	0	0
			29	29		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase





- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.54Å 60.03Å 79.99Å 79.25° 81.06° 75.33°	Depositor
Resolution (Å)	78.08 – 1.62 78.08 – 1.62	Depositor EDS
% Data completeness (in resolution range)	96.3 (78.08-1.62) 96.3 (78.08-1.62)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.242 , 0.277 0.254 , 0.286	Depositor DCC
R_{free} test set	5590 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6025	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, EDO, K1V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	0/1490	1.04	1/2032 (0.0%)
1	B	1.05	1/1492 (0.1%)	1.04	0/2035
1	C	1.12	1/1429 (0.1%)	1.09	2/1958 (0.1%)
1	D	1.09	1/1430 (0.1%)	1.09	7/1954 (0.4%)
All	All	1.09	3/5841 (0.1%)	1.07	10/7979 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	84	ARG	C-O	-7.55	1.18	1.25
1	C	147	THR	N-CA	5.77	1.53	1.46
1	B	85	PRO	CA-C	5.08	1.56	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	94	PHE	CA-C-N	-6.50	112.81	119.83
1	D	94	PHE	C-N-CA	-6.50	112.81	119.83
1	C	190	HIS	N-CA-CB	6.19	119.96	109.87
1	D	188	GLU	N-CA-C	6.15	121.84	113.97
1	C	85	PRO	N-CA-C	6.12	118.17	110.70
1	D	193	VAL	N-CA-C	-6.06	99.58	108.97
1	A	188	GLU	N-CA-C	6.01	120.77	112.90
1	D	85	PRO	N-CA-C	5.91	117.90	110.70
1	D	176	ASN	N-CA-C	5.55	119.15	112.38
1	D	26	GLY	N-CA-C	5.10	117.47	110.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1462	0	1433	16	0
1	B	1464	0	1414	13	0
1	C	1402	0	1298	13	0
1	D	1404	0	1338	21	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	24	0	0	1	0
4	A	4	0	6	0	0
4	B	8	0	12	0	0
4	C	4	0	6	0	0
5	B	1	0	0	0	0
6	A	74	0	0	0	0
6	B	72	0	0	0	0
6	C	69	0	0	3	0
6	D	29	0	0	0	0
All	All	6025	0	5507	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:CG	1.44	1.23
1:D:196[B]:ARG:HG3	1:D:196[B]:ARG:NH2	1.38	1.10
1:D:196[B]:ARG:CG	1:D:196[B]:ARG:NH2	2.16	0.80
1:C:125:GLU:OE1	6:C:401:HOH:O	2.12	0.68
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.77	0.65
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.79	0.65
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:N	2.45	0.64
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.83	0.62
1:B:104:THR:HG23	1:B:107:ALA:H	1.65	0.61
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.84	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.33	0.58
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:HG3	0.54	0.57
1:C:206:HIS:HD2	1:D:200:TYR:OH	1.88	0.56
6:C:401:HOH:O	1:D:206:HIS:HE1	1.87	0.56
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.40	0.55
1:C:104:THR:HG23	1:C:107:ALA:H	1.71	0.55
1:A:104:THR:HG23	1:A:107:ALA:H	1.71	0.54
1:D:120:LYS:H	1:D:155:ASN:HD21	1.56	0.54
1:C:15:GLN:N	6:C:403:HOH:O	2.43	0.52
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.46	0.51
1:A:191:LEU:C	1:A:191:LEU:HD23	2.36	0.51
1:C:132:MET:O	1:D:196[B]:ARG:NH1	2.44	0.50
1:A:134:PRO:HB2	1:B:84:ARG:HG3	1.96	0.48
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.49	0.48
1:C:200:TYR:OH	1:D:206:HIS:HD2	1.96	0.48
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:H	2.12	0.47
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.98	0.47
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.44	0.47
1:C:188:GLU:C	1:C:189:GLU:HG3	2.40	0.46
1:A:28:TRP:CD1	3:A:303:K1V:N1	2.83	0.46
1:A:200:TYR:OH	1:B:206:HIS:HD2	1.99	0.46
1:D:177:ASP:OD1	1:D:180:GLN:CB	2.64	0.46
1:A:39:PRO:HG2	1:B:167:PHE:CD2	2.52	0.45
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.51	0.45
1:D:64:VAL:O	1:D:94:PHE:HB3	2.15	0.45
1:B:194:ASP:CG	1:B:196:ARG:HH21	2.23	0.44
1:A:71:THR:HG23	1:A:151:ASP:OD2	2.17	0.44
1:A:200:TYR:CG	1:B:203:ALA:HB2	2.53	0.44
1:B:17:ILE:HD12	1:B:34:THR:CG2	2.48	0.44
1:C:191:LEU:HD12	1:C:191:LEU:C	2.42	0.43
1:D:81:LYS:HA	1:D:91:CYS:O	2.18	0.43
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:CB	2.79	0.43
1:B:158:PRO:O	1:B:160:PRO:HD3	2.19	0.42
1:A:104:THR:CG2	1:A:107:ALA:H	2.32	0.42
1:A:179:LEU:HD11	1:A:198:TYR:CZ	2.56	0.41
1:A:71:THR:O	1:A:73:HIS:HD2	2.04	0.41
1:D:182:LEU:O	1:D:186:VAL:HG23	2.21	0.41
1:D:196[B]:ARG:H	1:D:196[B]:ARG:CZ	2.34	0.41
1:D:121:GLY:HA3	1:D:146:VAL:CG1	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	188 (98%)	3 (2%)	1 (0%)	24	10
1	B	190/209 (91%)	185 (97%)	4 (2%)	1 (0%)	24	10
1	C	188/209 (90%)	180 (96%)	6 (3%)	2 (1%)	11	2
1	D	187/209 (90%)	176 (94%)	11 (6%)	0	100	100
All	All	757/836 (91%)	729 (96%)	24 (3%)	4 (0%)	24	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	24	SER
1	A	66	PRO
1	C	66	PRO
1	B	66	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/179 (86%)	150 (97%)	4 (3%)	40	16
1	B	154/179 (86%)	151 (98%)	3 (2%)	50	26
1	C	139/179 (78%)	135 (97%)	4 (3%)	37	13
1	D	143/179 (80%)	141 (99%)	2 (1%)	59	37
All	All	590/716 (82%)	577 (98%)	13 (2%)	45	21

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	33	LYS
1	A	70	ARG
1	A	191	LEU
1	B	31	LEU
1	B	43	THR
1	B	205	LYS
1	C	22	LEU
1	C	31	LEU
1	C	35	THR
1	C	164	ASP
1	D	43	THR
1	D	72	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	82	GLN
1	A	206	HIS
1	B	15	GLN
1	B	73	HIS
1	B	190	HIS
1	B	206	HIS
1	C	206	HIS
1	D	15	GLN
1	D	69	GLN
1	D	82	GLN
1	D	155	ASN
1	D	206	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 9 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	K1V	A	304	-	12,12,12	1.39	2 (16%)	15,16,16	1.35	3 (20%)
4	EDO	B	305	-	3,3,3	0.46	0	2,2,2	0.34	0
4	EDO	C	301	-	3,3,3	0.47	0	2,2,2	0.26	0
3	K1V	A	303	-	12,12,12	1.25	2 (16%)	15,16,16	1.26	2 (13%)
4	EDO	A	305	-	3,3,3	0.46	0	2,2,2	0.67	0
4	EDO	B	302	-	3,3,3	0.52	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	K1V	A	304	-	-	4/8/18/18	0/1/1/1
4	EDO	B	305	-	-	0/1/1/1	-
4	EDO	C	301	-	-	1/1/1/1	-
3	K1V	A	303	-	-	2/8/18/18	0/1/1/1
4	EDO	A	305	-	-	0/1/1/1	-
4	EDO	B	302	-	-	0/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	304	K1V	C1-N	2.80	1.43	1.35
3	A	303	K1V	C2-N	2.75	1.52	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	304	K1V	C7-N1	2.22	1.38	1.32
3	A	303	K1V	C1-N	2.12	1.41	1.35

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	304	K1V	C6-C5-C7	-3.06	104.33	110.07
3	A	303	K1V	C6-C5-C7	-3.05	104.35	110.07
3	A	304	K1V	C4-C3-C2	-2.28	107.76	110.75
3	A	303	K1V	O1-C7-N1	-2.08	119.37	123.04
3	A	304	K1V	O1-C7-N1	-2.02	119.47	123.04

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	303	K1V	C4-C5-C7-N1
3	A	303	K1V	C4-C5-C7-O1
3	A	304	K1V	C4-C5-C7-N1
3	A	304	K1V	C4-C5-C7-O1
3	A	304	K1V	C-C1-N-C2
3	A	304	K1V	O-C1-N-C2
4	C	301	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	K1V	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	0.76	22 (11%)	10 10	19, 28, 50, 63	7 (3%)
1	B	194/209 (92%)	1.01	27 (13%)	6 6	20, 32, 54, 72	4 (2%)
1	C	192/209 (91%)	1.68	62 (32%)	1 1	20, 35, 61, 74	4 (2%)
1	D	190/209 (90%)	2.55	109 (57%)	0 0	8, 42, 64, 77	9 (4%)
All	All	770/836 (92%)	1.49	220 (28%)	1 1	8, 34, 60, 77	24 (3%)

All (220) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	23	ILE	17.1
1	C	22	LEU	15.7
1	D	156	ALA	12.4
1	C	24	SER	11.4
1	B	191	LEU	10.5
1	C	25	GLU	10.5
1	D	157	ARG	10.4
1	D	196[A]	ARG	10.1
1	D	207	ALA	9.6
1	A	189	GLU	9.1
1	A	190	HIS	9.1
1	D	155	ASN	8.6
1	B	190	HIS	8.5
1	B	28	TRP	8.5
1	B	189	GLU	7.6
1	D	208	ASN	7.4
1	D	190	HIS	7.4
1	D	154	GLU	7.2
1	D	189	GLU	6.4
1	D	97	GLY	6.3
1	D	167	PHE	6.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	18	ILE	5.9
1	D	158	PRO	5.9
1	D	153	ALA	5.9
1	C	16	TYR	5.6
1	D	74	TYR	5.3
1	A	28	TRP	5.2
1	C	43	THR	5.1
1	D	168	VAL	5.1
1	B	58	THR	5.0
1	B	165	GLY	4.9
1	D	23	ILE	4.9
1	D	90	TYR	4.8
1	D	28	TRP	4.7
1	C	40	THR	4.6
1	C	31	LEU	4.5
1	C	17	ILE	4.5
1	D	68	LEU	4.4
1	C	49	VAL	4.4
1	D	40	THR	4.4
1	D	152	ASP	4.3
1	D	31	LEU	4.3
1	D	160	PRO	4.3
1	C	34	THR	4.2
1	C	135	GLY	4.2
1	D	173	LEU	4.2
1	B	159	LYS	4.1
1	D	171	ILE	4.1
1	B	162	PRO	4.1
1	C	21	GLU	4.0
1	C	35	THR	4.0
1	B	57	GLN	4.0
1	D	100	ASP	4.0
1	C	29	VAL	4.0
1	A	163	GLY	3.9
1	C	104	THR	3.9
1	D	53	THR	3.8
1	B	208	ASN	3.8
1	A	27	LYS	3.7
1	C	39	PRO	3.7
1	D	151	ASP	3.7
1	D	46	TRP	3.7
1	A	56	GLU	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	148	ILE	3.7
1	D	96	ALA	3.7
1	A	164	ASP	3.7
1	C	19	SER	3.7
1	D	67	VAL	3.6
1	C	136	LEU	3.6
1	D	119	TYR	3.6
1	D	70	ARG	3.5
1	B	14	LYS	3.5
1	C	28	TRP	3.5
1	B	163	GLY	3.5
1	D	22	LEU	3.5
1	C	55	LYS	3.5
1	C	48	SER	3.4
1	D	162	PRO	3.4
1	C	36	TYR	3.4
1	D	59	ALA	3.4
1	C	30	LYS	3.4
1	C	53	THR	3.4
1	C	58	THR	3.4
1	D	88	GLY	3.4
1	C	37	MET	3.4
1	D	27	LYS	3.3
1	D	164	ASP	3.3
1	D	18	ILE	3.3
1	C	20	GLU	3.3
1	D	78	VAL	3.3
1	A	207	ALA	3.3
1	C	207	ALA	3.3
1	C	52	THR	3.3
1	A	167	PHE	3.2
1	B	176	ASN	3.2
1	A	157	ARG	3.2
1	D	51	ARG	3.2
1	B	29	VAL	3.2
1	D	117	THR	3.2
1	D	192	THR	3.1
1	D	139	CYS	3.1
1	A	160	PRO	3.1
1	C	26	GLY	3.1
1	D	110	LEU	3.1
1	D	174	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	104	THR	3.0
1	C	32	GLU	3.0
1	A	72	LEU	3.0
1	B	18	ILE	3.0
1	C	184	ALA	3.0
1	D	76	CYS	3.0
1	D	83	PHE	3.0
1	A	57	GLN	2.9
1	B	22	LEU	2.9
1	D	79	LEU	2.9
1	D	113	LEU	2.9
1	D	89	GLY	2.9
1	C	15	GLN	2.9
1	D	101	ASP	2.9
1	D	123	ILE	2.9
1	B	104	THR	2.8
1	B	101	ASP	2.8
1	D	161	LYS	2.8
1	A	162	PRO	2.8
1	C	46	TRP	2.8
1	D	121	GLY	2.8
1	D	91	CYS	2.8
1	D	99	ILE	2.8
1	D	166	GLU	2.8
1	D	14	LYS	2.8
1	D	122	ASP	2.8
1	A	58	THR	2.8
1	D	29	VAL	2.7
1	C	165	GLY	2.7
1	C	161	LYS	2.7
1	C	57	GLN	2.7
1	D	102	GLY	2.7
1	D	69	GLN	2.7
1	D	34	THR	2.7
1	D	49	VAL	2.7
1	C	41	GLY	2.7
1	C	99	ILE	2.7
1	D	24	SER	2.6
1	C	102	GLY	2.6
1	D	26	GLY	2.6
1	D	197	VAL	2.6
1	C	42	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	85	PRO	2.6
1	B	102	GLY	2.6
1	D	109	ALA	2.6
1	D	72	LEU	2.5
1	D	149	ASN	2.5
1	C	47	GLU	2.5
1	D	170	VAL	2.5
1	D	124	ALA	2.5
1	A	41	GLY	2.5
1	C	45	THR	2.5
1	C	72	LEU	2.5
1	C	162	PRO	2.5
1	D	52	THR	2.5
1	D	73	HIS	2.5
1	D	87	MET	2.5
1	B	158	PRO	2.5
1	D	41	GLY	2.5
1	C	141	ILE	2.5
1	D	98	LEU	2.4
1	D	191	LEU	2.4
1	D	165	GLY	2.4
1	D	107	ALA	2.4
1	B	168	VAL	2.4
1	C	27	LYS	2.4
1	B	160	PRO	2.4
1	D	178	LEU	2.4
1	C	74	TYR	2.4
1	D	16	TYR	2.4
1	A	23	ILE	2.4
1	D	136	LEU	2.4
1	D	120	LYS	2.4
1	C	168	VAL	2.4
1	D	82	GLN	2.3
1	D	108	ALA	2.3
1	C	139	CYS	2.3
1	D	147	THR	2.3
1	A	31	LEU	2.3
1	C	163	GLY	2.3
1	D	105	PRO	2.3
1	A	40	THR	2.3
1	A	156	ALA	2.3
1	C	191	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	159	LYS	2.3
1	C	88	GLY	2.3
1	C	186	VAL	2.3
1	C	137	SER	2.2
1	B	171	ILE	2.2
1	C	134	PRO	2.2
1	C	153	ALA	2.2
1	B	100	ASP	2.2
1	D	163	GLY	2.2
1	B	167	PHE	2.2
1	D	66	PRO	2.2
1	D	118	GLY	2.2
1	B	157	ARG	2.2
1	C	190	HIS	2.2
1	D	62	VAL	2.1
1	D	112	GLU	2.1
1	C	59	ALA	2.1
1	D	111	ARG	2.1
1	A	191	LEU	2.1
1	D	86	PRO	2.1
1	D	92	ILE	2.1
1	A	158	PRO	2.1
1	D	84	ARG	2.1
1	D	94	PHE	2.1
1	D	71	THR	2.1
1	D	39	PRO	2.1
1	D	81	LYS	2.0
1	D	150	GLY	2.0
1	B	23	ILE	2.0
1	D	65	ILE	2.0
1	C	130	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	K1V	A	304	12/12	0.73	0.27	43,49,53,54	12
4	EDO	B	305	4/4	0.76	0.19	50,51,51,54	0
2	MG	B	304	1/1	0.82	0.14	73,73,73,73	0
2	MG	C	302	1/1	0.82	0.15	59,59,59,59	0
4	EDO	C	301	4/4	0.82	0.17	39,52,55,59	0
3	K1V	A	303	12/12	0.83	0.21	33,36,38,40	12
5	CL	B	301	1/1	0.84	0.17	67,67,67,67	0
2	MG	A	302	1/1	0.85	0.10	59,59,59,59	0
4	EDO	B	302	4/4	0.89	0.12	41,43,43,45	0
2	MG	D	301	1/1	0.90	0.08	38,38,38,38	0
2	MG	D	302	1/1	0.94	0.19	40,40,40,40	0
4	EDO	A	305	4/4	0.94	0.08	26,27,28,29	0
2	MG	A	301	1/1	0.99	0.03	26,26,26,26	0
2	MG	B	303	1/1	0.99	0.02	27,27,27,27	0
2	MG	C	303	1/1	0.99	0.04	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.