



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 05:13 PM UTC

PDB ID : 5QJT / pdb_00005qjt
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z969560582
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	
1	B	209	
1	C	209	
1	D	209	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1454	920	243	283	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1408	887	234	280	7			
1	D	190	Total	C	N	O	S	0	1	0
			1429	905	240	276	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is N,2-dimethyl-1,3-thiazole-5-carboxamide (CCD ID: K24) (formula: C₆H₈N₂OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			10	6	2	1	1		
3	D	1	Total	C	N	O	S	0	0
			10	6	2	1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	82	Total	O	0	0
			82	82		
6	B	74	Total	O	0	0
			74	74		
6	C	67	Total	O	0	0
			67	67		
6	D	33	Total	O	0	0
			33	33		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.45Å 59.87Å 80.23Å 79.16° 81.53° 75.51°	Depositor
Resolution (Å)	78.36 – 1.62 78.36 – 1.62	Depositor EDS
% Data completeness (in resolution range)	94.8 (78.36-1.62) 94.8 (78.36-1.62)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.220 , 0.251 0.231 , 0.259	Depositor DCC
R_{free} test set	5577 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6056	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K24, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	0/1482	1.05	0/2023
1	B	1.13	1/1492 (0.1%)	1.07	2/2035 (0.1%)
1	C	1.19	2/1435 (0.1%)	1.05	0/1966
1	D	1.17	2/1457 (0.1%)	1.10	4/1989 (0.2%)
All	All	1.17	5/5866 (0.1%)	1.07	6/8013 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	84	ARG	C-O	-6.80	1.18	1.24
1	C	147	THR	N-CA	5.91	1.53	1.46
1	C	104	THR	CA-C	5.40	1.59	1.53
1	B	173	LEU	CA-C	-5.17	1.48	1.52
1	D	86	PRO	CA-C	5.08	1.59	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	85	PRO	CA-C-N	-7.15	111.41	119.28
1	B	85	PRO	C-N-CA	-7.15	111.41	119.28
1	D	94	PHE	N-CA-C	-5.13	103.00	110.08
1	D	178	LEU	N-CA-C	5.06	116.48	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	85	PRO	CA-C-N	-5.01	113.88	119.19
1	D	85	PRO	C-N-CA	-5.01	113.88	119.19

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	52	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1454	0	1416	18	0
1	B	1464	0	1414	14	0
1	C	1408	0	1307	16	0
1	D	1429	0	1379	16	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	10	0	0	0	0
3	D	10	0	0	0	0
4	A	4	0	6	0	0
4	B	8	0	12	1	0
4	C	4	0	6	0	0
5	B	1	0	0	0	0
6	A	82	0	0	1	0
6	B	74	0	0	2	0
6	C	67	0	0	4	0
6	D	33	0	0	1	0
All	All	6056	0	5540	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.53	0.89
1:C:125:GLU:OE1	6:C:401:HOH:O	2.02	0.77
1:D:120:LYS:H	1:D:155:ASN:HD21	1.33	0.77
1:C:104:THR:HG23	1:C:107:ALA:H	1.50	0.75
1:C:112:GLU:OE2	6:C:402:HOH:O	2.12	0.68
6:C:401:HOH:O	1:D:206:HIS:HE1	1.80	0.65
1:A:104:THR:HG23	1:A:107:ALA:H	1.62	0.64
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.79	0.63
1:B:104:THR:HG23	1:B:107:ALA:H	1.65	0.61
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.84	0.59
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.85	0.58
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.34	0.57
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.88	0.56
1:D:104:THR:HG23	1:D:107:ALA:H	1.69	0.56
1:A:92:ILE:HD11	1:A:191:LEU:HD11	1.87	0.56
1:B:189:GLU:HB3	1:B:191:LEU:HD12	1.87	0.56
1:D:194:ASP:OD1	1:D:196[A]:ARG:HD3	2.08	0.54
1:A:112:GLU:OE1	6:A:401:HOH:O	2.18	0.54
1:A:71:THR:HG23	1:A:151:ASP:OD2	2.08	0.54
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.46	0.50
1:C:92:ILE:HD12	1:C:191:LEU:HD11	1.94	0.50
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.47	0.49
1:C:206:HIS:HD2	1:D:200:TYR:OH	1.95	0.49
4:B:302:EDO:C2	6:B:406:HOH:O	2.61	0.49
1:A:82:GLN:HG2	1:A:168:VAL:HG22	1.97	0.47
1:A:167:PHE:CD2	1:B:39:PRO:HG2	2.51	0.46
1:C:200:TYR:OH	1:D:206:HIS:HD2	1.97	0.46
1:A:200:TYR:OH	1:B:206:HIS:HD2	1.99	0.46
1:A:54:ARG:HD3	1:A:60:ASP:OD1	2.16	0.46
1:B:190:HIS:O	1:B:190:HIS:CG	2.68	0.46
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.51	0.45
1:A:179:LEU:HD23	1:A:205:LYS:HD2	1.98	0.44
1:B:175:LYS:HE2	6:B:472:HOH:O	2.18	0.44
1:A:125:GLU:HG3	1:A:145:THR:HB	2.00	0.44
1:A:206:HIS:HE1	1:B:125:GLU:OE1	2.00	0.44
1:C:206:HIS:HE1	1:D:125:GLU:OE1	2.00	0.44
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.48	0.44
1:A:104:THR:CG2	1:A:107:ALA:H	2.27	0.43
1:C:131:CYS:SG	1:D:194:ASP:HA	2.59	0.43
1:A:71:THR:O	1:A:73:HIS:HD2	2.02	0.43
1:D:70:ARG:HD3	6:D:411:HOH:O	2.19	0.42
1:C:15:GLN:N	6:C:405:HOH:O	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:87:MET:HE3	1:D:87:MET:HB3	1.93	0.42
1:C:132:MET:O	1:D:196[B]:ARG:NH1	2.53	0.41
1:B:58:THR:CB	1:B:142:HIS:NE2	2.83	0.41
1:C:136:LEU:HD13	1:C:136:LEU:O	2.21	0.41
1:B:158:PRO:O	1:B:160:PRO:HD3	2.21	0.40
1:D:37:MET:SD	1:D:41:GLY:O	2.79	0.40
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.57	0.40
1:D:64:VAL:O	1:D:94:PHE:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	187 (97%)	4 (2%)	1 (0%)	24	10
1	B	190/209 (91%)	186 (98%)	3 (2%)	1 (0%)	24	10
1	C	188/209 (90%)	186 (99%)	1 (0%)	1 (0%)	24	10
1	D	187/209 (90%)	181 (97%)	5 (3%)	1 (0%)	24	10
All	All	757/836 (91%)	740 (98%)	13 (2%)	4 (0%)	24	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	PRO
1	B	66	PRO
1	C	66	PRO
1	D	66	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/179 (85%)	147 (97%)	5 (3%)	33	11
1	B	154/179 (86%)	151 (98%)	3 (2%)	50	26
1	C	141/179 (79%)	139 (99%)	2 (1%)	59	37
1	D	148/179 (83%)	143 (97%)	5 (3%)	32	10
All	All	595/716 (83%)	580 (98%)	15 (2%)	42	17

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	43	THR
1	A	70	ARG
1	A	125	GLU
1	A	191	LEU
1	B	14	LYS
1	B	31	LEU
1	B	38	ASP
1	C	31	LEU
1	C	164	ASP
1	D	24	SER
1	D	25	GLU
1	D	43	THR
1	D	72	LEU
1	D	208	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	206	HIS
1	B	15	GLN
1	B	190	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	206	HIS
1	C	206	HIS
1	D	15	GLN
1	D	149	ASN
1	D	155	ASN
1	D	206	HIS
1	D	208	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 9 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	K24	A	303	-	10,10,10	2.39	1 (10%)	12,13,13	3.11	5 (41%)
4	EDO	B	302	-	3,3,3	0.68	0	2,2,2	0.41	0
3	K24	D	303	-	10,10,10	2.48	1 (10%)	12,13,13	2.76	3 (25%)
4	EDO	B	305	-	3,3,3	0.48	0	2,2,2	0.37	0
4	EDO	C	301	-	3,3,3	0.57	0	2,2,2	0.35	0
4	EDO	A	304	-	3,3,3	0.52	0	2,2,2	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	K24	A	303	-	-	2/6/6/6	0/1/1/1
4	EDO	B	302	-	-	0/1/1/1	-
3	K24	D	303	-	-	2/6/6/6	0/1/1/1
4	EDO	B	305	-	-	0/1/1/1	-
4	EDO	C	301	-	-	1/1/1/1	-
4	EDO	A	304	-	-	0/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	303	K24	C1-N	7.15	1.43	1.33
3	A	303	K24	C1-N	6.82	1.42	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	303	K24	C-N-C1	7.23	129.27	121.90
3	D	303	K24	C-N-C1	6.45	128.47	121.90
3	A	303	K24	C2-C1-N	5.65	126.57	116.88
3	D	303	K24	C2-C1-N	5.02	125.48	116.88
3	A	303	K24	O-C1-N	-3.66	116.65	122.84
3	D	303	K24	O-C1-N	-3.55	116.83	122.84
3	A	303	K24	S-C4-N1	2.55	116.36	114.63
3	A	303	K24	O-C1-C2	-2.12	116.77	120.62

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	303	K24	O-C1-N-C
3	A	303	K24	C2-C1-N-C
3	D	303	K24	O-C1-N-C
3	D	303	K24	C2-C1-N-C
4	C	301	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	302	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	0.55	17 (8%) 15 16	16, 28, 52, 62	6 (3%)
1	B	194/209 (92%)	0.68	25 (12%) 7 8	20, 33, 57, 71	1 (0%)
1	C	192/209 (91%)	0.96	38 (19%) 3 3	20, 33, 57, 64	0
1	D	190/209 (90%)	1.56	57 (30%) 1 1	13, 37, 61, 74	4 (2%)
All	All	770/836 (92%)	0.93	137 (17%) 4 3	13, 33, 57, 74	11 (1%)

All (137) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	207	ALA	12.5
1	D	208	ASN	9.8
1	A	191	LEU	9.7
1	A	189	GLU	8.4
1	A	190	HIS	7.9
1	D	154	GLU	6.4
1	D	40	THR	4.6
1	D	158	PRO	4.5
1	D	74	TYR	4.3
1	B	28	TRP	4.0
1	D	16	TYR	4.0
1	C	23	ILE	3.9
1	D	167	PHE	3.9
1	C	43	THR	3.8
1	B	58	THR	3.8
1	B	208	ASN	3.8
1	C	17	ILE	3.8
1	B	165	GLY	3.8
1	C	16	TYR	3.7
1	B	102	GLY	3.6
1	C	40	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	14	LYS	3.6
1	D	31	LEU	3.6
1	D	153	ALA	3.5
1	C	22	LEU	3.5
1	D	41	GLY	3.5
1	D	152	ASP	3.4
1	A	167	PHE	3.4
1	D	53	THR	3.4
1	B	191	LEU	3.3
1	C	29	VAL	3.3
1	D	29	VAL	3.3
1	C	104	THR	3.3
1	B	163	GLY	3.3
1	C	191	LEU	3.2
1	D	23	ILE	3.2
1	D	163	GLY	3.2
1	C	58	THR	3.2
1	D	52	THR	3.2
1	D	160	PRO	3.1
1	D	101	ASP	3.1
1	C	161	LYS	3.1
1	C	39	PRO	3.0
1	D	14	LYS	3.0
1	D	102	GLY	3.0
1	B	159	LYS	3.0
1	A	27	LYS	2.9
1	B	104	THR	2.9
1	D	164	ASP	2.9
1	B	176	ASN	2.9
1	C	184	ALA	2.9
1	C	42	LYS	2.8
1	A	72	LEU	2.8
1	D	159	LYS	2.8
1	C	207	ALA	2.8
1	D	88	GLY	2.8
1	B	158	PRO	2.8
1	C	136	LEU	2.8
1	D	70	ARG	2.7
1	D	171	ILE	2.7
1	A	29	VAL	2.7
1	C	18	ILE	2.7
1	D	100	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	39	PRO	2.7
1	D	39	PRO	2.7
1	C	28	TRP	2.7
1	D	90	TYR	2.7
1	D	97	GLY	2.7
1	D	196[A]	ARG	2.6
1	D	36	TYR	2.6
1	D	156	ALA	2.6
1	C	19	SER	2.6
1	C	74	TYR	2.6
1	B	18	ILE	2.6
1	D	28	TRP	2.6
1	C	102	GLY	2.6
1	D	59	ALA	2.6
1	B	46	TRP	2.6
1	B	162	PRO	2.6
1	A	40	THR	2.5
1	C	52	THR	2.5
1	A	41	GLY	2.5
1	C	34	THR	2.5
1	D	43	THR	2.5
1	B	190	HIS	2.5
1	D	118	GLY	2.5
1	D	155	ASN	2.4
1	D	151	ASP	2.4
1	C	35	THR	2.4
1	D	168	VAL	2.4
1	C	160	PRO	2.4
1	D	49	VAL	2.4
1	B	101	ASP	2.4
1	C	165	GLY	2.4
1	D	26	GLY	2.4
1	C	31	LEU	2.4
1	B	186	VAL	2.4
1	D	166	GLU	2.3
1	D	18	ILE	2.3
1	D	46	TRP	2.3
1	A	162	PRO	2.3
1	D	37	MET	2.3
1	D	68	LEU	2.3
1	C	24	SER	2.3
1	C	190	HIS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	162	PRO	2.2
1	D	25	GLU	2.2
1	C	27	LYS	2.2
1	C	167	PHE	2.2
1	D	119	TYR	2.2
1	C	53	THR	2.2
1	B	152	ASP	2.2
1	C	59	ALA	2.2
1	B	23	ILE	2.2
1	C	37	MET	2.1
1	D	107	ALA	2.1
1	A	163	GLY	2.1
1	A	56	GLU	2.1
1	B	160	PRO	2.1
1	C	187	ALA	2.1
1	D	73	HIS	2.1
1	A	22	LEU	2.1
1	A	158	PRO	2.1
1	C	36	TYR	2.1
1	A	207	ALA	2.1
1	D	22	LEU	2.1
1	C	46	TRP	2.1
1	A	23	ILE	2.1
1	D	123	ILE	2.1
1	D	165	GLY	2.0
1	D	21	GLU	2.0
1	B	22	LEU	2.0
1	C	72	LEU	2.0
1	A	104	THR	2.0
1	B	40	THR	2.0
1	B	167	PHE	2.0
1	D	76	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	B	302	4/4	0.82	0.15	41,45,48,48	0
5	CL	B	301	1/1	0.82	0.19	69,69,69,69	0
4	EDO	C	301	4/4	0.83	0.17	37,53,55,64	0
2	MG	A	302	1/1	0.85	0.12	63,63,63,63	0
3	K24	D	303	10/10	0.86	0.16	42,48,54,57	0
2	MG	B	304	1/1	0.89	0.12	77,77,77,77	0
3	K24	A	303	10/10	0.89	0.16	35,37,41,45	10
2	MG	D	301	1/1	0.90	0.06	34,34,34,34	0
4	EDO	A	304	4/4	0.94	0.09	26,27,27,27	0
4	EDO	B	305	4/4	0.96	0.08	34,35,35,36	0
2	MG	D	302	1/1	0.97	0.14	38,38,38,38	0
2	MG	C	302	1/1	0.98	0.07	43,43,43,43	0
2	MG	A	301	1/1	0.99	0.04	26,26,26,26	0
2	MG	B	303	1/1	0.99	0.03	31,31,31,31	0
2	MG	C	303	1/1	0.99	0.03	33,33,33,33	0

6.5 Other polymers

There are no such residues in this entry.