



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 01:25 PM UTC

PDB ID : 7QJY / pdb_00007qjy
EMDB ID : EMD-14026
Title : In vitro assembled 266/297 - 391 tau filaments with LiCl (9a)
Authors : Lovestam, S.; Scheres, S.H.W.
Deposited on : 2021-12-17
Resolution : 3.14 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14483 (2.64 - 3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	17% 83%
1	B	441	17% 83%
1	C	441	17% 83%
1	D	441	17% 83%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	441	 17%83%
1	F	441	 17%83%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Microtubule-associated protein tau.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	B	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	C	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	D	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	E	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	F	75	Total	C	N	O	S	0	0
			574	360	106	107	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Microtubule-associated protein tau



- Molecule 1: Microtubule-associated protein tau



[illegible]

- Molecule 1: Microtubule-associated protein tau

Chain C: 17% 83%

SER	PRO	SER	GLY	ARG	THR	HIS	SER	MET
ALA	GLY	ALA	GLY	LEU	PRO	VAL	GLU	ALA
SER	GLY	SER	LEU	THR	SER	GLN	THR	THR
LEU	S305	THR	ALA	GLY	ARG	ALA	ASP	PRO
ALA	PRO	GLU	PRO	GLY	MET	LYS	LYS	GLN
LYS	R379	VAL	VAL	VAL	VAL	VAL	SER	PHE
GLY	GLU	ASN	ASN	PRO	PRO	SER	THR	VAL
LEU	ALA	MET	PRO	PRO	LYS	LYS	THR	VAL
	LYS	PRO	ASP	GLY	SER	SER	ALA	MET
	ALA	ASP	LEU	ASP	GLY	LYS	GLU	GLN
	LYS	LEU	LYS	ARG	ASP	ASP	GLU	ASP
	THR	ASN	THR	SER	THR	GLY	ASP	HIS
	ASP	HIS	HIS	VAL	GLY	THR	VAL	ALA
	GLY	VAL	GLY	GLN	SER	SER	THR	SER
	THR	THR	THR	THR	TYR	SER	ALA	THR
	ALA	SER	GLY	LYS	SER	ASP	PRO	TYR
	VAL	LEU	ILE	LYS	SER	ASP	LEU	GLY
	THR	ILE	GLY	GLY	PRO	LYS	VAL	LEU
	VAL	SER	THR	SER	GLY	LYS	ASP	GLY
	VAL	GLY	THR	GLY	PRO	LYS	GLU	ASP
	SER	GLU	THR	THR	PRO	GLY	GLY	ARG
	PRO	ASN	THR	ASN	GLY	GLY	ALA	LYS
	VAL	GLN	SER	GLN	ARG	THR	ALA	THR
	GLY	PRO	ARG	GLY	THR	LYS	ALA	THR
	ASP	GLY	THR	THR	ARG	ILE	ALA	MET
	THR	GLY	GLY	THR	ALA	ALA	GLN	HIS
	PRO	GLY	PRO	PRO	THR	THR	GLN	GLN
	ARG	LYS	LYS	SER	PRO	HIS	ASP	ASP
	HIS	VAL	VAL	LEU	ARG	THR	GLN	GLN
	LEU	GLN	GLN	PRO	GLY	THR	GLU	GLU
	SER	ILE	ILE	THR	ALA	ALA	GLY	GLY
	ASN	ILE	ILE	PRO	ALA	PRO	ALA	LEU
	VAL	ASN	ASN	PRO	PRO	PRO	GLU	LYS
	SER	LYS	SER	THR	PRO	PRO	GLY	GLU
	SER	LYS	LYS	ARG	GLY	GLY	THR	ALA
	THR	LEU	LEU	GLU	GLN	THR	THR	ASP
	GLY	ASP	LEU	PRO	LYS	LYS	ALA	LEU
	SER	LEU	LEU	LYS	GLY	GLY	GLU	GLN
	ILE	SER	SER	LYS	GLN	ALA	ALA	SER
	ASP	MET	VAL	VAL	ASN	ASN	GLY	PRO
	VAL	ASP	GLN	VAL	ALA	ALA	ILE	LEU
	ASN	ASP	SER	VAL	THR	THR	GLY	GLY
	SER	SER	ASN	GLN	ALA	ALA	THR	THR
	THR	ASP	GLN	VAL	ALA	THR	GLY	GLY
	VAL	ASN	THR	VAL	THR	THR	GLY	THR
	THR	SER	THR	VAL	ARG	ILE	GLY	LEU
	ASN	SER	ASN	GLN	ALA	ALA	GLY	GLN
	VAL	THR	THR	THR	THR	THR	GLY	THR
	THR	LYS	LYS	VAL	LYS	LYS	LEU	ASP
	ALA	LYS	THR	THR	ALA	THR	GLY	GLY
	LEU	ASN	THR	SER	THR	PRO	GLU	SER
	THR	ASN	LEU	ASN	ILE	PRO	ASP	GLY
	ALA	ILE	ALA	ILE	THR	PRO	GLU	GLU
	ASP	LYS	THR	SER	ALA	ALA	ALA	GLY
	GLU	HIS	VAL	THR	PRO	PRO	ALA	PRO
	VAL	THR	THR	LYS	LYS	LYS	GLY	GLY

- Molecule 1: Microtubule-associated protein tau

Chain D:  17% 83%

[illegible]

- Molecule 1: Microtubule-associated protein tau

SER	PRO	SER	THR	HIS	SER	MET
ALA	GLY	ARG	PRO	VAL	GLU	ALA
LEU	GLY	GLN	PRO	THR	THR	THR
ALA	SS905	THR	SER	GLN	SER	PRO
LYS	GLY	ALA	GLY	ARG	ALA	GLN
GLN	RE379	PRO	GLU	MET	LYS	GLU
GLY	ASN	VAL	PRO	VAL	SER	PHE
LEU	ALA	MET	PRO	SER	THR	GLU
	LYS	PRO	LYS	LYS	THR	VAL
	ALA	ASP	GLY	LYS	ALA	GLU
	LYS	LEU	ASP	ASP	GLU	ASP
	THR	LYS	ARG	GLY	ASP	HIS
	ASP	ASN	SER	THR	VAL	ALA
	HIS	VAL	GLY	GLY	THR	GLY
	GLY	LYS	TYR	SER	ALA	THR
	ALA	SER	SER	ASP	PRO	TYR
	GLU	LYS	SER	ASP	LEU	THR
	ILE	ILE	PRO	LYS	VAL	LEU
	VAL	GLY	GLY	LYS	ASP	GLY
	TVR	SER	SER	ALA	GLU	ASP
	LYS	THR	PRO	LYS	GLY	ARG
	SER	GLU	GLY	GLY	ALA	LYS
	PRO	ASN	THR	ALA	PRO	ASP
	VAL	LEU	PRO	ASP	GLY	GLN
	VAL	LYS	GLY	GLY	LYS	GLY
	SER	HIS	SER	LYS	GLN	TVR
	GLY	GLN	ARG	THR	ALA	GLY
	ASP	PRO	SER	LYS	ALA	THR
	THR	GLY	ARG	ILE	ALA	MET
	SER	GLY	THR	ALA	GLN	HIS
	PRO	GLY	PRO	THR	PRO	GLN
	ARG	LYS	SER	PRO	HIS	ASP
	HIS	VAL	LEU	ARG	THR	GLY
	LEU	GLN	PRO	GLY	GLU	GLY
	SER	ILE	THR	ALA	ILE	LEU
	ASN	ILE	PRO	ALA	PRO	ASP
	VAL	ASN	PRO	PRO	GLU	THR
	VAL	LYS	THR	PRO	GLY	ASP
	SER	LYS	ARG	GLY	THR	ALA
	SER	LEU	GLU	GLN	THR	GLY
	GLY	ASP	PRO	LYS	ALA	LEU
	SER	LEU	LYS	GLY	GLU	LYS
	ILE	ASN	LYS	GLN	GLU	GLU
	ASP	ASN	VAL	ALA	ALA	SER
	MET	VAL	ALA	ASN	GLY	PRO
	VAL	GLN	VAL	ALA	ILE	LEU
	ASP	GLN	VAL	THR	THR	GLN
	SER	LYS	VAL	THR	ASP	LEU
	THR	LEU	VAL	THR	THR	GLY
	GLY	ASP	PRO	LYS	GLY	LEU
	SER	LEU	LYS	GLY	GLU	GLY
	ILE	ASN	VAL	GLN	GLU	SER
	ASP	ASN	VAL	ALA	ALA	PRO
	MET	VAL	ALA	ASN	GLY	GLY
	VAL	GLN	VAL	ALA	ILE	LEU
	ASP	GLN	VAL	THR	GLY	GLN
	SER	LYS	PRO	ALA	THR	ASP
	THR	LYS	LYS	LYS	SER	GLY
	GLY	ASP	THR	THR	LEU	ASP
	SER	ASN	SER	THR	GLU	GLY
	PRO	ASN	PRO	ILE	ASP	SER
	GLN	GLY	PRO	THR	THR	GLU
	LEU	SER	PRO	ALA	PRO	GLY
	ALA	LYS	LYS	LYS	LEU	ASP
	THR	ASN	SER	THR	GLU	GLY
	LEU	ASN	PRO	PRO	ASP	GLY
	ALA	ILE	SER	PRO	SER	GLU
	ASP	LYS	SER	ALA	GLU	GLY
	GLU	HIS	LYS	PRO	ALA	PRO
	VAL	VAL	LYS	THR	THR	GLY

- Chain F: 17% 83%

GLU	VAL	SER	GLY	PRO	SER	THR	HIS	SER	MET
VAL	SER	GLY	GLY	ARG	ARG	PRO	VAL	GLU	ALA
ALA	ALA	GLN	GLN	LEU	GLN	SER	GLN	THR	THR
SER	SER	GLY	THR	THR	ALA	SER	ARG	ASP	PRO
LEU	LEU	ALA	ALA	GLY	GLY	GLY	ARG	ALA	GLN
ALA	ALA	PRO	PRO	GLU	VAL	GLU	MET	LYS	GLU
LYS	LYS	VAL	VAL	VAL	VAL	PRO	VAL	THR	PHE
GLN	GLN	PRO	PRO	PRO	PRO	PRO	LYS	GLU	GLU
GLY	GLY	MET	MET	MET	LYS	LYS	LYS	THR	VAL
LEU	LEU	ASN	ASN	ASP	PRO	SER	SER	THR	MET
		LYS	LEU	ASP	GLY	GLY	LYS	ALA	GLU
		ALA	ALA	LYS	LYS	ARG	ASP	GLY	ASP
		THR	THR	ASN	ASN	SER	THR	VAL	HIS
		ASP	THR	VAL	VAL	LYS	GLY	THR	ALA
		HIS	SER	SER	SER	TYR	SER	ALA	THR
		GLY	GLY	SER	SER	SER	ASP	PRO	TYR
		ALA	ALA	ILE	ILE	PRO	LYS	VAL	GLY
		GLU	GLU	GLY	GLY	GLY	LYS	ASP	GLY
		ILE	ILE	SER	SER	SER	ALA	GLU	ASP
		VAL	VAL	THR	THR	PRO	LYS	GLY	ARG
		TYR	GLU	GLY	GLY	GLY	GLY	ALA	LYS
		LYS	ASN	ASN	THR	THR	ALA	PRO	ASP
		SER	LEU	LEU	PRO	PRO	GLY	GLY	GLN
		VAL	HIS	HIS	SER	SER	LYS	GLN	GLY
		VAL	VAL	GLN	PRO	SER	LYS	ALA	THR
		SER	SER	PRO	GLY	ARG	THR	ALA	THR
		GLY	GLY	GLY	THR	ARG	ILE	ALA	MET
		ASP	THR	GLY	GLY	THR	THR	GLN	HIS
		THR	THR	THR	THR	PRO	PRO	PRO	GLN
		VAL	VAL	ILE	ILE	PRO	ALA	ILE	GLY
		SER	SER	ASN	ASN	PRO	PRO	PRO	ASP
		ASN	ASN	LYS	LYS	THR	PRO	GLY	THR
		VAL	VAL	LYS	LYS	ARG	GLY	THR	ALA
		SER	SER	LEU	GLN	GLY	GLN	THR	GLY
		SER	SER	ASP	ASP	PRO	LYS	ALA	LEU
		THR	THR	LEU	LEU	LYS	GLY	GLU	LYS
		GLY	GLY	SER	SER	VAL	GLN	GLY	GLU
		ILE	ILE	VAL	ASN	VAL	ASN	GLY	SER
		ASP	ASP	GLN	GLN	VAL	ALA	ILE	PRO
		MET	MET	VAL	VAL	VAL	THR	GLY	GLN
		VAL	VAL	VAL	VAL	VAL	THR	ILE	THR
		THR	THR	LYS	LYS	PRO	PRO	THR	THR
		GLN	GLN	LYS	LYS	LYS	LYS	LEU	ASP
		LEU	LEU	ASP	ASP	SER	THR	GLY	GLY
		ALA	ALA	ASN	ASN	SER	PRO	ASP	SER
		THR	THR	ILE	ILE	SER	PRO	GLU	GLU
		LEU	LEU	LYS	LYS	SER	ALA	ALA	GLU
		ALA	ALA	HIS	HIS	ALA	PRO	ALA	PRO
		ASP	ASP	VAL	VAL	LYS	LYS	GLY	GLY

4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-1.26°, rise=4.74 Å, axial sym=C2	Depositor
Number of segments used	24754	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.068	Depositor
Minimum map value	-0.031	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.0119	Depositor
Map size (Å)	279.04, 279.04, 279.04	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.09, 1.09, 1.09	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/583	1.23	0/780
1	B	0.69	0/583	1.22	0/780
1	C	0.72	0/583	1.26	0/780
1	D	0.72	0/583	1.24	0/780
1	E	0.71	0/583	1.22	0/780
1	F	0.68	0/583	1.25	0/780
All	All	0.71	0/3498	1.24	0/4680

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	310	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	574	0	597	0	0
1	B	574	0	597	0	0
1	C	574	0	597	0	0
1	D	574	0	597	0	0
1	E	574	0	597	0	0
1	F	574	0	597	0	0
All	All	3444	0	3582	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/441 (17%)	68 (93%)	5 (7%)	0	100	100
1	B	73/441 (17%)	70 (96%)	3 (4%)	0	100	100
1	C	73/441 (17%)	68 (93%)	5 (7%)	0	100	100
1	D	73/441 (17%)	72 (99%)	1 (1%)	0	100	100
1	E	73/441 (17%)	67 (92%)	6 (8%)	0	100	100
1	F	73/441 (17%)	68 (93%)	5 (7%)	0	100	100
All	All	438/2646 (17%)	413 (94%)	25 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	66/358 (18%)	66 (100%)	0	100	100
1	B	66/358 (18%)	66 (100%)	0	100	100
1	C	66/358 (18%)	66 (100%)	0	100	100
1	D	66/358 (18%)	65 (98%)	1 (2%)	57	71
1	E	66/358 (18%)	66 (100%)	0	100	100
1	F	66/358 (18%)	66 (100%)	0	100	100
All	All	396/2148 (18%)	395 (100%)	1 (0%)	84	85

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	325	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	336	GLN
1	E	336	GLN
1	F	336	GLN
1	E	374	HIS
1	C	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14026. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 140




Y Index: 115




Z Index: 122


6.3.2 Raw map



X Index: 117



Y Index: 115




Z Index: 122


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 340 nm³; this corresponds to an approximate mass of 307 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.318 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.318 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.14	-	-
Author-provided FSC curve	3.14	3.67	3.19
Unmasked-calculated*	3.60	7.75	3.94

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.60 differs from the reported value 3.14 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-14026 and PDB model 7QJY. Per-residue inclusion information can be found in section [3](#) on page [5](#).


9.1 Map-model overlay [i](#)



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0119 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0119) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7760	0.5620
A	0.7790	0.5660
B	0.7750	0.5520
C	0.7840	0.5690
D	0.7750	0.5580
E	0.7710	0.5640
F	0.7730	0.5620

