Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:04 PM UTC

PDB ID	:	2QKN / pdb_00002qkn
Title	:	Crystal structure of Maize cytokinin oxidase/dehydrogenase complexed with phenylurea inhibitor CPPU
Authors	:	Briozzo, P.
Deposited on	:	2007-07-11
Resolution	:	2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	6%, 80%, 15%
2	B	2	100%
2	C	2	50%, 50%
2	D	2	50%, 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	245	A	1537[B]	-	-	X	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 4284 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytokinin dehydrogenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	0	0
			3784	2402	665	706	11			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ALA	GLY	variant	UNP Q9T0N8
A	168	THR	ASN	variant	UNP Q9T0N8
A	254	LEU	PHE	variant	UNP Q9T0N8

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is 1-(2-chloropyridin-4-yl)-3-phenylurea (CCD ID: 245) (formula: $C_{12}H_{10}ClN_3O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0	1
			17	12	1	3	1		
5	A	1	Total	C	Cl	N	O	0	1
			17	12	1	3	1		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	295	Total	O	0	0
			295	295		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytokinin dehydrogenase 1

Chain A:



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

NAG1
NAG2

4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	249.54Å 50.52Å 51.26Å 90.00° 94.01° 90.00°	Depositor
Resolution (Å)	30.00 – 2.15 30.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	10.0 (30.00-2.15) 92.3 (30.00-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.54 (at 2.08Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.191 , 0.231 0.210 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å^2)	27.6	Xtriage
Anisotropy	0.746	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 47.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4284	wwPDB-VP
Average B, all atoms (Å^2)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 245, FAD, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.40	2/3873 (0.1%)	0.90	9/5277 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	462	VAL	C-N	7.97	1.50	1.33
1	A	400	MET	SD-CE	-5.42	1.66	1.79

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	257	THR	N-CA-C	-6.47	105.13	114.12
1	A	227	GLY	N-CA-C	-5.91	107.09	115.30
1	A	250	ARG	N-CA-C	-5.89	100.23	109.25
1	A	454	ALA	N-CA-C	-5.85	99.74	109.46
1	A	168	THR	N-CA-C	-5.50	100.81	109.50
1	A	105	HIS	CA-CB-CG	-5.29	108.50	113.80
1	A	366	GLN	N-CA-C	5.14	118.13	109.95
1	A	414	VAL	N-CA-C	5.11	116.28	111.48
1	A	526	LEU	N-CA-C	5.05	117.79	109.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3784	0	3702	55	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
2	D	28	0	25	2	0
3	A	28	0	26	0	0
4	A	53	0	29	0	0
5	A	34	0	20	12	0
6	A	6	0	8	0	0
7	A	295	0	0	5	0
All	All	4284	0	3860	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1537[B]:245:H8	5:A:1537[B]:245:O1	1.79	0.80
5:A:1536[A]:245:H11	5:A:1536[A]:245:O1	1.81	0.79
5:A:1537[B]:245:O1	5:A:1537[B]:245:H11	1.81	0.79
1:A:141:TYR:CE1	1:A:239:ARG:HD2	2.17	0.79
1:A:143:ASP:OD1	1:A:239:ARG:HD3	1.83	0.79
5:A:1536[A]:245:O1	5:A:1536[A]:245:H8	1.85	0.74
1:A:463:ALA:HB3	1:A:466:ASP:HB2	1.73	0.69
1:A:141:TYR:CD1	1:A:239:ARG:HD2	2.29	0.66
1:A:524:ARG:HG3	1:A:524:ARG:HH11	1.61	0.65
1:A:47:LEU:HD22	1:A:73:LEU:HD13	1.78	0.64
1:A:492:LEU:CD1	5:A:1537[B]:245:CL7	2.83	0.64
1:A:136:SER:HB3	1:A:141:TYR:CE1	2.33	0.63
1:A:90:SER:O	1:A:92:PRO:HD3	1.98	0.63
1:A:253:ARG:HD2	1:A:306:PHE:CD2	2.33	0.63
5:A:1537[B]:245:H4	7:A:1611:HOH:O	2.00	0.62
1:A:463:ALA:HB3	1:A:466:ASP:CB	2.32	0.59
1:A:284:MET:HE3	1:A:353:VAL:HG21	1.84	0.58
1:A:48:ARG:HD3	1:A:74:TYR:CE2	2.39	0.57
5:A:1536[A]:245:H15	7:A:1611:HOH:O	2.07	0.55
5:A:1537[B]:245:O1	5:A:1537[B]:245:C11	2.50	0.53
1:A:479:ARG:HD2	7:A:1881:HOH:O	2.11	0.51
1:A:398:LEU:HG	1:A:400:MET:HE2	1.93	0.49
1:A:428:LEU:HD22	1:A:457:LEU:HD22	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:LEU:CB	1:A:47:LEU:HD23	2.42	0.49
1:A:495:HIS:HD2	7:A:1811:HOH:O	1.95	0.49
1:A:41:LEU:HB2	1:A:47:LEU:HD23	1.94	0.48
1:A:284:MET:N	1:A:284:MET:HE2	2.27	0.48
1:A:386:ASN:OD1	1:A:392:ARG:NH1	2.47	0.48
2:D:1:NAG:O3	2:D:1:NAG:H83	2.13	0.47
1:A:47:LEU:CD2	1:A:73:LEU:HD13	2.44	0.46
1:A:66:SER:HA	1:A:112:PHE:CE2	2.50	0.46
1:A:73:LEU:HG	1:A:75:PRO:HD3	1.98	0.46
1:A:164:PRO:HD2	1:A:167:TRP:CH2	2.51	0.46
1:A:338:ASN:O	1:A:339:ALA:C	2.59	0.45
5:A:1536[A]:245:O1	5:A:1536[A]:245:C8	2.51	0.45
1:A:88:ALA:HA	1:A:94:TRP:CD2	2.51	0.45
1:A:398:LEU:HD21	1:A:400:MET:HE1	1.99	0.45
1:A:401:PHE:HA	1:A:453:TYR:O	2.16	0.45
2:D:1:NAG:H61	2:D:2:NAG:HN2	1.82	0.45
1:A:336:TYR:CD2	1:A:346:VAL:HG21	2.52	0.44
1:A:136:SER:HB3	1:A:141:TYR:CD1	2.53	0.44
1:A:254:LEU:O	1:A:329:SER:HA	2.17	0.44
1:A:254:LEU:HD12	1:A:364:ALA:HA	1.99	0.44
1:A:48:ARG:HB3	1:A:50:ASP:OD1	2.17	0.44
1:A:129:ALA:HA	1:A:130:PRO:C	2.43	0.43
1:A:382:GLU:HG3	1:A:391:TRP:CZ2	2.53	0.43
1:A:252:VAL:HG22	1:A:253:ARG:N	2.34	0.43
1:A:462:VAL:HG23	1:A:463:ALA:N	2.34	0.43
1:A:492:LEU:HD11	5:A:1536[A]:245:H13	2.01	0.42
1:A:397:TRP:CZ3	5:A:1537[B]:245:CL7	3.10	0.42
1:A:427:PRO:HG2	1:A:458:LEU:HD12	2.00	0.42
1:A:516:MET:HA	1:A:516:MET:HE2	2.02	0.42
1:A:463:ALA:HB1	1:A:464:PRO:HD2	2.02	0.42
5:A:1537[B]:245:O1	5:A:1537[B]:245:C8	2.47	0.42
1:A:142:VAL:O	1:A:142:VAL:HG13	2.20	0.41
1:A:398:LEU:HG	1:A:400:MET:CE	2.50	0.41
1:A:524:ARG:HH11	1:A:524:ARG:CG	2.30	0.41
1:A:103:ARG:HA	1:A:103:ARG:HD3	1.90	0.41
1:A:214:SER:HA	1:A:222:PHE:CD1	2.55	0.41
1:A:283:PRO:HB2	1:A:284:MET:HE2	2.02	0.41
1:A:465:ASN:C	7:A:1892:HOH:O	2.63	0.41
1:A:47:LEU:HD22	1:A:73:LEU:CD1	2.49	0.41
1:A:102:GLY:O	1:A:103:ARG:NH1	2.53	0.41
1:A:318:LEU:HD21	1:A:360:VAL:HG21	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ASP:HB2	1:A:237:ARG:HB3	2.03	0.40
1:A:411:ASP:OD2	1:A:416:LYS:NZ	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	492/516 (95%)	473 (96%)	16 (3%)	3 (1%)	21 15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	PRO
1	A	182	ALA
1	A	282	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	386/395 (98%)	381 (99%)	5 (1%)	61 68

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LEU
1	A	95	PRO
1	A	216	GLN
1	A	254	LEU
1	A	419	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	120	ASN
1	A	335	ASN
1	A	379	HIS
1	A	495	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	B	1	2,1	14,14,15	0.56	0	17,19,21	0.78	1 (5%)
2	NAG	B	2	2	14,14,15	0.51	0	17,19,21	0.81	1 (5%)
2	NAG	C	1	2,1	14,14,15	0.50	0	17,19,21	0.67	0
2	NAG	C	2	2	14,14,15	0.53	0	17,19,21	0.71	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
2	NAG	D	1	2,1	14,14,15	0.58	0	17,19,21	0.69	1 (5%)
2	NAG	D	2	2	14,14,15	0.47	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	4/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	2	NAG	C2-N2-C7	-2.40	119.69	122.90
2	B	1	NAG	C2-N2-C7	-2.17	119.99	122.90
2	C	2	NAG	C2-N2-C7	-2.08	120.11	122.90
2	D	1	NAG	C2-N2-C7	-2.01	120.21	122.90

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
2	C	2	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O7-C7-N2-C2
2	C	2	NAG	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	2	0
2	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain B



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain C



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain D



Bond lengths



Bond angles



Torsions



Rings

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	245	A	1536[A]	-	18,18,18	1.54	2 (11%)	22,23,23	1.88	4 (18%)
4	FAD	A	1535	1	58,58,58	1.71	10 (17%)	85,89,89	0.92	2 (2%)
6	GOL	A	718	-	5,5,5	0.37	0	5,5,5	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	NAG	A	1608	1	14,14,15	0.49	0	17,19,21	0.67	1 (5%)
3	NAG	A	1601	1	14,14,15	0.48	0	17,19,21	0.60	0
5	245	A	1537[B]	-	18,18,18	1.54	2 (11%)	22,23,23	1.89	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	245	A	1536[A]	-	-	0/8/8/8	0/2/2/2
4	FAD	A	1535	1	-	0/34/50/50	0/6/6/6
6	GOL	A	718	-	-	0/4/4/4	-
3	NAG	A	1608	1	-	4/6/23/26	0/1/1/1
3	NAG	A	1601	1	-	0/6/23/26	0/1/1/1
5	245	A	1537[B]	-	-	0/8/8/8	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1535	FAD	C4X-N5	5.55	1.42	1.30
4	A	1535	FAD	C9-C8	5.13	1.46	1.39
4	A	1535	FAD	C8M-C8	4.17	1.58	1.51
5	A	1536[A]	245	C3-N2	-4.11	1.33	1.41
5	A	1537[B]	245	C10-N9	-4.01	1.33	1.41
5	A	1537[B]	245	C3-N2	-4.01	1.33	1.41
5	A	1536[A]	245	C10-N9	-3.98	1.33	1.41
4	A	1535	FAD	C4A-N3A	3.56	1.41	1.34
4	A	1535	FAD	C9A-C5X	3.05	1.46	1.41
4	A	1535	FAD	C9A-N10	2.99	1.46	1.41
4	A	1535	FAD	C10-N1	2.67	1.38	1.33
4	A	1535	FAD	C9-C9A	2.45	1.43	1.39
4	A	1535	FAD	C6-C5X	2.28	1.43	1.40
4	A	1535	FAD	P-O5'	2.28	1.68	1.59

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	1537[B]	245	C8-C7-N6	-5.96	120.41	125.59
5	A	1536[A]	245	C8-C7-N6	-5.93	120.43	125.59
5	A	1536[A]	245	C5-N6-C7	4.23	121.67	115.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	1537[B]	245	C5-N6-C7	4.22	121.65	115.58
4	A	1535	FAD	O2A-PA-O3P	-3.55	97.67	107.27
5	A	1536[A]	245	C4-C5-N6	-2.74	120.62	123.97
5	A	1537[B]	245	C4-C5-N6	-2.66	120.72	123.97
4	A	1535	FAD	C4X-C10-N10	2.54	120.12	116.48
5	A	1537[B]	245	CL7-C7-N6	2.16	120.03	115.98
3	A	1608	NAG	C2-N2-C7	-2.06	120.14	122.90
5	A	1537[B]	245	C3-N2-C1	-2.02	122.48	126.61
5	A	1536[A]	245	CL7-C7-N6	2.00	119.73	115.98

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1608	NAG	C8-C7-N2-C2
3	A	1608	NAG	O7-C7-N2-C2
3	A	1608	NAG	O5-C5-C6-O6
3	A	1608	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1536[A]	245	5	0
5	A	1537[B]	245	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FAD A 1535



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	496/516 (96%)	0.24	29 (5%) 29 32	17, 30, 50, 60	14 (2%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	ALA	5.5
1	A	464	PRO	5.5
1	A	129	ALA	5.4
1	A	128	ALA	5.1
1	A	465	ASN	4.3
1	A	512	ARG	3.8
1	A	383	VAL	3.8
1	A	392	ARG	3.7
1	A	210	MET	3.4
1	A	462	VAL	3.3
1	A	387	LYS	3.1
1	A	384	ALA	2.9
1	A	390	LEU	2.8
1	A	103	ARG	2.8
1	A	385	LEU	2.8
1	A	51	SER	2.7
1	A	47	LEU	2.6
1	A	33	PRO	2.6
1	A	388	LEU	2.6
1	A	367	ARG	2.6
1	A	130	PRO	2.6
1	A	273	ARG	2.4
1	A	463	ALA	2.4
1	A	381	GLU	2.2
1	A	498	ARG	2.2
1	A	389	GLY	2.1
1	A	160	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	497	ASP	2.1
1	A	391	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NAG	B	2	14/15	0.57	0.33	53,54,55,56	14
2	NAG	D	2	14/15	0.57	0.34	50,53,54,55	14
2	NAG	C	2	14/15	0.67	0.28	45,47,49,49	14
2	NAG	B	1	14/15	0.74	0.24	46,48,51,52	14
2	NAG	D	1	14/15	0.77	0.16	39,42,44,47	14
2	NAG	C	1	14/15	0.83	0.15	32,37,42,42	14

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	NAG	A	1608	14/15	0.64	0.32	54,55,57,58	14
5	245	A	1537[B]	17/17	0.66	0.15	15,20,28,29	17
5	245	A	1536[A]	17/17	0.73	0.13	14,21,29,30	17
6	GOL	A	718	6/6	0.85	0.13	33,37,41,46	0
3	NAG	A	1601	14/15	0.87	0.10	29,33,37,37	14
4	FAD	A	1535	53/53	0.97	0.06	18,23,26,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 1535:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.