



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:42 PM UTC

PDB ID : 4QKU / pdb_00004qku
Title : Crystal structure of a putative hydrolase from Burkholderia cenocepacia
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2014-06-09
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	 4% 90% 5% 5%
1	B	441	 3% 93% . .
1	C	441	 2% 92% . 5%
1	D	441	 2% 93% . .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13340 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	421	Total	C	N	O	S	0	1	0
			3140	1986	559	582	13			
1	B	427	Total	C	N	O	S	0	0	0
			3190	2017	565	595	13			
1	C	420	Total	C	N	O	S	0	0	0
			3150	1991	561	586	12			
1	D	427	Total	C	N	O	S	0	0	0
			3209	2026	570	600	13			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP B4ECX4
A	-6	ALA	-	expression tag	UNP B4ECX4
A	-5	HIS	-	expression tag	UNP B4ECX4
A	-4	HIS	-	expression tag	UNP B4ECX4
A	-3	HIS	-	expression tag	UNP B4ECX4
A	-2	HIS	-	expression tag	UNP B4ECX4
A	-1	HIS	-	expression tag	UNP B4ECX4
A	0	HIS	-	expression tag	UNP B4ECX4
B	-7	MET	-	expression tag	UNP B4ECX4
B	-6	ALA	-	expression tag	UNP B4ECX4
B	-5	HIS	-	expression tag	UNP B4ECX4
B	-4	HIS	-	expression tag	UNP B4ECX4
B	-3	HIS	-	expression tag	UNP B4ECX4
B	-2	HIS	-	expression tag	UNP B4ECX4
B	-1	HIS	-	expression tag	UNP B4ECX4
B	0	HIS	-	expression tag	UNP B4ECX4
C	-7	MET	-	expression tag	UNP B4ECX4
C	-6	ALA	-	expression tag	UNP B4ECX4
C	-5	HIS	-	expression tag	UNP B4ECX4
C	-4	HIS	-	expression tag	UNP B4ECX4
C	-3	HIS	-	expression tag	UNP B4ECX4

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP B4ECX4
C	-1	HIS	-	expression tag	UNP B4ECX4
C	0	HIS	-	expression tag	UNP B4ECX4
D	-7	MET	-	expression tag	UNP B4ECX4
D	-6	ALA	-	expression tag	UNP B4ECX4
D	-5	HIS	-	expression tag	UNP B4ECX4
D	-4	HIS	-	expression tag	UNP B4ECX4
D	-3	HIS	-	expression tag	UNP B4ECX4
D	-2	HIS	-	expression tag	UNP B4ECX4
D	-1	HIS	-	expression tag	UNP B4ECX4
D	0	HIS	-	expression tag	UNP B4ECX4

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0
2	B	1	Total Na 1 1	0	0
2	C	1	Total Na 1 1	0	0
2	D	1	Total Na 1 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	132	Total O 132 132	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	158	Total 158	O 158	0	0
5	C	167	Total 167	O 167	0	0
5	D	149	Total 149	O 149	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: hydrolase



- Molecule 1: hydrolase




- Molecule 1: hydrolase



- Molecule 1: hydrolase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.64Å 110.55Å 116.16Å 90.00° 106.61° 90.00°	Depositor
Resolution (Å)	50.00 – 2.45 50.00 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.45) 99.1 (50.00-2.45)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.208 , 0.249 0.213 , 0.253	Depositor DCC
R_{free} test set	3174 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13340	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.27 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8980e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/3220	0.82	2/4395 (0.0%)
1	B	0.74	0/3269	0.82	0/4465
1	C	0.71	0/3226	0.81	0/4402
1	D	0.74	0/3288	0.84	2/4487 (0.0%)
All	All	0.73	0/13003	0.82	4/17749 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	158	ASP	CA-C-N	5.28	124.73	119.24
1	D	158	ASP	C-N-CA	5.28	124.73	119.24
1	A	170	MET	CG-SD-CE	5.22	112.38	100.90
1	A	112	ARG	CG-CD-NE	-5.02	100.96	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3140	0	3027	16	0
1	B	3190	0	3075	11	0
1	C	3150	0	3060	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3209	0	3105	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	4	0	6	0	0
3	B	4	0	6	0	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
4	A	5	0	0	0	0
4	B	5	0	0	1	0
4	C	10	0	0	0	0
4	D	5	0	0	0	0
5	A	132	0	0	2	0
5	B	158	0	0	0	0
5	C	167	0	0	0	0
5	D	149	0	0	1	0
All	All	13340	0	12291	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:ILE:CG2	1:B:262:LEU:HD12	2.32	0.60
1:A:18:TRP:CD2	1:A:112:ARG:HB2	2.37	0.59
1:D:45:ALA:HB1	5:D:656:HOH:O	2.06	0.56
1:C:262:LEU:HD12	1:D:172:ILE:CG2	2.36	0.55
1:A:170:MET:HE2	1:A:347:HIS:HE1	1.71	0.55
1:A:262:LEU:HD12	1:B:172:ILE:CG2	2.39	0.53
1:B:218:GLY:HA2	1:B:361:MET:O	2.10	0.52
1:D:407:LEU:HB2	1:D:423:CYS:HB3	1.93	0.50
1:B:148:HIS:NE2	4:B:503:SO4:O2	2.39	0.50
1:B:407:LEU:HB2	1:B:423:CYS:HB3	1.94	0.50
1:A:407:LEU:HB2	1:A:423:CYS:HB3	1.93	0.49
1:D:218:GLY:HA2	1:D:361:MET:O	2.13	0.49
1:A:218:GLY:HA2	1:A:361:MET:O	2.13	0.49
1:A:146:LYS:NZ	5:A:688:HOH:O	2.46	0.48
1:C:407:LEU:HB2	1:C:423:CYS:HB3	1.94	0.48
1:C:172:ILE:CG2	1:D:262:LEU:HD12	2.43	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:170:MET:HE1	1:D:257:TRP:CD1	2.48	0.48
1:C:218:GLY:HA2	1:C:361:MET:O	2.14	0.48
1:A:170:MET:HE2	1:A:347:HIS:CE1	2.50	0.47
1:A:432:SER:HB2	1:C:90:ARG:NH2	2.30	0.47
1:A:431:ARG:NH2	5:A:641:HOH:O	2.45	0.46
1:C:213:ILE:HG23	1:C:399:ILE:HD11	1.98	0.46
1:D:213:ILE:HG23	1:D:399:ILE:HD11	1.97	0.46
1:A:172:ILE:HG23	1:B:262:LEU:HD12	1.98	0.45
1:B:6:ASP:OD1	1:B:7:TRP:N	2.49	0.45
1:A:213:ILE:HG23	1:A:399:ILE:HD11	1.98	0.44
1:B:213:ILE:HG23	1:B:399:ILE:HD11	1.98	0.44
1:D:381:THR:HG22	1:D:388:VAL:HG23	2.01	0.42
1:C:262:LEU:HD12	1:D:172:ILE:HG23	1.99	0.42
1:B:182:VAL:O	1:B:273:THR:HA	2.20	0.42
1:B:381:THR:HG22	1:B:388:VAL:HG23	2.02	0.42
1:A:106:ARG:O	1:A:112:ARG:NH1	2.51	0.41
1:A:182:VAL:O	1:A:273:THR:HA	2.20	0.41
1:C:381:THR:HG22	1:C:388:VAL:HG23	2.02	0.41
1:C:182:VAL:O	1:C:273:THR:HA	2.21	0.41
1:B:213:ILE:HG22	1:B:247:ASN:HD21	1.86	0.41
1:D:182:VAL:O	1:D:273:THR:HA	2.22	0.40
1:A:255:GLN:N	1:A:267[B]:SER:OG	2.54	0.40
1:A:381:THR:HG22	1:A:388:VAL:HG23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	418/441 (95%)	407 (97%)	11 (3%)	0	100	100
1	B	425/441 (96%)	412 (97%)	13 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	416/441 (94%)	404 (97%)	12 (3%)	0	100	100
1	D	425/441 (96%)	413 (97%)	12 (3%)	0	100	100
All	All	1684/1764 (96%)	1636 (97%)	48 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/347 (90%)	307 (98%)	6 (2%)	50	64
1	B	320/347 (92%)	318 (99%)	2 (1%)	78	86
1	C	318/347 (92%)	316 (99%)	2 (1%)	78	86
1	D	324/347 (93%)	322 (99%)	2 (1%)	78	86
All	All	1275/1388 (92%)	1263 (99%)	12 (1%)	70	81

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ARG
1	A	154	SER
1	A	156	PHE
1	A	170	MET
1	A	213	ILE
1	A	304	HIS
1	B	213	ILE
1	B	304	HIS
1	C	213	ILE
1	C	304	HIS
1	D	213	ILE
1	D	304	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	194	GLN
1	D	194	GLN
1	D	344	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	C	504	-	4,4,4	0.53	0	6,6,6	0.20	0
4	SO4	D	503	-	4,4,4	0.58	0	6,6,6	0.33	0
4	SO4	B	503	-	4,4,4	0.39	0	6,6,6	0.30	0
3	EDO	D	502	-	3,3,3	0.67	0	2,2,2	0.21	0
3	EDO	A	502	-	3,3,3	0.57	0	2,2,2	0.04	0
4	SO4	A	503	-	4,4,4	0.44	0	6,6,6	0.57	0
4	SO4	C	503	-	4,4,4	0.42	0	6,6,6	0.52	0
3	EDO	B	502	-	3,3,3	0.37	0	2,2,2	0.31	0
3	EDO	C	502	-	3,3,3	0.63	0	2,2,2	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	502	-	-	0/1/1/1	-
3	EDO	B	502	-	-	1/1/1/1	-
3	EDO	C	502	-	-	0/1/1/1	-
3	EDO	D	502	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	503	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	421/441 (95%)	0.42	19 (4%)	38 37	12, 24, 53, 73	1 (0%)
1	B	427/441 (96%)	0.19	14 (3%)	49 50	10, 20, 37, 61	0
1	C	420/441 (95%)	0.13	7 (1%)	69 71	11, 21, 38, 55	0
1	D	427/441 (96%)	0.17	9 (2%)	63 65	9, 20, 37, 64	0
All	All	1695/1764 (96%)	0.23	49 (2%)	53 55	9, 21, 43, 73	1 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	42	ALA	4.7
1	A	155	MET	4.6
1	B	414	ASP	4.6
1	D	305	ALA	4.5
1	D	43	LYS	4.5
1	A	156	PHE	4.2
1	A	107	ASP	3.8
1	C	161	ASN	3.7
1	C	154	SER	3.6
1	D	433	ALA	3.5
1	D	384	GLY	3.4
1	A	199	ASP	3.3
1	B	198	PRO	3.2
1	A	305	ALA	3.2
1	B	199	ASP	3.2
1	D	382	TRP	3.2
1	B	415	GLY	3.1
1	A	111	LEU	2.9
1	A	71	PRO	2.9
1	A	72	ALA	2.8
1	C	162	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	156	PHE	2.8
1	A	306	GLY	2.8
1	B	260	VAL	2.7
1	B	413	GLY	2.7
1	B	197	LEU	2.7
1	A	162	ALA	2.6
1	D	72	ALA	2.6
1	A	154	SER	2.5
1	B	416	TYR	2.4
1	D	44	GLY	2.4
1	A	382	TRP	2.4
1	A	74	ALA	2.4
1	C	382	TRP	2.3
1	A	200	GLN	2.2
1	C	119	ALA	2.2
1	D	198	PRO	2.2
1	A	432	SER	2.1
1	B	43	LYS	2.1
1	B	200	GLN	2.1
1	C	147	GLU	2.1
1	B	78	ALA	2.1
1	A	108	ASP	2.1
1	A	113	ASP	2.1
1	C	153	GLY	2.0
1	B	159	PRO	2.0
1	A	106	ARG	2.0
1	A	117	LEU	2.0
1	B	160	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	D	503	5/5	0.81	0.20	64,64,65,66	0
4	SO4	C	504	5/5	0.82	0.20	64,70,72,77	0
4	SO4	A	503	5/5	0.89	0.14	54,60,61,64	0
3	EDO	D	502	4/4	0.90	0.13	21,24,26,27	0
4	SO4	C	503	5/5	0.91	0.15	45,47,49,52	0
3	EDO	C	502	4/4	0.94	0.07	13,13,14,15	0
2	NA	A	501	1/1	0.94	0.13	16,16,16,16	0
2	NA	B	501	1/1	0.94	0.05	15,15,15,15	0
4	SO4	B	503	5/5	0.95	0.14	44,45,48,49	0
3	EDO	A	502	4/4	0.96	0.07	15,16,16,17	0
3	EDO	B	502	4/4	0.97	0.06	17,18,18,18	0
2	NA	D	501	1/1	0.97	0.03	21,21,21,21	0
2	NA	C	501	1/1	0.98	0.04	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.