



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 09:51 PM UTC

PDB ID : 5QK1 / pdb_00005qk1
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z2027049478
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.
Deposited on : 2018-10-31
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	10% 80% 10% • 7%
1	B	209	11% 80% 12% • 7%
1	C	209	18% 78% 13% • 8%
1	D	209	25% 72% 18% • 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1462	925	245	284	8			
1	B	194	Total	C	N	O	S	0	0	0
			1464	924	244	288	8			
1	C	192	Total	C	N	O	S	0	0	0
			1425	898	237	283	7			
1	D	190	Total	C	N	O	S	0	1	0
			1433	909	242	274	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is 5-(1,3-thiazol-2-yl)-1H-1,2,4-triazole (CCD ID: K34) (formula: C₅H₄N₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N S 10 5 4 1	0	0
4	B	1	Total C N S 10 5 4 1	0	0
4	C	1	Total C N S 10 5 4 1	0	0
4	D	1	Total C N S 10 5 4 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	84	Total	O	0	0
			84	84		
6	B	79	Total	O	0	0
			79	79		
6	C	77	Total	O	0	0
			77	77		
6	D	35	Total	O	0	0
			35	35		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.30Å 59.89Å 80.36Å 79.21° 81.58° 75.63°	Depositor
Resolution (Å)	78.50 – 1.49 78.50 – 1.49	Depositor EDS
% Data completeness (in resolution range)	93.5 (78.50-1.49) 93.5 (78.50-1.49)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 1.49Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.218 , 0.252 0.231 , 0.264	Depositor DCC
R_{free} test set	7183 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6128	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG, EDO, K34

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.43	5/1490 (0.3%)	1.24	6/2032 (0.3%)
1	B	1.42	6/1492 (0.4%)	1.23	4/2035 (0.2%)
1	C	1.44	5/1452 (0.3%)	1.21	1/1984 (0.1%)
1	D	1.48	10/1461 (0.7%)	1.29	8/1992 (0.4%)
All	All	1.44	26/5895 (0.4%)	1.24	19/8043 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	84	ARG	C-O	-9.77	1.15	1.24
1	D	17	ILE	C-O	-6.95	1.16	1.24
1	B	127	SER	C-O	-6.65	1.15	1.23
1	D	80	VAL	CA-C	-6.08	1.45	1.52
1	D	182	LEU	CA-C	-6.08	1.45	1.52
1	D	68	LEU	N-CA	5.97	1.53	1.46
1	C	104	THR	CA-C	5.92	1.60	1.53
1	A	174	PRO	CA-C	-5.81	1.45	1.52
1	C	91	CYS	CA-C	-5.72	1.45	1.52
1	D	175	LYS	CA-C	5.71	1.60	1.52
1	C	136	LEU	CA-C	-5.50	1.45	1.52
1	D	26	GLY	N-CA	5.49	1.48	1.44
1	D	85	PRO	CA-C	-5.47	1.47	1.52
1	D	180	GLN	C-O	-5.47	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	92	ILE	C-O	-5.47	1.18	1.24
1	A	124	ALA	CA-C	-5.40	1.46	1.52
1	B	35	THR	C-O	-5.38	1.17	1.23
1	A	183	ASP	N-CA	5.32	1.52	1.46
1	B	80	VAL	N-CA	-5.32	1.40	1.46
1	C	79	LEU	CA-CB	-5.32	1.45	1.53
1	A	104	THR	CB-CG2	-5.25	1.35	1.52
1	D	95	PRO	N-CA	5.19	1.53	1.47
1	B	196	ARG	NE-CZ	5.16	1.38	1.33
1	B	201	ALA	CA-CB	-5.13	1.45	1.53
1	A	43	THR	CA-C	5.10	1.59	1.52
1	B	120	LYS	CA-CB	5.03	1.60	1.53

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	206	HIS	N-CA-C	7.01	121.10	112.54
1	D	85	PRO	CA-C-N	-6.78	112.01	119.19
1	D	85	PRO	C-N-CA	-6.78	112.01	119.19
1	D	85	PRO	N-CA-C	6.49	118.62	110.70
1	B	38	ASP	CA-C-N	-6.10	113.45	120.04
1	B	38	ASP	C-N-CA	-6.10	113.45	120.04
1	D	94	PHE	N-CA-C	-5.69	102.23	110.08
1	A	188	GLU	N-CA-C	5.62	120.26	112.90
1	D	74	TYR	N-CA-C	-5.56	103.05	110.55
1	D	178	LEU	N-CA-C	5.53	117.31	111.28
1	D	132	MET	N-CA-C	5.43	118.37	111.69
1	A	73	HIS	CA-C-O	5.27	127.06	121.11
1	B	114	GLU	N-CA-C	5.25	116.68	111.07
1	C	85	PRO	N-CA-C	5.21	117.05	110.70
1	A	70	ARG	NE-CZ-NH2	-5.12	114.60	119.20
1	B	196	ARG	CG-CD-NE	-5.07	100.84	112.00
1	A	104	THR	CA-C-N	-5.04	113.46	119.05
1	A	104	THR	C-N-CA	-5.04	113.46	119.05
1	A	186	VAL	N-CA-C	-5.03	105.80	110.53

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	52	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1462	0	1433	18	0
1	B	1464	0	1414	15	0
1	C	1425	0	1347	16	0
1	D	1433	0	1397	18	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
4	C	10	0	0	0	0
4	D	10	0	0	0	0
5	A	4	0	6	0	0
5	B	4	0	6	0	0
5	C	8	0	12	0	0
5	D	4	0	6	0	0
6	A	84	0	0	0	0
6	B	79	0	0	0	0
6	C	77	0	0	5	0
6	D	35	0	0	2	0
All	All	6128	0	5621	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.53	0.90
1:C:125:GLU:OE1	6:C:401:HOH:O	1.95	0.84
1:C:104:THR:HG23	1:C:107:ALA:H	1.43	0.83
1:D:188:GLU:HG2	1:D:189:GLU:HG3	1.71	0.73
1:D:120:LYS:H	1:D:155:ASN:HD21	1.36	0.73
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.73	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:LEU:C	1:A:191:LEU:HD23	2.17	0.69
1:A:104:THR:HG23	1:A:107:ALA:H	1.59	0.67
1:D:194:ASP:OD1	1:D:196[A]:ARG:HD3	1.97	0.65
1:C:51:ARG:NE	6:C:402:HOH:O	2.30	0.64
6:C:402:HOH:O	1:D:136:LEU:HD22	1.97	0.64
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.85	0.58
6:C:401:HOH:O	1:D:206:HIS:HE1	1.86	0.58
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.35	0.57
1:C:206:HIS:HE1	1:D:125:GLU:OE1	1.89	0.56
1:A:125:GLU:HG3	1:A:145:THR:HB	1.87	0.56
1:B:104:THR:HG23	1:B:107:ALA:H	1.70	0.56
1:D:104:THR:HG23	1:D:107:ALA:H	1.71	0.56
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.88	0.55
1:D:150:GLY:O	6:D:401:HOH:O	2.18	0.55
1:C:200:TYR:OH	1:D:206:HIS:HD2	1.90	0.54
1:D:71:THR:O	1:D:73:HIS:ND1	2.40	0.54
1:A:200:TYR:OH	1:B:206:HIS:HD2	1.91	0.53
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.44	0.52
1:B:92:ILE:CD1	1:B:191:LEU:HD13	2.32	0.52
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.44	0.51
1:C:206:HIS:HD2	1:D:200:TYR:OH	1.94	0.51
1:A:132:MET:HE1	1:A:196:ARG:CZ	2.42	0.50
1:A:191:LEU:C	1:A:191:LEU:CD2	2.85	0.49
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.48	0.49
1:C:15:GLN:N	6:C:406:HOH:O	2.45	0.48
1:A:206:HIS:HE1	1:B:125:GLU:OE1	1.97	0.48
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.97	0.47
1:A:104:THR:CG2	1:A:107:ALA:H	2.25	0.47
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.50	0.46
1:B:23:ILE:HD12	1:B:31:LEU:HD13	1.98	0.46
1:A:167:PHE:CD2	1:B:39:PRO:HG2	2.51	0.45
1:A:71:THR:O	1:A:73:HIS:HD2	2.00	0.45
1:C:92:ILE:HD12	1:C:191:LEU:HD11	1.99	0.45
1:D:191:LEU:C	1:D:191:LEU:HD12	2.41	0.45
1:C:152:ASP:OD1	1:C:154:GLU:N	2.46	0.44
1:A:71:THR:HG23	1:A:151:ASP:OD2	2.19	0.43
1:C:81:LYS:HA	1:C:91:CYS:O	2.20	0.42
1:B:82:GLN:HG2	1:B:168:VAL:HG22	2.01	0.42
1:C:127:SER:HB2	1:C:128:PRO:HD2	2.01	0.42
1:A:54:ARG:HD3	1:A:60:ASP:OD1	2.20	0.42
1:D:181:ARG:HD3	6:D:405:HOH:O	2.18	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:149:ASN:C	1:D:149:ASN:HD22	2.28	0.41
1:B:92:ILE:HD11	1:B:191:LEU:CD1	2.36	0.41
1:D:49:VAL:O	1:D:137:SER:HA	2.21	0.41
1:B:58:THR:CB	1:B:142:HIS:NE2	2.84	0.40
1:A:38:ASP:OD1	1:A:38:ASP:C	2.64	0.40
1:A:203:ALA:HB2	1:B:200:TYR:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	187 (97%)	4 (2%)	1 (0%)	24	8
1	B	190/209 (91%)	186 (98%)	3 (2%)	1 (0%)	24	8
1	C	188/209 (90%)	183 (97%)	4 (2%)	1 (0%)	24	8
1	D	187/209 (90%)	184 (98%)	3 (2%)	0	100	100
All	All	757/836 (91%)	740 (98%)	14 (2%)	3 (0%)	30	12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	PRO
1	C	66	PRO
1	B	66	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/179 (86%)	148 (96%)	6 (4%)	28	5
1	B	154/179 (86%)	151 (98%)	3 (2%)	50	22
1	C	146/179 (82%)	140 (96%)	6 (4%)	27	5
1	D	149/179 (83%)	144 (97%)	5 (3%)	32	7
All	All	603/716 (84%)	583 (97%)	20 (3%)	33	8

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	27	LYS
1	A	43	THR
1	A	70	ARG
1	A	125	GLU
1	A	191	LEU
1	B	20	GLU
1	B	31	LEU
1	B	43	THR
1	C	31	LEU
1	C	43	THR
1	C	45	THR
1	C	71	THR
1	C	164	ASP
1	C	189	GLU
1	D	38	ASP
1	D	43	THR
1	D	72	LEU
1	D	168	VAL
1	D	188	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	73	HIS
1	A	180	GLN
1	A	206	HIS
1	B	82	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	190	HIS
1	B	206	HIS
1	B	208	ASN
1	C	15	GLN
1	C	206	HIS
1	D	15	GLN
1	D	149	ASN
1	D	155	ASN
1	D	190	HIS
1	D	206	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 9 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	K34	D	303	-	10,11,11	1.52	3 (30%)	10,14,14	1.73	3 (30%)
4	K34	A	304	-	10,11,11	1.73	4 (40%)	10,14,14	1.71	2 (20%)
5	EDO	A	305	-	3,3,3	0.66	0	2,2,2	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	C	305	-	3,3,3	0.53	0	2,2,2	0.79	0
4	K34	C	304	-	10,11,11	1.76	4 (40%)	10,14,14	1.81	3 (30%)
5	EDO	C	301	-	3,3,3	0.87	0	2,2,2	0.40	0
5	EDO	D	304	-	3,3,3	0.59	0	2,2,2	0.22	0
5	EDO	B	301	-	3,3,3	0.86	0	2,2,2	0.29	0
4	K34	B	304	-	10,11,11	1.39	2 (20%)	10,14,14	1.94	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	K34	D	303	-	-	0/1/4/4	0/2/2/2
4	K34	A	304	-	-	0/1/4/4	0/2/2/2
5	EDO	A	305	-	-	0/1/1/1	-
5	EDO	C	305	-	-	0/1/1/1	-
4	K34	C	304	-	-	0/1/4/4	0/2/2/2
5	EDO	C	301	-	-	1/1/1/1	-
5	EDO	D	304	-	-	0/1/1/1	-
5	EDO	B	301	-	-	0/1/1/1	-
4	K34	B	304	-	-	0/1/4/4	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	304	K34	C-N	2.78	1.43	1.38
4	A	304	K34	C3-N1	2.71	1.36	1.34
4	C	304	K34	C3-N3	2.69	1.40	1.33
4	D	303	K34	C3-N3	2.61	1.40	1.33
4	C	304	K34	C3-N1	2.60	1.36	1.34
4	B	304	K34	C-N	2.41	1.42	1.38
4	B	304	K34	N1-N2	-2.40	1.32	1.36
4	A	304	K34	C3-N3	2.24	1.39	1.33
4	C	304	K34	C4-N2	2.22	1.37	1.32
4	D	303	K34	C-N	2.20	1.42	1.38
4	D	303	K34	C4-N3	2.16	1.40	1.36
4	C	304	K34	C4-N3	2.16	1.40	1.36
4	A	304	K34	N1-N2	-2.05	1.33	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	304	K34	C3-N1-N2	-3.69	106.80	110.26
4	B	304	K34	C3-N1-N2	-3.16	107.29	110.26
4	C	304	K34	C3-N1-N2	-3.08	107.36	110.26
4	B	304	K34	C2-C3-N3	-2.97	123.42	126.49
4	C	304	K34	C2-C3-N3	-2.85	123.54	126.49
4	D	303	K34	C2-C3-N3	-2.77	123.63	126.49
4	D	303	K34	N2-C4-N3	-2.74	107.47	113.49
4	B	304	K34	N2-C4-N3	-2.64	107.70	113.49
4	D	303	K34	C3-N1-N2	-2.62	107.80	110.26
4	A	304	K34	N2-C4-N3	-2.59	107.81	113.49
4	C	304	K34	N2-C4-N3	-2.47	108.06	113.49
4	B	304	K34	S-C2-N	2.07	116.33	114.37

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	0.66	21 (10%)	11 11	16, 28, 54, 67	12 (6%)
1	B	194/209 (92%)	0.72	22 (11%)	10 10	20, 32, 59, 73	2 (1%)
1	C	192/209 (91%)	1.03	38 (19%)	3 2	21, 34, 59, 68	8 (4%)
1	D	190/209 (90%)	1.47	53 (27%)	1 1	12, 37, 62, 77	9 (4%)
All	All	770/836 (92%)	0.97	134 (17%)	4 3	12, 33, 59, 77	31 (4%)

All (134) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	153	ALA	8.4
1	B	208	ASN	5.9
1	D	154	GLU	5.8
1	D	74	TYR	5.6
1	C	23	ILE	5.5
1	D	28	TRP	5.5
1	A	191	LEU	5.3
1	A	28	TRP	5.1
1	A	57	GLN	5.1
1	A	167	PHE	5.0
1	D	29	VAL	4.8
1	A	56	GLU	4.8
1	D	16	TYR	4.7
1	B	28	TRP	4.5
1	C	18	ILE	4.4
1	B	46	TRP	4.3
1	C	16	TYR	4.0
1	D	14	LYS	3.9
1	D	23	ILE	3.9
1	D	40	THR	3.9
1	A	189	GLU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	191	LEU	3.7
1	C	17	ILE	3.7
1	D	167	PHE	3.7
1	C	40	THR	3.6
1	A	190	HIS	3.6
1	D	43	THR	3.5
1	D	158	PRO	3.5
1	C	190	HIS	3.4
1	C	43	THR	3.4
1	D	102	GLY	3.4
1	C	28	TRP	3.3
1	C	58	THR	3.3
1	C	39	PRO	3.3
1	C	207	ALA	3.3
1	B	167	PHE	3.2
1	C	104	THR	3.2
1	D	68	LEU	3.2
1	A	158	PRO	3.1
1	A	157	ARG	3.1
1	D	156	ALA	3.1
1	C	35	THR	3.1
1	A	72	LEU	3.1
1	A	163	GLY	3.1
1	B	176	ASN	3.1
1	D	97	GLY	3.1
1	C	153	ALA	3.1
1	B	40	THR	3.0
1	D	22	LEU	3.0
1	D	118	GLY	3.0
1	C	161	LYS	3.0
1	D	100	ASP	3.0
1	D	148	ILE	2.9
1	D	53	THR	2.9
1	D	41	GLY	2.9
1	D	163	GLY	2.9
1	B	22	LEU	2.9
1	D	152	ASP	2.9
1	C	22	LEU	2.9
1	A	14	LYS	2.9
1	B	58	THR	2.8
1	D	159	LYS	2.8
1	D	164	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	164	ASP	2.8
1	D	119	TYR	2.8
1	B	14	LYS	2.8
1	A	187	ALA	2.7
1	B	163	GLY	2.7
1	C	136	LEU	2.7
1	C	184	ALA	2.7
1	B	160	PRO	2.7
1	D	31	LEU	2.7
1	D	88	GLY	2.7
1	C	168	VAL	2.7
1	B	104	THR	2.7
1	A	27	LYS	2.6
1	C	45	THR	2.6
1	D	101	ASP	2.6
1	B	162	PRO	2.6
1	C	57	GLN	2.6
1	D	99	ILE	2.6
1	D	36	TYR	2.6
1	D	162	PRO	2.6
1	A	40	THR	2.5
1	D	90	TYR	2.5
1	C	167	PHE	2.5
1	C	162	PRO	2.5
1	A	23	ILE	2.5
1	D	166	GLU	2.5
1	C	191	LEU	2.5
1	D	208	ASN	2.5
1	D	165	GLY	2.4
1	D	70	ARG	2.4
1	D	196[A]	ARG	2.4
1	A	165	GLY	2.4
1	B	165	GLY	2.4
1	C	102	GLY	2.4
1	C	36	TYR	2.4
1	C	29	VAL	2.4
1	B	159	LYS	2.3
1	C	27	LYS	2.3
1	A	58	THR	2.3
1	C	49	VAL	2.3
1	D	34	THR	2.3
1	D	104	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	31	LEU	2.3
1	C	74	TYR	2.3
1	C	187	ALA	2.2
1	B	18	ILE	2.2
1	D	171	ILE	2.2
1	A	162	PRO	2.2
1	B	102	GLY	2.2
1	B	190	HIS	2.2
1	A	160	PRO	2.2
1	D	27	LYS	2.2
1	C	165	GLY	2.2
1	D	98	LEU	2.1
1	C	30	LYS	2.1
1	C	160	PRO	2.1
1	C	172	SER	2.1
1	D	19	SER	2.1
1	D	135	GLY	2.1
1	C	72	LEU	2.1
1	D	173	LEU	2.1
1	C	52	THR	2.1
1	B	168	VAL	2.1
1	B	158	PRO	2.1
1	D	37	MET	2.1
1	D	160	PRO	2.1
1	D	96	ALA	2.1
1	D	18	ILE	2.1
1	D	73	HIS	2.1
1	B	31	LEU	2.0
1	D	110	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	B	301	4/4	0.71	0.18	42,48,53,54	0
5	EDO	C	301	4/4	0.81	0.15	37,45,48,57	0
3	CL	A	303	1/1	0.83	0.17	68,68,68,68	0
2	MG	A	302	1/1	0.85	0.13	65,65,65,65	0
2	MG	C	302	1/1	0.87	0.20	48,48,48,48	0
4	K34	C	304	10/10	0.87	0.13	40,44,54,55	10
4	K34	D	303	10/10	0.88	0.15	37,44,51,57	10
2	MG	D	301	1/1	0.89	0.09	32,32,32,32	0
4	K34	A	304	10/10	0.89	0.12	33,37,42,46	10
2	MG	B	303	1/1	0.91	0.09	62,62,62,62	0
5	EDO	C	305	4/4	0.91	0.12	37,38,39,39	4
4	K34	B	304	10/10	0.92	0.13	30,32,37,40	10
5	EDO	D	304	4/4	0.93	0.09	33,34,35,35	4
2	MG	D	302	1/1	0.94	0.18	43,43,43,43	0
5	EDO	A	305	4/4	0.95	0.08	26,27,28,28	0
2	MG	C	303	1/1	0.98	0.04	33,33,33,33	0
2	MG	B	302	1/1	0.99	0.04	30,30,30,30	0
2	MG	A	301	1/1	0.99	0.03	27,27,27,27	0

6.5 Other polymers

There are no such residues in this entry.