



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 03:58 PM UTC

PDB ID : 1QLE / pdb_00001qle
Title : CRYO-STRUCTURE OF THE PARACOCCLUS DENITRIFICANS FOUR-SUBUNIT CYTOCHROME C OXIDASE IN THE COMPLETELY OXIDIZED STATE COMPLEXED WITH AN ANTIBODY FV FRAGMENT
Authors : Harrenga, A.; Michel, H.
Deposited on : 1999-08-30
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	538	32% (green), 58% (yellow), 9% (orange), 1% (red), 1% (grey)
2	B	252	45% (green), 45% (yellow), 9% (orange), 1% (red), 1% (grey)
3	C	273	44% (green), 47% (yellow), 10% (orange), 1% (red), 1% (grey)
4	D	43	63% (green), 35% (yellow), 1% (orange), 1% (red), 1% (grey)
5	H	119	29% (green), 62% (yellow), 8% (orange), 1% (red), 1% (grey)
6	L	108	36% (green), 50% (yellow), 14% (orange), 1% (red), 1% (grey)

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 10762 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE I-BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	538	4269	2860	671	705	33	0	0	0

- Molecule 2 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	252	1977	1295	319	355	8	0	0	0

- Molecule 3 is a protein called CYTOCHROME C OXIDASE POLYPEPTIDE III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	273	2181	1483	339	348	11	0	0	0

- Molecule 4 is a protein called CCYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	43	332	214	58	59	1	0	0	0


- Molecule 5 is a protein called HEAVY CHAIN ANTIBODY FV FRAGMENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	H	119	938	589	157	186	6	0	0	0

- Molecule 6 is a protein called LIGHT CHAIN ANTIBODY FV FRAGMENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	L	108	832	530	135	165	2	0	0	0

- Molecule 7 is HEME-A (CCD ID: HEA) (formula: $C_{49}H_{56}FeN_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
7	A	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		
7	A	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		

- Molecule 8 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cu	0	0
			1	1		


- Molecule 9 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Ca	0	0
			1	1		

- Molecule 10 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total	Mn	0	0
			1	1		

- Molecule 11 is DINUCLEAR COPPER ION (CCD ID: CUA) (formula: Cu_2).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	1	Total	Cu	0	0
			2	2		

- Molecule 12 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: $C_{44}H_{88}NO_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	C	1	Total	C	N	O	P	0	0
			54	44	1	8	1		
12	C	1	Total	C	N	O	P	0	0
			54	44	1	8	1		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: CYTOCHROME C OXIDASE POLYPEPTIDE I-BETA




- Molecule 2: CYTOCHROME C OXIDASE POLYPEPTIDE II





- Molecule 3: CYTOCHROME C OXIDASE POLYPEPTIDE III

Chain C: 44% 47% 10%




- Molecule 4: CCYTOCHROME C OXIDASE

Chain D: 63% 35%




- Molecule 5: HEAVY CHAIN ANTIBODY FV FRAGMENT

Chain H: 29% 62% 8%



- Molecule 6: LIGHT CHAIN ANTIBODY FV FRAGMENT

Chain L: 36% 50% 14%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 4	Depositor
Cell constants a, b, c, α , β , γ	205.20Å 205.20Å 81.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 3.00	Depositor
% Data completeness (in resolution range)	66.2 (6.00-3.00)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 0.3	Depositor
R, R_{free}	0.235 , 0.309	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	10762	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PC1, HEA, CU, MN, CUA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	3/4428 (0.1%)	1.12	20/6042 (0.3%)
2	B	0.71	1/2034 (0.0%)	1.13	7/2787 (0.3%)
3	C	0.66	0/2267	1.05	5/3103 (0.2%)
4	D	0.62	0/339	0.99	0/457
5	H	0.70	0/960	0.95	0/1298
6	L	0.65	0/853	1.05	6/1156 (0.5%)
All	All	0.70	4/10881 (0.0%)	1.08	38/14843 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	200	ILE	CA-CB	5.68	1.60	1.54
2	B	203	TRP	NE1-CE2	-5.41	1.31	1.37
1	A	443	TRP	NE1-CE2	-5.27	1.31	1.37
1	A	323	TRP	NE1-CE2	-5.09	1.31	1.37

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	195	VAL	N-CA-C	8.73	116.19	107.55
1	A	257	ASP	CA-C-N	8.12	127.68	119.24
1	A	257	ASP	C-N-CA	8.12	127.68	119.24
1	A	528	ASP	N-CA-C	6.93	119.68	111.71
3	C	238	GLN	N-CA-C	6.75	118.72	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4269	0	4184	396	0
2	B	1977	0	1963	151	0
3	C	2181	0	2151	184	0
4	D	332	0	331	12	0
5	H	938	0	894	75	0
6	L	832	0	807	56	0
7	A	120	0	108	28	0
8	A	1	0	0	0	0
9	A	1	0	0	0	0
10	A	1	0	0	0	0
11	B	2	0	0	0	0
12	C	108	0	176	26	0
All	All	10762	0	10614	817	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 817 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4:LEU:HD11	2:B:242:TRP:CD1	1.25	1.68
2:B:4:LEU:CD1	2:B:242:TRP:CD1	2.15	1.27
3:C:2:HIS:O	3:C:3:VAL:HG22	1.13	1.25
5:H:6:GLU:OE1	5:H:96:CYS:N	1.66	1.25
2:B:159:ASP:O	2:B:161:PRO:HD3	1.37	1.22

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	536/538 (100%)	447 (83%)	71 (13%)	18 (3%)	3	17
2	B	250/252 (99%)	198 (79%)	47 (19%)	5 (2%)	6	28
3	C	271/273 (99%)	215 (79%)	51 (19%)	5 (2%)	6	31
4	D	41/43 (95%)	33 (80%)	5 (12%)	3 (7%)	1	4
5	H	117/119 (98%)	99 (85%)	16 (14%)	2 (2%)	7	32
6	L	106/108 (98%)	92 (87%)	13 (12%)	1 (1%)	14	48
All	All	1321/1333 (99%)	1084 (82%)	203 (15%)	34 (3%)	4	23

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	PHE
1	A	213	THR
1	A	442	GLU
1	A	19	PHE
1	A	102	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/441 (100%)	390 (88%)	51 (12%)	5	23
2	B	211/211 (100%)	183 (87%)	28 (13%)	4	18
3	C	220/220 (100%)	194 (88%)	26 (12%)	5	22
4	D	34/34 (100%)	28 (82%)	6 (18%)	2	10
5	H	101/101 (100%)	87 (86%)	14 (14%)	3	17
6	L	92/92 (100%)	78 (85%)	14 (15%)	3	14
All	All	1099/1099 (100%)	960 (87%)	139 (13%)	4	20

5 of 139 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	H	44	ARG
5	H	82	GLN
6	L	42	LYS
1	A	495	ILE
1	A	491	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	111	ASN
3	C	159	HIS
5	H	82	GLN
1	A	469	GLN
1	A	463	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	HEA	A	602	1	67,67,67	1.16	5 (7%)	81,103,103	1.25	11 (13%)
11	CUA	B	301	2	0,1,1	-	-	-		
12	PC1	C	301	-	53,53,53	1.53	2 (3%)	59,61,61	1.02	2 (3%)
7	HEA	A	601	1	67,67,67	1.15	5 (7%)	81,103,103	1.07	4 (4%)
12	PC1	C	302	-	53,53,53	1.42	3 (5%)	59,61,61	0.90	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	HEA	A	602	1	-	13/36/76/76	-
12	PC1	C	302	-	-	8/57/57/57	-
12	PC1	C	301	-	-	17/57/57/57	-
7	HEA	A	601	1	-	9/36/76/76	-

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	C	301	PC1	O21-C21	8.09	1.57	1.34
12	C	302	PC1	O31-C31	6.75	1.53	1.33
12	C	301	PC1	O31-C31	6.40	1.52	1.33
12	C	302	PC1	O21-C21	6.35	1.52	1.34
7	A	602	HEA	CMA-C3A	4.69	1.55	1.45

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	C	301	PC1	C3-C2-C1	-4.78	100.64	111.78
7	A	601	HEA	C17-C18-C19	-3.10	120.53	127.62
12	C	302	PC1	C11-C12-N	-2.85	106.68	115.82
12	C	301	PC1	C11-C12-N	-2.65	107.33	115.82
12	C	302	PC1	C2-O21-C21	-2.60	111.58	117.80

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	602	HEA	C2A-C3A-CMA-OMA

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	A	602	HEA	C21-C22-C23-C24
12	C	301	PC1	O22-C21-O21-C2
12	C	301	PC1	C22-C21-O21-C2
12	C	302	PC1	O13-C11-C12-N


There are no ring outliers.

4 monomers are involved in 54 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	602	HEA	10	0
12	C	301	PC1	10	0
7	A	601	HEA	18	0
12	C	302	PC1	16	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.