



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 02:03 PM UTC

PDB ID : 1QM3 / pdb_00001qm3
Title : Human prion protein fragment 121-230
Authors : Zahn, R.; Liu, A.; Luhrs, T.; Wuthrich, K.
Deposited on : 1999-09-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	112	

2 Ensemble composition and analysis

This entry contains 20 models. Model 12 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:125-A:228 (104)	1.07	12

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 7 single-model clusters were found.

Cluster number	Models
1	2, 6, 19
2	1, 9, 10
3	8, 12, 20
4	7, 14
5	13, 16
Single-model clusters	3; 4; 5; 11; 15; 17; 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1688 atoms, of which 811 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PRION PROTEIN.

Mol	Chain	Residues	Atoms						Trace
1	A	104	Total	C	H	N	O	S	0
			1688	544	811	153	171	9	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	119	GLY	-	cloning artifact	UNP P04156
A	120	SER	-	cloning artifact	UNP P04156

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PRION PROTEIN




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: PRION PROTEIN



4.2.2 Score per residue for model 2

- Molecule 1: PRION PROTEIN



4.2.3 Score per residue for model 3

- Molecule 1: PRION PROTEIN

Chain A:  78% 14% • 7%



4.2.4 Score per residue for model 4

- Molecule 1: PRION PROTEIN


Chain A:  72% 19% • 7%



4.2.5 Score per residue for model 5

- Molecule 1: PRION PROTEIN

Chain A:  76% 17% • 7%



4.2.6 Score per residue for model 6

- Molecule 1: PRION PROTEIN


Chain A:  76% 16% • 7%



4.2.7 Score per residue for model 7

- Molecule 1: PRION PROTEIN

Chain A:  77% 15% • 7%



4.2.8 Score per residue for model 8

- Molecule 1: PRION PROTEIN

Chain A:  71% 19% 7%



4.2.9 Score per residue for model 9

- Molecule 1: PRION PROTEIN

Chain A:  79% 13% 7%



4.2.10 Score per residue for model 10

- Molecule 1: PRION PROTEIN

Chain A:  75% 18% 7%



4.2.11 Score per residue for model 11

- Molecule 1: PRION PROTEIN

Chain A:  70% 22% 7%



4.2.12 Score per residue for model 12 (medoid)

- Molecule 1: PRION PROTEIN


Chain A:  79% 12% 7%



4.2.13 Score per residue for model 13

- Molecule 1: PRION PROTEIN

Chain A:  76% 14% 7%



4.2.14 Score per residue for model 14

- Molecule 1: PRION PROTEIN

Chain A:  70% 21% 7%



4.2.15 Score per residue for model 15

- Molecule 1: PRION PROTEIN

Chain A:  78% 14% 7%



4.2.16 Score per residue for model 16

- Molecule 1: PRION PROTEIN

Chain A:  75% 14% 7%



4.2.17 Score per residue for model 17


- Molecule 1: PRION PROTEIN

Chain A:  73% 17% 7%




4.2.18 Score per residue for model 18

- Molecule 1: PRION PROTEIN




4.2.19 Score per residue for model 19

- Molecule 1: PRION PROTEIN



4.2.20 Score per residue for model 20

- Molecule 1: PRION PROTEIN



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
OPALp	refinement	
DYANA	structure solution	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.71±0.02	0±0/897 (0.0± 0.0%)	1.47±0.04	4±2/1210 (0.3± 0.2%)
All	All	0.71	0/17940 (0.0%)	1.47	72/24200 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.6±1.2
All	All	0	33

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	125	LEU	CA-C-N	12.72	131.41	121.61	17	4
1	A	125	LEU	C-N-CA	12.72	131.41	121.61	17	4
1	A	159	ASN	CA-CB-CG	6.79	119.39	112.60	14	2
1	A	187	HIS	CA-C-N	6.44	129.21	120.38	13	5
1	A	187	HIS	C-N-CA	6.44	129.21	120.38	13	5
1	A	166	MET	CA-C-N	6.36	129.04	120.65	4	2
1	A	166	MET	C-N-CA	6.36	129.04	120.65	4	2
1	A	140	HIS	CB-CG-CD2	-6.29	123.03	131.20	18	4
1	A	193	THR	CB-CA-C	6.17	121.12	110.94	10	3
1	A	157	TYR	CA-C-N	6.12	126.08	119.78	1	2
1	A	157	TYR	C-N-CA	6.12	126.08	119.78	1	2
1	A	192	THR	N-CA-C	-6.06	104.75	111.36	18	1
1	A	147	ASP	CA-CB-CG	5.99	118.59	112.60	18	2
1	A	173	ASN	CA-CB-CG	-5.97	106.62	112.60	5	1
1	A	210	VAL	CA-C-N	5.97	128.53	120.65	4	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	210	VAL	C-N-CA	5.97	128.53	120.65	4	2
1	A	179	CYS	CA-CB-SG	5.75	127.62	114.40	20	2
1	A	178	ASP	CA-CB-CG	5.71	118.31	112.60	19	2
1	A	146	GLU	CA-C-N	5.61	128.07	120.44	18	1
1	A	146	GLU	C-N-CA	5.61	128.07	120.44	18	1
1	A	198	PHE	CA-CB-CG	5.51	119.31	113.80	19	1
1	A	164	ARG	CB-CA-C	5.43	117.39	109.08	17	1
1	A	150	TYR	CA-C-N	5.42	127.86	120.54	13	2
1	A	150	TYR	C-N-CA	5.42	127.86	120.54	13	2
1	A	177	HIS	CA-CB-CG	5.42	119.22	113.80	19	1
1	A	209	VAL	CB-CA-C	5.39	118.59	111.70	8	1
1	A	126	GLY	N-CA-C	-5.38	104.84	111.45	6	1
1	A	147	ASP	N-CA-C	-5.33	105.11	111.03	1	1
1	A	189	VAL	CA-C-N	5.30	127.69	120.54	13	2
1	A	189	VAL	C-N-CA	5.30	127.69	120.54	13	2
1	A	191	THR	CA-CB-CG2	5.25	119.42	110.50	4	1
1	A	155	HIS	CB-CG-CD2	-5.16	124.49	131.20	18	1
1	A	210	VAL	CG1-CB-CG2	-5.09	99.61	110.80	4	1
1	A	136	ARG	CA-C-N	5.06	124.82	119.76	2	2
1	A	136	ARG	C-N-CA	5.06	124.82	119.76	2	2
1	A	181	ASN	CA-CB-CG	-5.02	107.58	112.60	8	1
1	A	223	GLN	OE1-CD-NE2	-5.01	117.59	122.60	1	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	151	ARG	Sidechain	4
1	A	156	ARG	Sidechain	4
1	A	157	TYR	Sidechain	4
1	A	208	ARG	Sidechain	3
1	A	163	TYR	Sidechain	3
1	A	148	ARG	Sidechain	3
1	A	228	ARG	Sidechain	2
1	A	226	TYR	Sidechain	2
1	A	162	TYR	Sidechain	1
1	A	225	TYR	Sidechain	1
1	A	169	TYR	Sidechain	1
1	A	153	ASN	Peptide	1
1	A	218	TYR	Sidechain	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	128	TYR	Sidechain	1
1	A	125	LEU	Peptide	1
1	A	136	ARG	Sidechain	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	877	811	813	1±1
All	All	17540	16220	16260	26

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:176:VAL:HG22	1:A:214:CYS:HB3	0.55	1.78	6	3
1:A:134:MET:HE1	1:A:217:GLN:CD	0.53	2.28	18	1
1:A:176:VAL:HG21	1:A:215:ILE:HG13	0.51	1.82	17	1
1:A:129:MET:HE2	1:A:129:MET:H	0.50	1.66	5	1
1:A:209:VAL:HG12	1:A:213:MET:HE2	0.48	1.84	18	2
1:A:191:THR:HG21	1:A:198:PHE:CE2	0.47	2.43	17	2
1:A:176:VAL:HG22	1:A:214:CYS:CB	0.47	2.39	17	1
1:A:197:ASN:HD22	1:A:197:ASN:C	0.46	2.19	8	2
1:A:206:MET:O	1:A:210:VAL:HG23	0.46	2.10	14	2
1:A:180:VAL:HA	1:A:210:VAL:HG12	0.45	1.89	14	2
1:A:176:VAL:HA	1:A:179:CYS:SG	0.44	2.53	20	2
1:A:187:HIS:CD2	1:A:206:MET:HE1	0.44	2.48	15	1
1:A:159:ASN:HA	1:A:213:MET:HE3	0.43	1.91	8	1
1:A:157:TYR:CE2	1:A:206:MET:HG3	0.43	2.48	4	1
1:A:130:LEU:HD12	1:A:161:VAL:O	0.42	2.14	11	1
1:A:150:TYR:CD1	1:A:157:TYR:CD2	0.41	3.08	1	1
1:A:191:THR:HG23	1:A:196:GLU:O	0.40	2.16	17	1
1:A:137:PRO:HD2	1:A:209:VAL:HG13	0.40	1.93	18	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/112 (91%)	92±2 (90±2%)	9±2 (9±2%)	1±1 (1±1%)	14	63
All	All	2040/2240 (91%)	1835 (90%)	183 (9%)	22 (1%)	14	63

All 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	166	MET	5
1	A	196	GLU	3
1	A	168	GLU	3
1	A	167	ASP	3
1	A	172	GLN	3
1	A	170	SER	1
1	A	189	VAL	1
1	A	132	SER	1
1	A	157	TYR	1
1	A	171	ASN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/101 (96%)	81±4 (84±4%)	16±4 (16±4%)	4	40
All	All	1940/2020 (96%)	1626 (84%)	314 (16%)	4	40

All 63 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	190	THR	20
1	A	192	THR	20
1	A	193	THR	14
1	A	187	HIS	12
1	A	191	THR	12
1	A	186	GLN	11
1	A	197	ASN	10
1	A	227	GLN	10
1	A	159	ASN	10
1	A	163	TYR	9
1	A	151	ARG	8
1	A	148	ARG	7
1	A	167	ASP	7
1	A	207	GLU	7
1	A	136	ARG	7
1	A	166	MET	6
1	A	172	GLN	6
1	A	125	LEU	6
1	A	173	ASN	6
1	A	178	ASP	6
1	A	128	TYR	6
1	A	196	GLU	6
1	A	144	ASP	5
1	A	226	TYR	5
1	A	134	MET	5
1	A	217	GLN	5
1	A	213	MET	5
1	A	221	GLU	5
1	A	164	ARG	4
1	A	212	GLN	4
1	A	200	GLU	4
1	A	168	GLU	4
1	A	154	MET	3
1	A	169	TYR	3
1	A	135	SER	3
1	A	130	LEU	3
1	A	228	ARG	3
1	A	174	ASN	3
1	A	185	LYS	3
1	A	194	LYS	3
1	A	155	HIS	3
1	A	201	THR	3
1	A	202	ASP	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	147	ASP	2
1	A	222	SER	2
1	A	129	MET	2
1	A	138	ILE	2
1	A	219	GLU	2
1	A	143	SER	2
1	A	204	LYS	2
1	A	177	HIS	2
1	A	179	CYS	2
1	A	206	MET	1
1	A	181	ASN	1
1	A	188	THR	1
1	A	175	PHE	1
1	A	216	THR	1
1	A	182	ILE	1
1	A	171	ASN	1
1	A	211	GLU	1
1	A	170	SER	1
1	A	156	ARG	1
1	A	176	VAL	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided