



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:32 PM UTC

PDB ID : 1QM6 / pdb_00001qm6
Title : Closed form of Clostridium perfringens alpha-toxin strain NCTC8237
Authors : Naylor, C.E.; Miller, J.; Titball, R.W.; Basak, A.K.
Deposited on : 1999-09-21
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	5% 84% 15% .
1	B	370	6% 85% 14% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOLIPASE C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	0	0
			3006	1901	497	599	9			
1	B	370	Total	C	N	O	S	0	0	0
			3006	1901	497	599	9			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total	O	0	0
			43	43		
3	B	46	Total	O	0	0
			46	46		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHOSPHOLIPASE C



• Molecule 1: PHOSPHOLIPASE C



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	149.54Å 149.54Å 193.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.50 30.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	90.2 (30.00-2.50) 90.2 (30.00-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.39Å)	Xtriage
Refinement program	CNS 0.5	Depositor
R, R_{free}	0.220 , 0.247 0.220 , 0.243	Depositor DCC
R_{free} test set	1291 reflections (4.43%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 28.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6105	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.78 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.8532e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3084	0.89	10/4170 (0.2%)
1	B	0.50	0/3084	0.88	6/4170 (0.1%)
All	All	0.49	0/6168	0.89	16/8340 (0.2%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	355	LYS	N-CA-C	7.26	120.25	108.41
1	B	114	TYR	N-CA-C	6.77	118.66	111.28
1	A	208	ALA	N-CA-C	6.68	121.51	113.23
1	B	208	ALA	N-CA-C	6.64	121.46	113.23
1	B	355	LYS	N-CA-C	6.44	118.90	108.41
1	A	150	LYS	N-CA-C	5.98	118.57	111.33
1	A	114	TYR	N-CA-C	5.91	117.72	111.28
1	A	210	MET	N-CA-C	5.74	118.31	111.71
1	B	86	LEU	N-CA-C	5.72	119.78	112.34
1	B	150	LYS	N-CA-C	5.67	118.19	111.33
1	A	86	LEU	N-CA-C	5.33	119.27	112.34
1	B	262	ILE	N-CA-C	5.28	115.91	108.36
1	A	262	ILE	N-CA-C	5.18	115.77	108.36
1	A	174	ALA	N-CA-C	5.17	119.91	111.37
1	A	54	TYR	N-CA-C	5.15	120.61	113.45
1	A	159	LYS	N-CA-C	5.03	117.41	111.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3006	0	2822	31	0
1	B	3006	0	2822	30	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	43	0	0	1	0
3	B	46	0	0	2	0
All	All	6105	0	5644	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:ASP:HB2	1:A:300:MET:HE1	1.52	0.89
1:B:63:ASP:HB2	1:B:300:MET:HE1	1.52	0.89
1:B:45:MET:HE1	1:B:48:LEU:HD23	1.72	0.69
1:B:143:VAL:HG11	1:B:150:LYS:HG3	1.75	0.68
1:A:45:MET:HE1	1:A:48:LEU:HD23	1.76	0.65
1:B:171:THR:HA	1:B:176:TYR:CD2	2.32	0.65
1:A:171:THR:HA	1:A:176:TYR:CD2	2.32	0.65
1:A:86:LEU:HD22	1:A:310:LYS:HB3	1.79	0.64
1:A:143:VAL:HG11	1:A:150:LYS:HG3	1.80	0.64
1:A:330:LYS:HE3	1:A:363:GLY:HA2	1.83	0.59
1:B:143:VAL:HG11	1:B:150:LYS:CG	2.34	0.58
1:B:330:LYS:HE3	1:B:363:GLY:HA2	1.85	0.57
1:B:345:ILE:HD11	1:B:369:ILE:HD11	1.88	0.55
1:A:143:VAL:HG11	1:A:150:LYS:CG	2.37	0.55
1:A:345:ILE:HD11	1:A:369:ILE:HD11	1.89	0.55
1:B:290:TRP:CE2	1:B:309:PHE:HB3	2.42	0.54
1:B:250:SER:HB2	3:B:2044:HOH:O	2.09	0.53
1:A:290:TRP:CE2	1:A:309:PHE:HB3	2.44	0.52
1:B:45:MET:HE3	1:B:48:LEU:HB3	1.95	0.49
1:A:3:GLY:HA3	3:A:2005:HOH:O	2.12	0.49
1:A:95:GLU:OE2	1:A:193:ALA:HB3	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:MET:HE3	1:A:48:LEU:HB3	1.96	0.47
1:A:131:ILE:HD11	1:A:193:ALA:HB1	1.96	0.47
1:A:242:ASP:HA	1:A:247:ASN:HB2	1.95	0.47
1:B:45:MET:CE	1:B:48:LEU:HD23	2.42	0.47
1:B:137:PRO:HB3	1:B:204:TYR:OH	2.15	0.47
1:A:90:ILE:HB	1:A:308:THR:HB	1.97	0.47
1:A:137:PRO:HB3	1:A:204:TYR:OH	2.14	0.47
1:A:45:MET:CE	1:A:48:LEU:HD23	2.44	0.46
1:B:90:ILE:HB	1:B:308:THR:HB	1.98	0.46
1:B:95:GLU:OE2	1:B:193:ALA:HB3	2.15	0.46
1:B:144:ASP:OD2	1:B:148:HIS:HD2	1.99	0.46
1:A:56:ASP:OD2	1:A:148:HIS:HE1	1.98	0.45
1:B:66:GLN:HG3	1:B:68:HIS:CE1	2.52	0.45
1:B:277:TYR:O	1:B:326:ILE:HA	2.17	0.45
1:B:56:ASP:OD2	1:B:148:HIS:HE1	1.99	0.45
1:B:86:LEU:HD22	1:B:310:LYS:HB3	1.98	0.45
1:A:182:ASN:HB3	1:A:188:TRP:HB2	2.00	0.44
1:B:45:MET:O	1:B:49:GLN:HG3	2.17	0.44
1:A:277:TYR:O	1:A:326:ILE:HA	2.18	0.44
1:A:158:ARG:HH11	1:A:158:ARG:HG3	1.83	0.43
1:B:131:ILE:HD11	1:B:193:ALA:HB1	1.99	0.43
1:B:158:ARG:HH11	1:B:158:ARG:HG3	1.84	0.43
1:B:3:GLY:HA3	3:B:2005:HOH:O	2.18	0.43
1:A:66:GLN:HG3	1:A:68:HIS:CE1	2.54	0.43
1:A:144:ASP:OD2	1:A:148:HIS:HD2	2.02	0.43
1:A:283:LYS:HE3	1:A:319:ASP:O	2.19	0.43
1:B:55:PRO:HB2	1:B:126:HIS:CD2	2.55	0.42
1:B:283:LYS:HE3	1:B:319:ASP:O	2.19	0.42
1:A:45:MET:O	1:A:49:GLN:HG3	2.19	0.42
1:B:26:LEU:HG	1:B:30:GLU:HG3	2.02	0.42
1:A:55:PRO:HB2	1:A:126:HIS:CD2	2.55	0.41
1:B:182:ASN:HB3	1:B:188:TRP:HB2	2.02	0.41
1:B:62:TYR:HH	1:B:306:THR:H	1.68	0.41
1:A:26:LEU:HG	1:A:30:GLU:HG3	2.03	0.41
1:A:13:MET:C	1:A:13:MET:SD	3.04	0.41
1:A:257:GLU:HG2	1:A:310:LYS:HA	2.03	0.41
1:B:22:LEU:HA	1:B:22:LEU:HD12	1.83	0.41
1:A:262:ILE:HG23	1:A:343:ILE:HG12	2.03	0.40
1:A:86:LEU:CD2	1:A:310:LYS:HB3	2.47	0.40
1:B:344:LYS:HA	1:B:353:VAL:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/370 (100%)	353 (96%)	13 (4%)	2 (0%)	24	43
1	B	368/370 (100%)	353 (96%)	14 (4%)	1 (0%)	36	55
All	All	736/740 (100%)	706 (96%)	27 (4%)	3 (0%)	30	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	VAL
1	A	83	SER
1	B	83	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/319 (100%)	314 (98%)	5 (2%)	55	79
1	B	319/319 (100%)	314 (98%)	5 (2%)	55	79
All	All	638/638 (100%)	628 (98%)	10 (2%)	55	79

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	VAL
1	A	22	LEU
1	A	29	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	39	GLU
1	A	66	GLN
1	B	15	VAL
1	B	22	LEU
1	B	29	ASN
1	B	39	GLU
1	B	66	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	110	GLN
1	A	148	HIS
1	A	165	ASN
1	A	294	ASN
1	A	297	ASN
1	A	348	ASN
1	B	77	ASN
1	B	110	GLN
1	B	148	HIS
1	B	165	ASN
1	B	182	ASN
1	B	294	ASN
1	B	297	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/370 (100%)	0.07	17 (4%) 37 33	7, 15, 38, 65	0
1	B	370/370 (100%)	0.14	21 (5%) 29 26	7, 15, 38, 65	0
All	All	740/740 (100%)	0.11	38 (5%) 33 29	7, 15, 38, 65	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	251	VAL	7.0
1	A	251	VAL	5.4
1	B	61	ALA	5.4
1	B	249	PRO	5.1
1	A	61	ALA	4.9
1	B	60	ASN	4.6
1	B	252	GLY	4.6
1	A	248	ASP	4.6
1	A	64	LEU	4.3
1	B	64	LEU	4.1
1	B	66	GLN	3.9
1	B	62	TYR	3.5
1	B	322	GLN	3.3
1	B	247	ASN	3.3
1	B	248	ASP	3.1
1	B	63	ASP	3.0
1	A	246	GLY	2.8
1	B	246	GLY	2.8
1	A	62	TYR	2.7
1	A	60	ASN	2.7
1	A	249	PRO	2.7
1	A	252	GLY	2.6
1	A	157	GLU	2.6
1	A	66	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	59	LYS	2.5
1	A	250	SER	2.5
1	A	253	LYS	2.4
1	B	368	ASN	2.4
1	A	247	ASN	2.3
1	B	331	TYR	2.3
1	B	110	GLN	2.3
1	B	335	SER	2.2
1	A	191	GLU	2.2
1	B	160	GLU	2.1
1	B	250	SER	2.1
1	B	283	LYS	2.1
1	A	284	ASP	2.0
1	A	63	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	400	1/1	0.97	0.15	21,21,21,21	0
2	ZN	B	401	1/1	0.98	0.15	22,22,22,22	0
2	ZN	A	400	1/1	0.99	0.08	13,13,13,13	0
2	ZN	A	401	1/1	0.99	0.10	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.