



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 07:30 PM UTC

PDB ID : 1QMD / pdb_00001qmd
Title : calcium bound closed form alpha-toxin from Clostridium perfringens
Authors : Naylor, C.E.; Miller, J.; Titball, R.W.; Basak, A.K.
Deposited on : 1999-09-27
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	4% 85% 14%
1	B	370	3% 84% 16%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOLIPASE C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	2	0
			3014	1906	498	600	10			
1	B	370	Total	C	N	O	S	0	2	0
			3014	1906	498	600	10			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		
3	B	3	Total	Ca	0	0
			3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	161	Total	O	0	0
			161	161		
4	B	152	Total	O	0	0
			152	152		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHOSPHOLIPASE C



• Molecule 1: PHOSPHOLIPASE C



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	149.89Å 149.89Å 192.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.20 50.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-2.20) 98.0 (50.00-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.44 (at 2.20Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.198 , 0.235 0.197 , 0.233	Depositor DCC
R_{free} test set	2108 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.660	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for -2/3*h-1/3*k+2/3*l,-1/3*h-2/3*k-2/3*l,2/3*h-2/3*k+1/3*l 0.001 for -h,1/3*h-1/3*k+2/3*l,2/3*h+4/3*k+1/3*l 0.000 for -1/3*h+1/3*k-2/3*l,-k,-4/3*h-2/3*k+1/3*l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6351	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.1898e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/3100	0.90	8/4191 (0.2%)
1	B	0.43	0/3100	0.91	9/4191 (0.2%)
All	All	0.43	0/6200	0.91	17/8382 (0.2%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	159	LYS	N-CA-C	6.01	118.60	111.33
1	A	159	LYS	N-CA-C	5.76	118.30	111.33
1	A	208	ALA	N-CA-C	5.67	120.19	113.28
1	B	150	LYS	N-CA-C	5.60	118.16	111.71
1	B	208	ALA	N-CA-C	5.59	120.38	113.50
1	B	355	LYS	N-CA-C	5.53	117.54	108.52
1	A	174	ALA	N-CA-C	5.53	120.44	111.37
1	B	174	ALA	N-CA-C	5.47	120.40	111.37
1	B	114	TYR	N-CA-C	5.44	118.13	111.82
1	B	86	LEU	N-CA-C	5.43	118.70	111.75
1	A	86	LEU	N-CA-C	5.39	118.66	111.75
1	A	355	LYS	N-CA-C	5.33	117.20	108.52
1	A	150	LYS	N-CA-C	5.32	117.08	111.28
1	B	251	VAL	N-CA-C	5.19	120.13	109.34
1	A	54	TYR	N-CA-C	5.16	120.63	113.45
1	B	331	TYR	N-CA-C	-5.04	105.49	111.69
1	A	369	ILE	N-CA-C	-5.02	105.53	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3014	0	2829	33	0
1	B	3014	0	2829	27	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	161	0	0	5	0
4	B	152	0	0	3	0
All	All	6351	0	5658	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:GLN:HE22	1:A:165[B]:ASN:HD21	1.30	0.79
1:A:253:LYS:HE3	1:A:348:ASN:HB3	1.68	0.75
1:A:11:HIS:HD2	4:A:2006:HOH:O	1.72	0.71
1:B:278:PHE:HE1	1:B:369:ILE:HD11	1.58	0.67
1:A:106:ARG:O	1:A:110:GLN:HG2	1.97	0.65
1:B:281:LYS:HE2	4:B:2141:HOH:O	1.96	0.65
1:B:248:ASP:HB3	1:B:251:VAL:HG13	1.79	0.64
1:A:17:GLN:HE22	1:A:165[B]:ASN:ND2	1.99	0.59
1:A:249:PRO:O	1:A:253:LYS:HE2	2.03	0.59
1:A:17:GLN:NE2	1:A:165[B]:ASN:ND2	2.54	0.54
1:A:250:SER:HB2	4:A:2152:HOH:O	2.06	0.54
1:A:23:GLU:OE2	1:A:42:LYS:HD2	2.08	0.53
1:B:242:ASP:HA	1:B:247:ASN:HD22	1.73	0.53
1:A:17:GLN:NE2	1:A:165[B]:ASN:HD21	2.03	0.53
1:A:86:LEU:HD22	1:A:310:LYS:HB3	1.91	0.53
1:B:249:PRO:HA	4:B:2097:HOH:O	2.08	0.53
1:B:45:MET:O	1:B:49:GLN:HG3	2.09	0.52
1:A:29:ASN:HD22	1:A:29:ASN:H	1.56	0.52
1:B:250:SER:HB2	4:B:2145:HOH:O	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:LYS:O	1:A:194:ARG:HB2	2.10	0.51
1:A:158:ARG:O	1:A:161:GLN:HG2	2.11	0.51
1:B:190:LYS:O	1:B:194:ARG:HB2	2.11	0.51
1:A:171:THR:HA	1:A:176:TYR:CD2	2.46	0.51
1:A:29:ASN:HD22	1:A:29:ASN:N	2.09	0.50
1:A:143:VAL:HG11	1:A:150:LYS:HG3	1.93	0.49
1:B:290:TRP:CE2	1:B:309:PHE:HB3	2.48	0.49
1:B:26:LEU:HG	1:B:30:GLU:HG3	1.95	0.49
1:B:135:TYR:CE2	1:B:149:VAL:HG11	2.48	0.49
1:A:249:PRO:HB2	4:A:2098:HOH:O	2.13	0.48
1:A:252:GLY:O	1:A:317:LYS:HE3	2.14	0.48
1:A:290:TRP:CE2	1:A:309:PHE:HB3	2.49	0.48
1:A:78:PHE:CE2	1:A:91:PRO:HG2	2.50	0.46
1:B:98:ILE:HD11	1:B:128:PHE:HA	1.97	0.46
1:B:131:ILE:HD11	1:B:193:ALA:HB1	1.98	0.46
1:B:344:LYS:HA	1:B:353:VAL:O	2.16	0.45
1:A:355:LYS:HE2	1:A:367:TYR:OH	2.16	0.45
1:A:22:LEU:O	1:A:26:LEU:HB2	2.16	0.44
1:A:135:TYR:CE2	1:A:149:VAL:HG11	2.52	0.44
1:A:178:ASP:HA	1:A:181:LYS:HE2	1.99	0.44
1:B:151:PHE:HA	1:B:217:TRP:CZ3	2.52	0.44
1:A:307:TYR:HA	4:A:2051:HOH:O	2.17	0.44
1:A:350:LYS:HD3	4:A:2122:HOH:O	2.18	0.44
1:B:330:LYS:CE	1:B:363:GLY:HA2	2.47	0.43
1:B:77:ASN:HD22	1:B:80:LYS:NZ	2.17	0.43
1:B:23:GLU:OE2	1:B:42:LYS:HD2	2.19	0.43
1:B:330:LYS:HE3	1:B:363:GLY:HA2	2.00	0.43
1:B:171:THR:HA	1:B:176:TYR:CD2	2.54	0.42
1:B:280:ILE:HG22	1:B:324:MET:HG2	2.01	0.42
1:B:2:ASP:HB3	1:B:156:GLU:OE1	2.20	0.42
1:A:209:SER:OG	1:A:212:HIS:HD2	2.03	0.42
1:B:86:LEU:HD12	1:B:312:LYS:HG2	2.02	0.42
1:A:125:MET:HE1	1:A:235:TYR:CG	2.55	0.42
1:A:143:VAL:HG11	1:A:150:LYS:CG	2.50	0.41
1:B:47:GLU:HG2	1:B:115:LYS:O	2.19	0.41
1:A:345:ILE:HB	1:A:353:VAL:HB	2.02	0.41
1:B:162:TYR:CD2	1:B:225:LEU:HD12	2.56	0.41
1:A:162:TYR:CD2	1:A:225:LEU:HD12	2.56	0.41
1:B:107:TYR:O	1:B:110:GLN:HG3	2.21	0.41
1:B:22:LEU:HD23	1:B:22:LEU:HA	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/370 (100%)	358 (97%)	11 (3%)	1 (0%)	36	42
1	B	370/370 (100%)	357 (96%)	12 (3%)	1 (0%)	36	42
All	All	740/740 (100%)	715 (97%)	23 (3%)	2 (0%)	36	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	VAL
1	B	251	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/319 (101%)	316 (98%)	5 (2%)	55	71
1	B	321/319 (101%)	316 (98%)	5 (2%)	55	71
All	All	642/638 (101%)	632 (98%)	10 (2%)	55	71

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	HIS
1	A	29	ASN
1	A	160	GLU
1	A	258	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	315	ASN
1	B	29	ASN
1	B	39	GLU
1	B	253	LYS
1	B	258	LEU
1	B	268	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	HIS
1	A	17	GLN
1	A	29	ASN
1	A	76	ASN
1	A	186	ASN
1	A	212	HIS
1	A	322	GLN
1	B	49	GLN
1	B	76	ASN
1	B	77	ASN
1	B	186	ASN
1	B	207	HIS
1	B	212	HIS
1	B	247	ASN
1	B	294	ASN
1	B	322	GLN
1	B	368	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	370/370 (100%)	0.05	13 (3%)	47	44	9, 25, 49, 88	2 (0%)
1	B	370/370 (100%)	0.03	12 (3%)	50	47	10, 25, 49, 93	2 (0%)
All	All	740/740 (100%)	0.04	25 (3%)	48	45	9, 25, 49, 93	4 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	250	SER	6.5
1	B	249	PRO	6.4
1	A	249	PRO	5.8
1	A	248	ASP	5.7
1	B	251	VAL	5.5
1	B	247	ASN	5.1
1	B	248	ASP	5.0
1	A	251	VAL	5.0
1	B	63	ASP	3.8
1	A	250	SER	3.8
1	B	246	GLY	3.5
1	A	247	ASN	3.5
1	A	246	GLY	3.4
1	B	35	ARG	3.0
1	A	64	LEU	2.9
1	A	252	GLY	2.6
1	B	331	TYR	2.5
1	B	283	LYS	2.5
1	B	252	GLY	2.5
1	A	35	ARG	2.4
1	A	29	ASN	2.4
1	B	334	PHE	2.3
1	A	253	LYS	2.3
1	A	283	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	110	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	A	404	1/1	0.94	0.05	30,30,30,30	0
3	CA	B	405	1/1	0.94	0.07	51,51,51,51	0
3	CA	A	405	1/1	0.96	0.06	42,42,42,42	0
3	CA	A	403	1/1	0.97	0.04	27,27,27,27	0
3	CA	B	403	1/1	0.98	0.03	31,31,31,31	0
3	CA	B	404	1/1	0.98	0.03	29,29,29,29	0
2	ZN	A	401	1/1	0.98	0.11	21,21,21,21	0
2	ZN	A	402	1/1	0.99	0.04	17,17,17,17	0
2	ZN	B	401	1/1	0.99	0.03	16,16,16,16	0
2	ZN	B	402	1/1	0.99	0.09	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.