



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:32 PM UTC

PDB ID : 4QN4 / pdb_00004qn4
Title : Crystal structure of Neuraminidase N6
Authors : Sun, X.; Li, Q.; Wu, Y.; Liu, Y.; Qi, J.; Vavricka, C.J.; Gao, G.F.
Deposited on : 2014-06-17
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	% 96% ..
1	B	391	% 95% ..
2	C	3	33% 67%
2	E	3	67% 33%
3	D	6	50% 33% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	6	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: green (50%), yellow (33%), and orange (17%).

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7354 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	389	Total	C	N	O	S	0	0	0
			3004	1875	532	571	26			
1	B	389	Total	C	N	O	S	0	1	0
			3010	1878	533	573	26			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	PHE	LYS	conflict	UNP Q2FCL6
B	2	PHE	LYS	conflict	UNP Q2FCL6

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	E	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	6	Total	C	N	O	0	0	0
			72	40	2	30			
3	F	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		
5	A	1	Total	C	O	0	0
			8	6	2		
5	B	1	Total	C	O	0	0
			8	6	2		
5	B	1	Total	C	O	0	0
			8	6	2		
5	B	1	Total	C	O	0	0
			8	6	2		
5	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	532	Total	O	0	0
			532	532		
7	B	524	Total	O	0	0
			524	524		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neuraminidase

Chain A: 




- Molecule 1: Neuraminidase

Chain B: 



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 




- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 



- Molecule 3: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] α -D-mannopyranose-(1-6)- β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain F:  50% 33% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	138.14Å 138.14Å 150.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.69 – 1.80 47.69 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (47.69-1.80) 98.0 (47.69-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.141 , 0.156 0.142 , 0.157	Depositor DCC
R_{free} test set	6603 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7354	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.91 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.8485e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, MAN, MPD, GOL, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3080	0.73	1/4182 (0.0%)
1	B	0.38	0/3086	0.73	1/4190 (0.0%)
All	All	0.39	0/6166	0.73	2/8372 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	GLN	N-CA-C	6.36	118.29	111.36
1	B	148	GLN	N-CA-C	5.83	117.72	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3004	0	2882	10	0
1	B	3010	0	2886	12	0
2	C	39	0	34	1	0
2	E	39	0	34	2	0
3	D	72	0	61	1	0
3	F	72	0	61	1	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
5	A	16	0	28	1	0
5	B	32	0	56	1	0
6	A	6	0	8	0	0
6	B	6	0	8	0	0
7	A	532	0	0	5	0
7	B	524	0	0	9	2
All	All	7354	0	6058	27	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:ASN:ND2	7:A:1009:HOH:O	2.00	0.87
7:B:1004:HOH:O	2:E:3:BMA:O6	1.96	0.83
1:A:375:ARG:NH1	7:A:941:HOH:O	2.09	0.78
7:A:1020:HOH:O	2:C:3:BMA:O6	2.12	0.68
7:B:803:HOH:O	3:F:3:BMA:O3	2.12	0.67
7:B:1019:HOH:O	2:E:3:BMA:O6	2.18	0.56
1:B:7:ASN:ND2	7:B:983:HOH:O	2.26	0.54
1:B:322:GLN:NE2	7:B:958:HOH:O	2.17	0.54
1:B:259:ASP:OD2	7:B:1097:HOH:O	2.18	0.53
1:B:353:ARG:CZ	7:B:1036:HOH:O	2.57	0.52
1:A:42:TYR:CG	1:A:150:SER:HA	2.45	0.51
1:B:175:LYS:NZ	1:B:189:GLU:OE2	2.36	0.51
1:B:42:TYR:CG	1:B:150:SER:HA	2.46	0.50
1:B:353:ARG:NH1	7:B:1036:HOH:O	2.45	0.48
1:A:210:LYS:HE3	5:A:513:MPD:H4	1.97	0.46
1:B:110:SER:HB2	1:B:129:TYR:CZ	2.51	0.45
1:B:210:LYS:HE3	5:B:515:MPD:H4	1.98	0.45
1:A:332:ILE:HB	1:A:334:TYR:CZ	2.51	0.45
1:B:332:ILE:HB	1:B:334:TYR:CZ	2.52	0.44
1:A:32:ASP:CG	1:A:32:ASP:O	2.60	0.44
3:D:1:NAG:H61	3:D:2:NAG:C7	2.49	0.43
1:B:338:LYS:NZ	7:B:1105:HOH:O	2.52	0.42
1:A:110:SER:HB2	1:A:129:TYR:CZ	2.55	0.41
1:A:45:CYS:HA	1:A:50:CYS:HA	2.03	0.41
1:B:32:ASP:O	1:B:32:ASP:CG	2.63	0.41
1:A:375:ARG:NH2	7:A:809:HOH:O	2.54	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:375:ARG:HD2	7:A:941:HOH:O	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:992:HOH:O	7:B:1060:HOH:O[3_555]	1.74	0.46
7:B:993:HOH:O	7:B:1083:HOH:O[3_555]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/391 (99%)	372 (96%)	15 (4%)	0	100	100
1	B	388/391 (99%)	373 (96%)	15 (4%)	0	100	100
All	All	775/782 (99%)	745 (96%)	30 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/333 (99%)	331 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	332/333 (100%)	332 (100%)	0	100	100
All	All	663/666 (100%)	663 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	48	GLN
1	A	251	ASN
1	A	255	ASN
1	A	303	ASN
1	A	323	ASN
1	B	303	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.60	0	17,19,21	0.89	1 (5%)
2	NAG	C	2	2	14,14,15	0.57	0	17,19,21	0.82	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	C	3	2	11,11,12	0.60	0	15,15,17	0.77	0
3	NAG	D	1	1,3	14,14,15	0.59	0	17,19,21	1.51	2 (11%)
3	NAG	D	2	3	14,14,15	0.65	0	17,19,21	0.74	0
3	BMA	D	3	3	11,11,12	0.71	0	15,15,17	1.17	1 (6%)
3	MAN	D	4	3	11,11,12	0.60	0	15,15,17	0.75	0
3	MAN	D	5	3	11,11,12	0.60	0	15,15,17	0.68	0
3	MAN	D	6	3	11,11,12	0.51	0	15,15,17	0.78	0
2	NAG	E	1	2,1	14,14,15	0.62	0	17,19,21	0.74	0
2	NAG	E	2	2	14,14,15	0.55	0	17,19,21	0.82	0
2	BMA	E	3	2	11,11,12	0.62	0	15,15,17	0.91	0
3	NAG	F	1	1,3	14,14,15	0.66	0	17,19,21	1.22	3 (17%)
3	NAG	F	2	3	14,14,15	0.61	0	17,19,21	0.78	0
3	BMA	F	3	3	11,11,12	0.51	0	15,15,17	1.29	2 (13%)
3	MAN	F	4	3	11,11,12	0.56	0	15,15,17	0.73	0
3	MAN	F	5	3	11,11,12	0.56	0	15,15,17	0.56	0
3	MAN	F	6	3	11,11,12	0.58	0	15,15,17	0.87	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	1/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	0/2/19/22	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	BMA	E	3	2	-	0/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
3	MAN	F	5	3	-	0/2/19/22	0/1/1/1
3	MAN	F	6	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	O5-C1-C2	-4.22	104.76	111.29
3	F	3	BMA	C1-O5-C5	3.44	116.80	112.19
3	D	3	BMA	C1-O5-C5	3.27	116.57	112.19
3	D	1	NAG	C6-C5-C4	-2.82	106.09	113.02
3	F	1	NAG	O5-C1-C2	-2.49	107.43	111.29
3	F	3	BMA	O2-C2-C3	-2.28	105.43	110.15
3	F	1	NAG	C6-C5-C4	-2.24	107.52	113.02
3	F	6	MAN	C1-O5-C5	2.22	115.17	112.19
2	C	1	NAG	O5-C1-C2	-2.12	108.01	111.29
3	F	1	NAG	C2-N2-C7	-2.01	120.21	122.90

There are no chirality outliers.

All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	D	1	NAG	C4-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6
2	C	3	BMA	O5-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	2	NAG	1	0
3	F	3	BMA	1	0
2	E	3	BMA	2	0
3	D	1	NAG	1	0
2	C	3	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	512	-	5,5,5	0.35	0	5,5,5	0.38	0
5	MPD	A	511	-	7,7,7	0.23	0	9,10,10	0.25	0
5	MPD	A	513	-	7,7,7	0.33	0	9,10,10	0.40	0
6	GOL	B	514	-	5,5,5	0.39	0	5,5,5	0.36	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MPD	B	513	-	7,7,7	0.25	0	9,10,10	0.28	0
5	MPD	B	515	-	7,7,7	0.32	0	9,10,10	0.37	0
5	MPD	B	511	-	7,7,7	0.29	0	9,10,10	0.61	0
5	MPD	B	512	-	7,7,7	0.25	0	9,10,10	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	512	-	-	0/4/4/4	-
5	MPD	A	511	-	-	0/5/5/5	-
5	MPD	A	513	-	-	3/5/5/5	-
6	GOL	B	514	-	-	0/4/4/4	-
5	MPD	B	513	-	-	0/5/5/5	-
5	MPD	B	515	-	-	3/5/5/5	-
5	MPD	B	511	-	-	0/5/5/5	-
5	MPD	B	512	-	-	2/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	513	MPD	CM-C2-C3-C4
5	B	512	MPD	C1-C2-C3-C4
5	B	515	MPD	CM-C2-C3-C4
5	A	513	MPD	C2-C3-C4-O4
5	B	515	MPD	C2-C3-C4-O4
5	A	513	MPD	C1-C2-C3-C4
5	B	512	MPD	CM-C2-C3-C4
5	B	515	MPD	C1-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	513	MPD	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	515	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	389/391 (99%)	-0.78	2 (0%) 87 88	4, 8, 18, 34	0
1	B	389/391 (99%)	-0.73	2 (0%) 87 88	5, 10, 19, 35	1 (0%)
All	All	778/782 (99%)	-0.75	4 (0%) 87 88	4, 9, 19, 35	1 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	375	ARG	2.8
1	B	375	ARG	2.7
1	B	48	GLN	2.4
1	A	48	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	1	14/15	0.84	0.14	22,30,36,36	0
3	NAG	F	1	14/15	0.87	0.13	26,30,39,40	0
3	BMA	F	3	11/12	0.92	0.09	15,19,26,32	0
3	MAN	F	6	11/12	0.92	0.09	16,21,27,37	0
2	BMA	E	3	11/12	0.93	0.08	14,23,27,37	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	F	2	14/15	0.93	0.08	15,20,25,31	0
2	BMA	C	3	11/12	0.93	0.09	15,23,29,33	0
3	NAG	D	2	14/15	0.93	0.08	11,16,22,24	0
3	MAN	F	5	11/12	0.94	0.07	17,21,24,24	0
3	BMA	D	3	11/12	0.94	0.08	14,17,22,29	0
3	MAN	D	6	11/12	0.95	0.08	13,17,29,35	0
3	MAN	D	5	11/12	0.95	0.07	14,19,23,23	0
3	MAN	D	4	11/12	0.96	0.06	13,15,17,17	0
3	MAN	F	4	11/12	0.96	0.06	15,17,18,19	0
2	NAG	C	1	14/15	0.97	0.05	5,8,11,13	0
2	NAG	E	2	14/15	0.97	0.05	8,10,17,19	0
2	NAG	C	2	14/15	0.97	0.06	7,12,18,21	0
2	NAG	E	1	14/15	0.98	0.05	6,7,11,13	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	A	512	6/6	0.89	0.13	17,21,25,27	0
5	MPD	B	512	8/8	0.90	0.12	12,20,24,29	0
5	MPD	B	515	8/8	0.91	0.12	15,20,23,25	0
5	MPD	B	511	8/8	0.91	0.10	11,16,21,31	0
6	GOL	B	514	6/6	0.91	0.11	20,22,24,24	0
5	MPD	A	513	8/8	0.92	0.11	10,16,22,26	0
5	MPD	A	511	8/8	0.94	0.10	16,19,21,24	0
5	MPD	B	513	8/8	0.95	0.09	17,19,22,28	0
4	CA	B	501	1/1	0.99	0.16	25,25,25,25	0
4	CA	A	501	1/1	1.00	0.14	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.