



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:49 AM UTC

PDB ID : 4QNP / pdb_00004qnp
Title : Crystal structure of the 2009 pandemic H1N1 influenza virus neuraminidase with a neutralizing antibody
Authors : Wan, H.Q.; Yang, H.; Shore, D.A.; Garten, R.J.; Couzens, L.; Gao, J.; Jiang, L.L.; Carney, P.J.; Villanueva, J.; Stevens, J.; Eichelberger, M.C.
Deposited on : 2014-06-18
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	% 93% 6% .
1	B	386	% 91% 8% .
2	F	212	86% 13% .
2	L	212	87% 11%
3	E	224	88% 12% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	224	% 84% 13% .
4	C	2	100%
4	D	2	50% 50%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			2983	1873	515	574	21			
1	B	386	Total	C	N	O	S	0	0	0
			2983	1873	515	574	21			

- Molecule 2 is a protein called neutralizing antibody, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	212	Total	C	N	O	S	0	0	0
			1635	1019	276	333	7			
2	L	211	Total	C	N	O	S	0	0	0
			1626	1014	275	330	7			

- Molecule 3 is a protein called neutralizing antibody, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	224	Total	C	N	O	S	0	0	0
			1704	1078	283	334	9			
3	H	224	Total	C	N	O	S	0	0	0
			1704	1078	283	334	9			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Ca	0	0
			2	2		
6	B	2	Total	Ca	0	0
			2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	40	Total 40	O 40	0	0
7	B	50	Total 50	O 50	0	0
7	F	17	Total 17	O 17	0	0
7	E	30	Total 30	O 30	0	0
7	L	14	Total 14	O 14	0	0
7	H	30	Total 30	O 30	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neuraminidase




- Molecule 1: Neuraminidase




- Molecule 2: neutralizing antibody, light chain





- Molecule 2: neutralizing antibody, light chain



- Molecule 3: neutralizing antibody, heavy chain



- Molecule 3: neutralizing antibody, heavy chain



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	144.29Å 202.95Å 143.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.85 – 2.80 47.85 – 2.80	Depositor EDS
% Data completeness (in resolution range)	92.9 (47.85-2.80) 92.9 (47.85-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.57 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.181 , 0.224 0.188 , 0.227	Depositor DCC
R_{free} test set	2458 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	49.3	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12932	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.96	0/3065	1.03	3/4166 (0.1%)
1	B	0.94	0/3065	1.03	5/4166 (0.1%)
2	F	0.84	0/1677	0.97	1/2282 (0.0%)
2	L	0.81	0/1668	0.94	2/2270 (0.1%)
3	E	0.81	1/1747 (0.1%)	0.92	1/2384 (0.0%)
3	H	0.83	0/1747	0.97	4/2384 (0.2%)
All	All	0.89	1/12969 (0.0%)	0.99	16/17652 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	H	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	50	ARG	NE-CZ	5.27	1.38	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	GLN	N-CA-C	7.96	119.95	111.28
3	H	116	ALA	N-CA-C	-7.80	100.23	110.53
1	B	227	GLN	N-CA-C	6.97	118.96	111.36
2	L	193	CYS	CB-CA-C	-6.78	97.18	109.38
3	H	116	ALA	CA-C-N	-6.56	112.28	121.67
3	H	116	ALA	C-N-CA	-6.56	112.28	121.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	193	CYS	CB-CA-C	-6.36	97.94	109.38
3	H	93	THR	N-CA-CB	6.33	119.29	109.85
2	L	179	THR	CB-CA-C	5.54	119.86	109.71
1	B	125	SER	CA-C-N	-5.38	114.28	119.87
1	B	125	SER	C-N-CA	-5.38	114.28	119.87
3	E	43	LYS	N-CA-C	5.31	120.41	113.30
1	A	181	ALA	N-CA-C	5.15	116.63	108.96
1	B	434	ASN	N-CA-C	5.11	117.46	108.56
1	A	389	ILE	N-CA-C	5.05	114.53	108.06
1	B	347	LYS	N-CA-C	-5.02	102.95	110.23

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	H	8	GLY	Peptide
3	H	9	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2983	0	2815	12	0
1	B	2983	0	2815	13	0
2	F	1635	0	1559	22	0
2	L	1626	0	1553	19	0
3	E	1704	0	1668	15	0
3	H	1704	0	1668	21	0
4	C	28	0	25	0	0
4	D	28	0	25	0	1
5	A	28	0	26	1	0
5	B	28	0	26	1	0
6	A	2	0	0	0	0
6	B	2	0	0	0	0
7	A	40	0	0	0	0
7	B	50	0	0	1	0
7	E	30	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	F	17	0	0	0	0
7	H	30	0	0	1	0
7	L	14	0	0	0	0
All	All	12932	0	12180	94	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:90:THR:O	3:H:93:THR:HG23	1.72	0.90
1:B:89:SER:HA	5:B:502:NAG:H82	1.66	0.76
3:E:145:SER:O	3:E:196:SER:OG	2.04	0.76
3:H:6:GLU:OE2	3:H:116:ALA:O	2.02	0.76
3:H:1:GLN:O	7:H:326:HOH:O	2.11	0.68
1:B:170:TYR:O	7:B:605:HOH:O	2.13	0.66
2:F:143:ILE:HG23	2:F:174:MET:HE3	1.76	0.66
2:F:185:TYR:CE2	2:F:210:ARG:HD2	2.31	0.65
1:A:89:SER:HA	5:A:504:NAG:H82	1.79	0.65
2:F:143:ILE:CG2	2:F:174:MET:HE3	2.27	0.64
3:E:32:TYR:O	3:E:74:ARG:NH2	2.27	0.63
3:E:66:VAL:HG13	3:E:70:PHE:HB2	1.81	0.62
3:H:66:VAL:HG13	3:H:70:PHE:HB2	1.81	0.62
1:A:399:TRP:CZ3	1:A:432:LYS:HD3	2.35	0.61
3:H:6:GLU:CD	3:H:116:ALA:O	2.45	0.59
1:B:399:TRP:CZ3	1:B:432:LYS:HD3	2.37	0.59
1:B:399:TRP:HZ3	1:B:432:LYS:HD3	1.67	0.59
2:F:34:TRP:HB2	2:F:47:ILE:HG23	1.87	0.57
2:F:47:ILE:HD11	2:F:50:THR:O	2.04	0.56
3:H:141:ASP:C	3:H:143:THR:HG23	2.32	0.55
1:A:399:TRP:HZ3	1:A:432:LYS:HD3	1.73	0.54
1:B:462:GLU:HA	1:B:462:GLU:OE1	2.08	0.54
3:E:155:GLY:HA2	3:E:185:LEU:HG	1.89	0.53
2:F:29:VAL:HG11	2:F:89:GLN:HG3	1.89	0.53
2:L:182:LYS:NZ	2:L:186:GLU:OE2	2.40	0.53
2:L:29:VAL:HG11	2:L:89:GLN:HG3	1.89	0.53
2:L:90:TRP:CD2	2:L:95:PRO:HB3	2.43	0.52
2:F:146:LYS:HB3	2:F:194:GLU:HB3	1.91	0.52
3:H:141:ASP:OD2	3:H:141:ASP:N	2.42	0.52
1:A:220:ARG:NH2	2:F:57:VAL:O	2.43	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:182:LYS:O	2:L:186:GLU:HG2	2.11	0.51
1:A:259:GLU:O	1:A:262:LYS:HG3	2.10	0.51
3:E:69:ARG:NH2	3:E:92:ASP:OD2	2.43	0.51
3:H:69:ARG:NH2	3:H:92:ASP:OD2	2.44	0.51
2:F:188:HIS:O	2:F:210:ARG:NH1	2.44	0.51
2:L:146:LYS:HB3	2:L:194:GLU:HB3	1.92	0.50
1:A:346:VAL:HG13	1:A:368:ARG:NE	2.26	0.50
3:E:58:TYR:HE1	3:E:74:ARG:HD2	1.77	0.50
2:F:162:TRP:NE1	2:F:174:MET:HE2	2.27	0.50
2:L:134:PHE:CD2	3:H:191:SER:HB3	2.46	0.50
3:E:145:SER:O	3:E:196:SER:N	2.43	0.49
2:F:90:TRP:CD2	2:F:95:PRO:HB3	2.48	0.49
2:L:166:ASP:OD1	2:L:167:SER:N	2.46	0.49
2:F:174:MET:HG2	2:F:175:SER:N	2.27	0.48
3:E:162:THR:O	3:E:208:VAL:HA	2.14	0.48
1:A:113:ASP:O	1:A:169:PRO:HD2	2.14	0.47
3:H:162:THR:O	3:H:208:VAL:HA	2.14	0.47
3:H:164:THR:OG1	3:H:207:ASN:HB2	2.13	0.47
3:H:181:LEU:HD11	3:H:184:ASP:HA	1.96	0.47
3:H:141:ASP:O	3:H:143:THR:HG23	2.15	0.47
1:B:346:VAL:HG13	1:B:368:ARG:NE	2.30	0.47
2:F:166:ASP:OD1	2:F:168:LYS:N	2.47	0.46
1:B:173:ARG:HD2	1:B:175:GLU:OE2	2.16	0.46
3:E:181:LEU:HD11	3:E:184:ASP:HA	1.97	0.46
3:E:58:TYR:CE1	3:E:74:ARG:HD2	2.51	0.46
3:E:164:THR:OG1	3:E:207:ASN:HB2	2.14	0.46
2:F:143:ILE:HG23	2:F:174:MET:CE	2.43	0.46
2:L:174:MET:HG2	2:L:175:SER:N	2.32	0.45
2:F:134:PHE:CD2	3:E:191:SER:HB3	2.52	0.45
2:L:60:ARG:NH2	2:L:81:ASP:OD1	2.46	0.45
2:L:166:ASP:OD1	2:L:168:LYS:N	2.48	0.45
1:B:241:VAL:HG22	1:B:255:ILE:HG22	1.99	0.45
1:B:113:ASP:O	1:B:169:PRO:HD2	2.18	0.44
2:F:166:ASP:OD1	2:F:167:SER:N	2.50	0.44
2:L:185:TYR:CE2	2:L:210:ARG:HD3	2.53	0.44
1:A:155:TYR:CE1	1:B:460:GLY:HA3	2.53	0.44
2:L:47:ILE:HD11	2:L:72:LEU:HD13	2.00	0.44
2:L:130:SER:OG	2:L:179:THR:HB	2.17	0.44
1:A:241:VAL:HG22	1:A:255:ILE:HG22	1.99	0.43
3:E:145:SER:O	3:E:196:SER:CB	2.66	0.43
3:H:90:THR:O	3:H:93:THR:CG2	2.56	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:60:ARG:NH2	2:F:81:ASP:OD1	2.43	0.42
3:E:174:VAL:HG22	3:E:192:VAL:HG23	2.02	0.42
2:L:168:LYS:N	2:L:168:LYS:HD3	2.32	0.42
2:L:134:PHE:CE2	3:H:191:SER:HB3	2.54	0.42
2:F:162:TRP:CE2	2:F:174:MET:HE2	2.54	0.42
3:H:95:MET:SD	3:H:117:GLY:HA3	2.59	0.42
1:A:229:SER:HB3	1:A:347:LYS:HE2	2.00	0.42
1:B:117:ILE:O	1:B:118:ARG:HG2	2.20	0.41
2:F:134:PHE:CE2	3:E:191:SER:HB3	2.55	0.41
2:L:182:LYS:O	2:L:186:GLU:CG	2.68	0.41
2:L:168:LYS:HD3	2:L:168:LYS:HA	1.86	0.41
3:H:174:VAL:HG22	3:H:192:VAL:HG23	2.03	0.41
1:A:117:ILE:O	1:A:118:ARG:HG2	2.21	0.41
2:L:185:TYR:CZ	2:L:210:ARG:HD3	2.55	0.41
1:A:220:ARG:HD2	2:F:59:PHE:CZ	2.55	0.41
2:F:60:ARG:HH21	2:F:81:ASP:CG	2.28	0.41
3:H:220:LYS:HD3	3:H:222:GLU:OE2	2.21	0.40
1:B:121:PHE:CG	1:B:229:SER:HA	2.57	0.40
1:B:441:SER:CB	1:B:459:ASP:OD1	2.70	0.40
2:L:48:TYR:HB3	3:H:110:TYR:CD1	2.57	0.40
3:H:143:THR:OG1	3:H:144:GLY:N	2.54	0.40
3:H:93:THR:O	3:H:94:ALA:HB2	2.20	0.40
2:F:47:ILE:HD11	2:F:50:THR:C	2.46	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:2:NAG:O6	4:D:2:NAG:O6[2_655]	2.12	0.08

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/386 (100%)	366 (95%)	18 (5%)	0	100	100
1	B	384/386 (100%)	366 (95%)	18 (5%)	0	100	100
2	F	210/212 (99%)	205 (98%)	5 (2%)	0	100	100
2	L	209/212 (99%)	203 (97%)	6 (3%)	0	100	100
3	E	222/224 (99%)	209 (94%)	13 (6%)	0	100	100
3	H	222/224 (99%)	205 (92%)	17 (8%)	0	100	100
All	All	1631/1644 (99%)	1554 (95%)	77 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/333 (100%)	321 (96%)	12 (4%)	31	66
1	B	333/333 (100%)	320 (96%)	13 (4%)	28	64
2	F	187/187 (100%)	181 (97%)	6 (3%)	34	70
2	L	186/187 (100%)	182 (98%)	4 (2%)	45	78
3	E	192/192 (100%)	184 (96%)	8 (4%)	26	61
3	H	192/192 (100%)	182 (95%)	10 (5%)	21	53
All	All	1423/1424 (100%)	1370 (96%)	53 (4%)	30	65

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	SER
1	A	94	VAL
1	A	131	THR
1	A	217	LYS
1	A	231	CYS
1	A	255	ILE
1	A	259	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	262	LYS
1	A	297	HIS
1	A	346	VAL
1	A	365	ILE
1	A	366	SER
1	B	94	VAL
1	B	125	SER
1	B	131	THR
1	B	217	LYS
1	B	231	CYS
1	B	255	ILE
1	B	257	ARG
1	B	297	HIS
1	B	346	VAL
1	B	365	ILE
1	B	366	SER
1	B	386	ASN
1	B	398	GLU
2	F	11	LEU
2	F	24	ARG
2	F	47	ILE
2	F	77	VAL
2	F	89	GLN
2	F	142	ASP
3	E	3	LYS
3	E	6	GLU
3	E	48	VAL
3	E	66	VAL
3	E	185	LEU
3	E	188	LEU
3	E	195	THR
3	E	224	ARG
2	L	19	VAL
2	L	89	GLN
2	L	142	ASP
2	L	179	THR
3	H	6	GLU
3	H	7	SER
3	H	48	VAL
3	H	66	VAL
3	H	74	ARG
3	H	93	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	H	141	ASP
3	H	182	GLN
3	H	188	LEU
3	H	195	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	341	ASN
1	B	309	ASN
2	F	36	GLN
3	E	56	ASN
3	E	78	GLN
2	L	89	GLN
2	L	123	GLN
3	H	78	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	1	1,4	14,14,15	0.66	0	17,19,21	1.37	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	2	4	14,14,15	0.78	0	17,19,21	1.59	4 (23%)
4	NAG	D	1	1,4	14,14,15	0.65	0	17,19,21	1.37	2 (11%)
4	NAG	D	2	4	14,14,15	0.78	0	17,19,21	1.59	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	C	2	4	-	2/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	C1-O5-C5	3.99	117.54	112.19
4	C	1	NAG	C1-O5-C5	3.98	117.52	112.19
4	D	1	NAG	C1-C2-N2	-3.32	105.21	110.43
4	C	1	NAG	C1-C2-N2	-3.31	105.22	110.43
4	D	2	NAG	O7-C7-C8	-2.99	116.73	122.05
4	C	2	NAG	O7-C7-C8	-2.97	116.76	122.05
4	C	2	NAG	C2-N2-C7	2.48	126.23	122.90
4	D	2	NAG	C2-N2-C7	2.46	126.20	122.90
4	C	2	NAG	O3-C3-C2	2.42	114.42	109.40
4	D	2	NAG	O3-C3-C2	2.40	114.39	109.40
4	D	2	NAG	C4-C3-C2	-2.17	107.84	111.02
4	C	2	NAG	C4-C3-C2	-2.16	107.86	111.02

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	2	NAG	O5-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	C	2	NAG	C4-C5-C6-O6
4	C	1	NAG	C4-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
4	C	1	NAG	O5-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	2	NAG	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	502	1	14,14,15	0.51	0	17,19,21	1.50	3 (17%)
5	NAG	A	503	1	14,14,15	0.87	0	17,19,21	2.23	2 (11%)
5	NAG	B	501	1	14,14,15	0.60	0	17,19,21	1.73	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	504	1	14,14,15	0.57	0	17,19,21	1.67	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	502	1	-	0/6/23/26	0/1/1/1
5	NAG	A	503	1	-	1/6/23/26	0/1/1/1
5	NAG	B	501	1	-	0/6/23/26	0/1/1/1
5	NAG	A	504	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	503	NAG	C1-O5-C5	7.63	122.41	112.19
5	A	503	NAG	O5-C5-C6	3.59	114.66	107.66
5	B	501	NAG	C1-C2-N2	3.53	115.99	110.43
5	B	501	NAG	O7-C7-C8	-3.21	116.35	122.05
5	B	502	NAG	C2-N2-C7	2.98	126.89	122.90
5	A	504	NAG	O5-C5-C6	2.95	113.41	107.66
5	A	504	NAG	C1-O5-C5	2.82	115.96	112.19
5	B	501	NAG	C1-O5-C5	2.79	115.92	112.19
5	B	501	NAG	O5-C5-C6	2.69	112.90	107.66
5	A	504	NAG	O7-C7-C8	-2.21	118.12	122.05
5	A	504	NAG	C3-C4-C5	-2.20	106.24	110.23
5	B	502	NAG	O5-C5-C6	2.12	111.78	107.66
5	B	502	NAG	O5-C5-C4	-2.11	105.70	110.83
5	A	504	NAG	C4-C3-C2	-2.06	108.01	111.02
5	A	504	NAG	O4-C4-C5	2.01	114.28	109.32

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	NAG	C1-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	502	NAG	1	0
5	A	504	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/386 (100%)	-0.38	3 (0%) 82 75	29, 40, 62, 109	0
1	B	386/386 (100%)	-0.32	2 (0%) 87 82	27, 40, 65, 107	0
2	F	212/212 (100%)	-0.21	0 100 100	32, 49, 69, 118	0
2	L	211/212 (99%)	-0.19	0 100 100	34, 48, 72, 94	0
3	E	224/224 (100%)	-0.13	0 100 100	31, 50, 92, 140	0
3	H	224/224 (100%)	-0.17	2 (0%) 81 74	31, 48, 79, 135	0
All	All	1643/1644 (99%)	-0.26	7 (0%) 88 84	27, 44, 76, 140	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	ILE	3.4
1	B	149	ILE	3.2
1	A	442	SER	2.5
1	B	148	THR	2.3
3	H	142	THR	2.2
1	A	148	THR	2.1
3	H	5	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	C	1	14/15	-	-	66,78,85,89	0
4	NAG	C	2	14/15	-	-	66,77,82,82	0
4	NAG	D	1	14/15	-	-	66,78,85,88	0
4	NAG	D	2	14/15	-	-	66,77,82,82	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	503	14/15	0.73	0.14	79,89,95,98	0
5	NAG	B	502	14/15	0.80	0.11	63,72,87,93	0
5	NAG	A	504	14/15	0.83	0.12	66,77,87,90	0
5	NAG	B	501	14/15	0.85	0.12	69,85,91,96	0
6	CA	A	506	1/1	0.89	0.09	91,91,91,91	0
6	CA	B	505	1/1	0.95	0.14	72,72,72,72	0
6	CA	B	506	1/1	0.95	0.09	88,88,88,88	0
6	CA	A	505	1/1	0.96	0.15	75,75,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.