



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:41 AM UTC

PDB ID : 6QN6 / pdb_00006qn6
Title : Three dimensional structure of human carbonic anhydrase IX in complex with benzenesulfonamide
Authors : Leitans, J.; Tars, K.
Deposited on : 2019-02-09
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	2% 79% 16% ..
1	B	257	3% 81% 14% ...
1	C	257	88% 8% ..
1	D	257	2% 81% 14% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8312 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	0	0
			1929	1230	339	356	4			
1	B	251	Total	C	N	O	S	0	0	0
			1939	1236	343	356	4			
1	C	251	Total	C	N	O	S	0	0	0
			1939	1236	343	356	4			
1	D	251	Total	C	N	O	S	0	0	0
			1935	1233	342	356	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q16790
A	0	PRO	-	expression tag	UNP Q16790
A	41	SER	CYS	engineered mutation	UNP Q16790
B	-1	GLY	-	expression tag	UNP Q16790
B	0	PRO	-	expression tag	UNP Q16790
B	41	SER	CYS	engineered mutation	UNP Q16790
C	-1	GLY	-	expression tag	UNP Q16790
C	0	PRO	-	expression tag	UNP Q16790
C	41	SER	CYS	engineered mutation	UNP Q16790
D	-1	GLY	-	expression tag	UNP Q16790
D	0	PRO	-	expression tag	UNP Q16790
D	41	SER	CYS	engineered mutation	UNP Q16790

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		

- Molecule 3 is {N}-butyl-4-chloranyl-2-(2-phenylethylsulfanyl)-5-sulfamoyl-benzamide (CCD ID: J8Q) (formula: C₁₉H₂₃ClN₂O₃S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	S	0	0
			27	19	1	2	3	2		
3	B	1	Total	C	Cl	N	O	S	0	0
			27	19	1	2	3	2		
3	C	1	Total	C	Cl	N	O	S	0	0
			27	19	1	2	3	2		
3	D	1	Total	C	Cl	N	O	S	0	0
			27	19	1	2	3	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	98	Total	O	0	0
			98	98		
4	B	97	Total	O	0	0
			97	97		
4	C	165	Total	O	0	0
			165	165		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	98	Total	O	0	0
			98	98		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase 9





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	152.19Å 152.19Å 172.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.90 – 2.25 47.90 – 2.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.90-2.25) 100.0 (47.90-2.25)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.171 , 0.217 (Not available) , 0.214	Depositor DCC
R_{free} test set	3447 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	44.7	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8312	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: J8Q, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.30	3/1987 (0.2%)	1.41	7/2717 (0.3%)
1	B	1.48	5/1997 (0.3%)	1.63	23/2728 (0.8%)
1	C	1.38	4/1997 (0.2%)	1.64	13/2728 (0.5%)
1	D	1.24	4/1993 (0.2%)	1.36	5/2724 (0.2%)
All	All	1.35	16/7974 (0.2%)	1.52	48/10897 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	1
1	D	0	2
All	All	0	6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	82	GLY	C-N	31.59	1.69	1.33
1	C	11	PRO	C-N	25.10	1.64	1.33
1	A	125	THR	C-N	23.61	1.66	1.33
1	B	125	THR	C-N	22.95	1.69	1.33
1	B	72	PRO	C-N	21.68	1.67	1.33
1	C	72	PRO	C-N	18.71	1.59	1.33
1	C	125	THR	C-N	16.08	1.57	1.33
1	D	82	GLY	C-N	15.57	1.51	1.33
1	D	72	PRO	C-N	15.43	1.56	1.33
1	A	82	GLY	C-N	14.77	1.51	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	11	PRO	C-N	11.98	1.61	1.33
1	D	125	THR	C-N	11.91	1.51	1.33
1	D	11	PRO	C-N	9.70	1.56	1.33
1	C	82	GLY	C-N	8.48	1.43	1.33
1	B	11	PRO	CA-C	6.99	1.58	1.52
1	B	11	PRO	C-N	6.33	1.48	1.33

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	11	PRO	CA-C-N	-32.12	88.23	120.03
1	C	11	PRO	C-N-CA	-32.12	88.23	120.03
1	B	11	PRO	CA-C-N	-24.26	89.51	119.84
1	B	11	PRO	C-N-CA	-24.26	89.51	119.84
1	B	82	GLY	O-C-N	-15.42	106.35	121.77
1	A	10	ASP	O-C-N	11.86	130.30	121.88
1	A	11	PRO	O-C-N	-11.19	99.83	121.10
1	B	125	THR	O-C-N	-10.86	109.52	122.22
1	B	10	ASP	C-N-CD	-10.67	81.25	125.00
1	D	11	PRO	C-N-CD	-10.20	83.17	125.00
1	C	213	ASN	CA-CB-CG	10.06	122.66	112.60
1	B	11	PRO	CA-C-O	-9.94	106.15	120.56
1	B	82	GLY	CA-C-N	-8.45	110.58	120.13
1	B	82	GLY	C-N-CA	-8.45	110.58	120.13
1	A	11	PRO	N-CA-C	8.12	133.22	112.10
1	B	72	PRO	CA-C-N	-8.03	109.92	122.69
1	B	72	PRO	C-N-CA	-8.03	109.92	122.69
1	C	11	PRO	O-C-N	7.63	135.59	121.10
1	B	11	PRO	C-N-CD	7.39	155.29	125.00
1	A	11	PRO	CA-C-N	-7.23	110.80	119.84
1	A	11	PRO	C-N-CA	-7.23	110.80	119.84
1	C	58	ARG	CG-CD-NE	-7.19	96.18	112.00
1	B	10	ASP	CA-C-N	7.05	127.64	120.38
1	B	10	ASP	C-N-CA	7.05	127.64	120.38
1	A	11	PRO	C-N-CD	-6.63	97.82	125.00
1	B	72	PRO	O-C-N	6.54	131.12	123.01
1	D	58	ARG	CG-CD-NE	-6.39	97.94	112.00
1	C	212	PHE	CA-C-N	6.34	128.69	120.44
1	C	212	PHE	C-N-CA	6.34	128.69	120.44
1	C	34	ARG	CB-CA-C	6.12	117.01	110.65
1	C	213	ASN	N-CA-C	-6.09	104.55	111.07
1	D	11	PRO	CA-C-O	-5.98	105.86	120.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	102	ARG	CB-CA-C	5.95	116.49	109.47
1	C	10	ASP	O-C-N	5.88	128.08	121.32
1	D	10	ASP	O-C-N	5.88	127.35	121.30
1	B	232	PRO	CA-N-CD	-5.86	103.79	112.00
1	B	11	PRO	CA-N-CD	-5.82	103.85	112.00
1	B	221	LYS	CA-C-N	5.78	128.34	120.54
1	B	221	LYS	C-N-CA	5.78	128.34	120.54
1	B	213	ASN	CA-CB-CG	5.76	118.36	112.60
1	B	247	ALA	CA-C-N	5.56	128.60	120.71
1	B	247	ALA	C-N-CA	5.56	128.60	120.71
1	C	89	ARG	CB-CA-C	5.40	119.06	109.72
1	C	212	PHE	CA-C-O	-5.30	115.22	121.16
1	B	15	PRO	CA-N-CD	-5.16	104.78	112.00
1	B	40	PHE	CA-C-O	-5.11	114.83	120.70
1	D	11	PRO	N-CA-C	5.03	125.19	112.10
1	C	197	SER	CA-C-O	-5.01	115.90	121.51

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	PRO	Mainchain
1	B	10	ASP	Peptide
1	B	11	PRO	Mainchain
1	C	82	GLY	Mainchain
1	D	11	PRO	Mainchain
1	D	82	GLY	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1929	0	1864	24	0
1	B	1939	0	1886	24	0
1	C	1939	0	1886	11	0
1	D	1935	0	1875	21	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	27	0	0	0	0
3	B	27	0	0	0	0
3	C	27	0	0	1	0
3	D	27	0	0	0	0
4	A	98	0	0	2	0
4	B	97	0	0	0	0
4	C	165	0	0	1	0
4	D	98	0	0	3	0
All	All	8312	0	7511	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:THR:C	1:B:127:ALA:N	1.69	1.50
1:B:82:GLY:C	1:B:84:PRO:N	1.68	1.47
1:B:82:GLY:C	1:B:84:PRO:CA	2.58	0.76
1:B:82:GLY:CA	1:B:84:PRO:N	2.48	0.75
1:C:205:GLN:NE2	4:C:401:HOH:O	2.16	0.68
1:B:125:THR:C	1:B:127:ALA:CA	2.67	0.67
1:B:11:PRO:O	1:B:15:PRO:C	2.25	0.67
1:C:10:ASP:HB3	1:C:11:PRO:HA	1.76	0.67
1:A:135:LEU:O	1:C:58:ARG:NH2	2.31	0.62
1:B:11:PRO:C	1:B:15:PRO:O	2.39	0.62
1:B:148:LEU:HD22	1:B:216:VAL:CG2	2.30	0.62
1:A:170:GLU:OE2	1:A:232:PRO:HA	2.01	0.60
1:D:214:GLN:NE2	4:D:402:HOH:O	2.35	0.58
1:A:131:VAL:HG23	4:A:456:HOH:O	2.02	0.57
1:A:11:PRO:HB2	1:A:15:PRO:HD2	1.89	0.54
1:A:54:PRO:O	1:A:178:GLY:HA3	2.08	0.54
1:A:15:PRO:HG2	1:A:18:ARG:HD3	1.89	0.54
1:A:193:GLN:HA	1:A:209:TRP:O	2.08	0.53
1:B:54:PRO:O	1:B:178:GLY:HA3	2.08	0.53
1:D:163:ARG:NH1	1:D:166:GLU:OE1	2.41	0.53
1:D:163:ARG:NH1	1:D:166:GLU:OE2	2.41	0.53
1:B:170:GLU:OE2	1:B:232:PRO:HD3	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:THR:CA	1:B:127:ALA:N	2.68	0.52
1:A:163:ARG:HD2	1:A:166:GLU:OE1	2.10	0.52
1:C:53:LEU:HD21	1:C:77:LEU:HD11	1.92	0.51
1:B:130:ARG:NH2	1:D:169:GLU:OE1	2.45	0.50
1:D:36:GLN:NE2	1:D:110:GLU:OE2	2.45	0.49
1:B:10:ASP:CG	1:B:15:PRO:HD3	2.37	0.49
1:A:32:ASP:OD1	1:A:108:THR:OG1	2.25	0.49
1:A:99:ALA:O	1:A:100:ALA:C	2.57	0.48
1:A:10:ASP:CB	1:A:11:PRO:HA	2.43	0.48
1:C:198:LEU:HD22	3:C:302:J8Q:CL1	2.50	0.48
1:D:93:LEU:C	1:D:93:LEU:HD12	2.40	0.47
1:D:30:PRO:HG3	1:D:106:GLU:HB3	1.96	0.47
1:D:205:GLN:NE2	4:D:401:HOH:O	2.34	0.47
1:D:163:ARG:NH1	1:D:166:GLU:CD	2.73	0.47
1:B:10:ASP:HB3	1:B:15:PRO:CD	2.45	0.46
1:D:136:GLY:HA2	1:D:205:GLN:O	2.15	0.46
1:A:169:GLU:O	1:A:172:SER:OG	2.33	0.46
1:D:98:GLY:HA2	1:D:243:LEU:HD23	1.97	0.46
1:D:167:ILE:HA	1:D:172:SER:OG	2.15	0.46
1:B:82:GLY:O	1:B:84:PRO:N	2.35	0.46
1:B:88:TYR:HB3	1:B:122:HIS:HB3	1.98	0.46
1:C:93:LEU:HA	1:C:119:HIS:O	2.16	0.46
1:B:167:ILE:HA	1:B:172:SER:OG	2.15	0.46
1:C:88:TYR:HB3	1:C:122:HIS:HB3	1.98	0.45
1:A:54(B):LEU:HA	1:A:55:PRO:C	2.41	0.45
1:A:56:GLU:C	1:A:57:LEU:HD12	2.42	0.45
1:A:98:GLY:N	1:A:104:GLY:HA3	2.32	0.44
1:B:48:GLU:O	1:B:79:MET:HA	2.17	0.44
1:A:132:ASP:HB2	1:C:173:GLU:O	2.17	0.44
1:A:16:TRP:N	1:A:17:PRO:CD	2.80	0.44
1:D:11:PRO:O	1:D:15:PRO:O	2.36	0.43
1:A:97:TRP:C	1:A:104:GLY:HA3	2.42	0.43
1:C:10:ASP:CB	1:C:11:PRO:HA	2.36	0.43
1:D:88:TYR:HB3	1:D:122:HIS:HB3	1.99	0.43
1:D:49:LEU:HB3	1:D:51:PHE:CD1	2.54	0.43
1:D:97:TRP:CE3	1:D:226:LEU:HD11	2.53	0.43
1:B:197:SER:HA	1:B:204:ALA:O	2.19	0.43
1:A:132:ASP:HA	1:A:135:LEU:HD12	2.00	0.42
1:B:59:LEU:HD11	1:B:66:VAL:HG22	2.00	0.42
1:B:227(A):ASP:O	1:B:240:ARG:NH2	2.53	0.42
1:D:163:ARG:HH11	1:D:166:GLU:CD	2.26	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:GLN:OE1	1:A:203:CYS:HB3	2.20	0.42
1:A:116:ALA:HB3	1:A:148:LEU:HB2	2.02	0.42
1:A:88:TYR:HB3	1:A:122:HIS:HB3	2.02	0.41
1:A:125:THR:HG21	4:A:439:HOH:O	2.19	0.41
1:B:53:LEU:HD21	1:B:77:LEU:HD11	2.02	0.41
1:A:77:LEU:O	1:A:89:ARG:HA	2.20	0.41
1:D:159:GLN:NE2	4:D:403:HOH:O	2.44	0.41
1:B:7:TYR:CD2	1:B:244:ASN:HA	2.55	0.41
1:C:219:SER:OG	1:C:222:GLN:HG3	2.21	0.41
1:D:58:ARG:HD3	1:D:173:GLU:OE2	2.20	0.41
1:D:11:PRO:HA	1:D:15:PRO:HD3	1.78	0.40
1:B:20:SER:HA	1:B:21:PRO:HD3	1.96	0.40
1:D:54:PRO:O	1:D:178:GLY:HA3	2.21	0.40
1:C:45:ARG:O	1:C:191:TYR:OH	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/257 (94%)	231 (96%)	8 (3%)	2 (1%)	16	14
1	B	249/257 (97%)	239 (96%)	8 (3%)	2 (1%)	16	14
1	C	239/257 (93%)	235 (98%)	4 (2%)	0	100	100
1	D	190/257 (74%)	179 (94%)	11 (6%)	0	100	100
All	All	919/1028 (89%)	884 (96%)	31 (3%)	4 (0%)	30	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	ALA
1	B	15	PRO
1	B	11	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/210 (96%)	194 (96%)	7 (4%)	32	39
1	B	203/210 (97%)	196 (97%)	7 (3%)	32	40
1	C	203/210 (97%)	202 (100%)	1 (0%)	81	87
1	D	202/210 (96%)	190 (94%)	12 (6%)	18	18
All	All	809/840 (96%)	782 (97%)	27 (3%)	33	42

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	LEU
1	A	92	GLN
1	A	110	GLU
1	A	170	GLU
1	A	216	VAL
1	A	241	LEU
1	A	248	THR
1	B	57	LEU
1	B	92	GLN
1	B	173	GLU
1	B	213	ASN
1	B	232	PRO
1	B	241	LEU
1	B	248	THR
1	C	92	GLN
1	D	17	PRO
1	D	34	ARG
1	D	36	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	50	LEU
1	D	66	VAL
1	D	87	GLU
1	D	92	GLN
1	D	132	ASP
1	D	163	ARG
1	D	227	SER
1	D	227(A)	ASP
1	D	248	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	159	GLN
1	A	193	GLN
1	A	224	HIS
1	B	36	GLN
1	B	64	HIS
1	B	67	GLN
1	B	159	GLN
1	C	64	HIS
1	C	67	GLN
1	C	242	GLN
1	D	213	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	J8Q	B	302	2	28,28,28	2.65	5 (17%)	38,38,38	2.02	10 (26%)
3	J8Q	D	302	2	28,28,28	2.78	7 (25%)	38,38,38	2.32	10 (26%)
3	J8Q	A	302	2	28,28,28	2.61	4 (14%)	38,38,38	2.94	14 (36%)
3	J8Q	C	302	2	28,28,28	2.59	6 (21%)	38,38,38	2.83	15 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	J8Q	B	302	2	-	2/21/21/21	0/2/2/2
3	J8Q	D	302	2	-	2/21/21/21	0/2/2/2
3	J8Q	A	302	2	-	2/21/21/21	0/2/2/2
3	J8Q	C	302	2	-	3/21/21/21	0/2/2/2

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	302	J8Q	C7-S4	-9.77	1.65	1.77
3	C	302	J8Q	C7-S4	-9.52	1.65	1.77
3	A	302	J8Q	C7-S4	-9.34	1.66	1.77
3	B	302	J8Q	C7-S4	-8.60	1.67	1.77
3	B	302	J8Q	C8-C7	7.58	1.50	1.40
3	D	302	J8Q	C8-C7	7.14	1.49	1.40
3	A	302	J8Q	C8-C7	6.49	1.48	1.40
3	D	302	J8Q	C11-C10	6.04	1.50	1.40
3	A	302	J8Q	C11-C10	5.88	1.50	1.40
3	B	302	J8Q	C11-C10	5.08	1.49	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	302	J8Q	C11-C10	4.76	1.48	1.40
3	C	302	J8Q	C8-C7	4.75	1.46	1.40
3	B	302	J8Q	C8-CL1	3.82	1.82	1.73
3	C	302	J8Q	C10-S14	-3.72	1.71	1.77
3	C	302	J8Q	C8-CL1	3.32	1.81	1.73
3	A	302	J8Q	C8-CL1	3.13	1.81	1.73
3	D	302	J8Q	C8-CL1	2.54	1.79	1.73
3	C	302	J8Q	S4-N1	2.49	1.65	1.60
3	D	302	J8Q	C10-S14	-2.30	1.73	1.77
3	D	302	J8Q	S4-N1	2.13	1.64	1.60
3	B	302	J8Q	O6-S4	2.13	1.47	1.43
3	D	302	J8Q	C12-C7	2.01	1.42	1.39

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	J8Q	O6-S4-O5	-9.73	103.83	118.80
3	C	302	J8Q	O6-S4-O5	-8.64	105.51	118.80
3	C	302	J8Q	O6-S4-C7	7.95	118.82	107.26
3	A	302	J8Q	O6-S4-C7	7.03	117.48	107.26
3	A	302	J8Q	C12-C7-S4	6.92	126.54	118.26
3	D	302	J8Q	O5-S4-N1	6.92	117.33	107.35
3	C	302	J8Q	C12-C7-S4	6.21	125.69	118.26
3	D	302	J8Q	O6-S4-O5	-5.77	109.92	118.80
3	B	302	J8Q	O6-S4-O5	-5.43	110.44	118.80
3	D	302	J8Q	O6-S4-N1	-4.84	100.37	107.35
3	B	302	J8Q	C12-C7-S4	4.82	124.02	118.26
3	A	302	J8Q	C11-C10-S14	4.55	125.66	120.36
3	B	302	J8Q	C7-S4-N1	4.45	116.28	108.28
3	C	302	J8Q	O5-S4-N1	-4.34	101.10	107.35
3	C	302	J8Q	C8-C7-S4	-4.29	118.32	123.47
3	A	302	J8Q	C15-S14-C10	4.26	111.91	103.05
3	D	302	J8Q	O24-C23-C11	-4.23	113.28	121.03
3	A	302	J8Q	C8-C7-S4	-4.22	118.41	123.47
3	D	302	J8Q	C12-C7-S4	3.88	122.90	118.26
3	B	302	J8Q	C12-C7-C8	-3.84	114.31	118.33
3	A	302	J8Q	C7-S4-N1	3.81	115.12	108.28
3	A	302	J8Q	C7-C8-CL1	-3.72	118.83	121.52
3	C	302	J8Q	C7-C8-CL1	-3.63	118.89	121.52
3	D	302	J8Q	C11-C23-N25	3.56	124.23	117.32
3	A	302	J8Q	C12-C7-C8	-3.35	114.82	118.33
3	B	302	J8Q	C15-S14-C10	3.23	109.77	103.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	302	J8Q	C15-S14-C10	3.22	109.75	103.05
3	A	302	J8Q	C8-C9-C10	3.16	123.55	120.27
3	B	302	J8Q	O24-C23-C11	-3.11	115.34	121.03
3	B	302	J8Q	C8-C9-C10	3.04	123.42	120.27
3	C	302	J8Q	C11-C10-S14	2.95	123.81	120.36
3	C	302	J8Q	C15-S14-C10	2.95	109.19	103.05
3	D	302	J8Q	C12-C7-C8	-2.93	115.27	118.33
3	B	302	J8Q	O5-S4-C7	2.83	111.37	107.26
3	C	302	J8Q	C12-C7-C8	-2.59	115.62	118.33
3	C	302	J8Q	C27-N25-C23	2.54	127.84	122.11
3	D	302	J8Q	O6-S4-C7	2.50	110.89	107.26
3	C	302	J8Q	O24-C23-N25	2.50	127.47	122.59
3	A	302	J8Q	C9-C10-C11	-2.39	114.88	119.62
3	B	302	J8Q	C9-C10-C11	-2.22	115.21	119.62
3	C	302	J8Q	C29-C28-C27	2.20	125.28	114.01
3	C	302	J8Q	C9-C8-C7	2.19	123.48	121.36
3	C	302	J8Q	C9-C10-S14	-2.19	119.28	122.77
3	A	302	J8Q	C9-C10-S14	-2.15	119.35	122.77
3	C	302	J8Q	O5-S4-C7	2.13	110.36	107.26
3	A	302	J8Q	O5-S4-N1	-2.12	104.30	107.35
3	D	302	J8Q	C7-C8-CL1	-2.05	120.03	121.52
3	A	302	J8Q	O24-C23-C11	-2.04	117.30	121.03
3	B	302	J8Q	C12-C11-C10	2.01	121.61	118.68

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	302	J8Q	N25-C27-C28-C29
3	D	302	J8Q	N25-C27-C28-C29
3	A	302	J8Q	N25-C27-C28-C29
3	C	302	J8Q	N25-C27-C28-C29
3	D	302	J8Q	C12-C7-S4-N1
3	B	302	J8Q	C12-C7-S4-N1
3	C	302	J8Q	C8-C7-S4-N1
3	A	302	J8Q	C12-C7-S4-N1
3	C	302	J8Q	C27-C28-C29-C30

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	302	J8Q	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand J8Q D 302




Bond lengths



Bond angles




Torsions




Rings


Ligand J8Q A 302




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	3
1	A	2
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	125:THR	C	127:ALA	N	1.69
1	B	82:GLY	C	84:PRO	N	1.68
1	A	125:THR	C	127:ALA	N	1.66
1	B	72:PRO	C	76:GLY	N	1.66
1	C	11:PRO	C	15:PRO	N	1.64
1	A	11:PRO	C	15:PRO	N	1.61

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/257 (97%)	-0.01	5 (1%) 65 65	36, 54, 86, 114	0
1	B	251/257 (97%)	-0.07	8 (3%) 50 50	33, 47, 96, 147	0
1	C	251/257 (97%)	-0.40	1 (0%) 88 89	30, 42, 68, 96	0
1	D	251/257 (97%)	-0.11	5 (1%) 65 65	36, 52, 81, 102	0
All	All	1004/1028 (97%)	-0.15	19 (1%) 66 67	30, 48, 86, 147	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	PRO	5.6
1	B	9	GLY	5.5
1	B	11	PRO	4.3
1	D	11	PRO	4.1
1	B	19	VAL	3.2
1	B	238	ASP	3.0
1	B	8	GLY	3.0
1	D	238	ASP	2.8
1	C	238	ASP	2.7
1	B	102	ARG	2.6
1	A	232	PRO	2.5
1	A	238	ASP	2.5
1	A	230	TRP	2.4
1	D	15	PRO	2.3
1	A	15	PRO	2.3
1	A	9	GLY	2.3
1	D	189	PHE	2.1
1	B	101	GLY	2.1
1	D	101	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	J8Q	D	302	27/27	0.95	0.13	40,54,82,93	0
3	J8Q	C	302	27/27	0.96	0.10	38,50,62,68	0
3	J8Q	A	302	27/27	0.97	0.10	41,52,93,99	0
3	J8Q	B	302	27/27	0.97	0.10	35,49,82,100	0
2	ZN	A	301	1/1	1.00	0.02	45,45,45,45	0
2	ZN	B	301	1/1	1.00	0.01	43,43,43,43	0
2	ZN	C	301	1/1	1.00	0.01	37,37,37,37	0
2	ZN	D	301	1/1	1.00	0.01	42,42,42,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around J8Q D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J8Q C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J8Q A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around J8Q B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.