



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:24 AM UTC

PDB ID : 6QNS / pdb_00006qns
Title : Crystal structure of the binding domain of Botulinum Neurotoxin type B mutant I1248W/V1249W in complex with human synaptotagmin 1 and GD1a receptors
Authors : Masuyer, G.; Yin, L.; Zhang, S.; Miyashita, S.I.; Dong, M.; Stenmark, P.
Deposited on : 2019-02-12
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	2% 88% 9% .
2	S	21	10% 52% 24% 24%
3	B	6	17% 67% 17%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Botulinum neurotoxin type B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	0	0
			3676	2377	601	690	8			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	850	GLY	-	expression tag	UNP P10844
A	851	SER	-	expression tag	UNP P10844
A	852	HIS	-	expression tag	UNP P10844
A	853	MET	-	expression tag	UNP P10844
A	854	ALA	-	expression tag	UNP P10844
A	855	SER	-	expression tag	UNP P10844
A	856	MET	-	expression tag	UNP P10844
A	1248	TRP	ILE	engineered mutation	UNP P10844
A	1249	TRP	VAL	engineered mutation	UNP P10844

- Molecule 2 is a protein called Synaptotagmin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	S	16	Total	C	N	O	S	0	0	0
			137	88	22	26	1			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[N-acetyl-alpha-neuraminic acid-(2-3)]beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	6	Total	C	N	O	0	0	0
			88	48	3	37			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	85	Total	O	0	0
			85	85		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Botulinum neurotoxin type B



- Molecule 2: Synaptotagmin-1



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[N-acetyl-alpha-neuraminic acid-(2-3)]beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.34Å 78.49Å 149.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.55 – 2.40 74.55 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (74.55-2.40) 99.9 (74.55-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.40Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.199 , 0.229 0.203 , 0.232	Depositor DCC
R_{free} test set	1205 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.844	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3986	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, SIA, NGA, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/3768	1.29	6/5082 (0.1%)
2	S	1.03	0/139	1.62	0/182
All	All	0.99	0/3907	1.30	6/5264 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	977	ASP	CA-CB-CG	7.52	120.12	112.60
1	A	1033	GLU	CB-CA-C	-5.42	102.70	110.62
1	A	1003	PHE	CA-CB-CG	5.23	119.03	113.80
1	A	976	ILE	CA-C-O	-5.05	114.92	120.43
1	A	1053	PHE	CA-C-N	5.00	130.01	122.75
1	A	1053	PHE	C-N-CA	5.00	130.01	122.75

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1270	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3676	0	3570	32	0
2	S	137	0	132	13	0
3	B	88	0	74	1	0
4	A	85	0	0	1	0
All	All	3986	0	3776	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1008:THR:C	1:A:1009:ASN:HD22	1.75	0.93
1:A:1203:GLU:OE1	2:S:44:LYS:NZ	2.09	0.85
2:S:51:LEU:H	2:S:51:LEU:HD23	1.45	0.80
1:A:1008:THR:O	1:A:1009:ASN:ND2	2.16	0.78
2:S:48:MET:HA	2:S:51:LEU:HD21	1.74	0.68
1:A:1070:ASN:C	1:A:1070:ASN:HD22	2.02	0.65
1:A:1057:LYS:HD2	4:A:1463:HOH:O	1.99	0.62
1:A:1105:ASN:HB3	1:A:1108:SER:OG	2.04	0.58
2:S:47:PHE:O	2:S:51:LEU:CD2	2.51	0.58
1:A:1203:GLU:OE1	2:S:44:LYS:CE	2.51	0.57
2:S:47:PHE:O	2:S:51:LEU:HD22	2.03	0.57
1:A:1184:LYS:HE2	1:A:1205:TYR:CE1	2.40	0.56
1:A:1128:ASN:O	1:A:1128:ASN:ND2	2.40	0.55
1:A:1008:THR:C	1:A:1009:ASN:ND2	2.58	0.54
3:B:5:SIA:O4	3:B:5:SIA:H112	2.08	0.54
1:A:1070:ASN:C	1:A:1070:ASN:ND2	2.67	0.52
1:A:1203:GLU:CD	2:S:44:LYS:HE3	2.36	0.51
1:A:1084:LEU:CD1	1:A:1210:ILE:HG12	2.41	0.50
1:A:1156:ILE:O	1:A:1156:ILE:HG13	2.12	0.50
1:A:923:PHE:HE2	1:A:925:ASP:O	1.94	0.49
1:A:923:PHE:CE2	1:A:925:ASP:O	2.67	0.47
1:A:1203:GLU:OE1	2:S:44:LYS:HE3	2.14	0.47
1:A:1120:GLU:OE1	1:A:1254:LYS:NZ	2.43	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1109:TYR:CE2	1:A:1141:ILE:HD12	2.49	0.46
1:A:1260:SER:HB3	1:A:1263:TYR:CD2	2.50	0.46
1:A:1031:ILE:HG23	1:A:1031:ILE:O	2.16	0.45
2:S:44:LYS:O	2:S:48:MET:HG2	2.16	0.45
1:A:1001:ARG:NH2	1:A:1281:GLN:OE1	2.50	0.44
1:A:1247:GLY:HA3	1:A:1250:PHE:O	2.18	0.44
1:A:1028:ILE:O	1:A:1031:ILE:HG22	2.18	0.44
1:A:1199:SER:HB3	2:S:40:PHE:CE2	2.52	0.44
2:S:51:LEU:H	2:S:51:LEU:CD2	2.24	0.43
1:A:973:TRP:CD2	1:A:1007:ILE:HG21	2.53	0.43
1:A:1152:ASN:ND2	1:A:1156:ILE:HD11	2.34	0.42
1:A:1191:GLU:OE2	2:S:51:LEU:HB2	2.18	0.42
1:A:1056:MET:HE2	1:A:1056:MET:HB3	1.89	0.42
1:A:1199:SER:CB	2:S:40:PHE:CE2	3.02	0.42
1:A:923:PHE:CZ	1:A:925:ASP:HB3	2.56	0.40
1:A:960:SER:HB3	1:A:1035:ILE:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/442 (97%)	410 (96%)	18 (4%)	0	100	100
2	S	14/21 (67%)	13 (93%)	1 (7%)	0	100	100
All	All	442/463 (96%)	423 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/415 (98%)	404 (100%)	1 (0%)	87	94
2	S	15/18 (83%)	15 (100%)	0	100	100
All	All	420/433 (97%)	419 (100%)	1 (0%)	87	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1070	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	862	ASN
1	A	979	ASN
1	A	1128	ASN
1	A	1129	GLN
1	A	1130	ASN
1	A	1152	ASN
1	A	1154	GLN
1	A	1175	ASN
1	A	1176	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	B	1	3	12,12,12	0.71	0	17,17,17	1.02	1 (5%)
3	GAL	B	2	3	11,11,12	0.51	0	15,15,17	1.30	4 (26%)
3	NGA	B	3	3	14,14,15	0.34	0	17,19,21	0.99	1 (5%)
3	GAL	B	4	3	11,11,12	0.49	0	15,15,17	0.86	0
3	SIA	B	5	3	20,20,21	0.77	1 (5%)	21,28,31	1.28	2 (9%)
3	SIA	B	6	3	20,20,21	0.82	1 (5%)	21,28,31	0.99	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	B	1	3	-	2/2/22/22	0/1/1/1
3	GAL	B	2	3	-	0/2/19/22	0/1/1/1
3	NGA	B	3	3	-	1/6/23/26	0/1/1/1
3	GAL	B	4	3	-	0/2/19/22	0/1/1/1
3	SIA	B	5	3	-	9/18/34/38	0/1/1/1
3	SIA	B	6	3	-	10/18/34/38	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	6	SIA	O1B-C1	-2.83	1.21	1.30
3	B	5	SIA	O1B-C1	-2.47	1.22	1.30

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	B	5	SIA	C8-C7-C6	3.11	118.89	113.05
3	B	5	SIA	O1B-C1-C2	2.81	120.01	112.71
3	B	6	SIA	O1B-C1-C2	2.75	119.86	112.71
3	B	2	GAL	O3-C3-C4	2.58	116.46	110.38
3	B	2	GAL	C2-C3-C4	-2.34	106.74	110.86
3	B	1	BGC	C1-O5-C5	2.28	118.06	113.65
3	B	3	NGA	C1-O5-C5	2.26	115.21	112.19
3	B	2	GAL	O3-C3-C2	-2.19	105.58	110.05
3	B	2	GAL	C1-C2-C3	2.14	112.75	109.64

There are no chirality outliers.

All (22) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	5	SIA	C5-C6-C7-C8
3	B	5	SIA	C5-C6-C7-O7
3	B	5	SIA	O6-C6-C7-C8
3	B	5	SIA	O6-C6-C7-O7
3	B	5	SIA	O8-C8-C9-O9
3	B	5	SIA	C11-C10-N5-C5
3	B	6	SIA	C5-C6-C7-C8
3	B	6	SIA	C5-C6-C7-O7
3	B	6	SIA	O6-C6-C7-C8
3	B	6	SIA	O6-C6-C7-O7
3	B	6	SIA	C7-C8-C9-O9
3	B	5	SIA	O10-C10-N5-C5
3	B	1	BGC	C4-C5-C6-O6
3	B	6	SIA	O8-C8-C9-O9
3	B	1	BGC	O5-C5-C6-O6
3	B	5	SIA	C7-C8-C9-O9
3	B	6	SIA	C11-C10-N5-C5
3	B	6	SIA	O10-C10-N5-C5
3	B	3	NGA	C4-C5-C6-O6
3	B	6	SIA	C4-C5-N5-C10
3	B	5	SIA	C4-C5-N5-C10
3	B	6	SIA	O1A-C1-C2-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	5	SIA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/442 (97%)	0.07	10 (2%) 61 57	38, 62, 123, 170	0
2	S	16/21 (76%)	0.77	2 (12%) 8 6	59, 88, 112, 115	0
All	All	446/463 (96%)	0.10	12 (2%) 56 52	38, 63, 123, 170	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	923	PHE	4.0
1	A	924	LEU	3.4
1	A	1031	ILE	3.0
1	A	1130	ASN	2.9
1	A	1155	SER	2.8
1	A	918	ILE	2.7
1	A	919	PHE	2.5
2	S	41	SER	2.5
2	S	39	ALA	2.3
1	A	1156	ILE	2.2
1	A	1033	GLU	2.2
1	A	917	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BGC	B	1	12/12	-	-	114,139,144,146	0
3	GAL	B	2	11/12	-	-	96,103,106,109	0
3	NGA	B	3	14/15	-	-	76,85,92,93	0
3	GAL	B	4	11/12	-	-	66,69,73,75	0
3	SIA	B	5	20/21	-	-	65,74,86,88	0
3	SIA	B	6	20/21	-	-	109,132,143,146	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.