



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 10:06 AM EDT

PDB ID : 2QOM / pdb_00002qom
Title : The crystal structure of the E.coli EspP autotransporter Beta-domain.
Authors : Barnard, T.J.; Dautin, N.; Lukacik, P.; Bernstein, H.D.; Buchanan, S.K.
Deposited on : 2007-07-20
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	
1	B	285	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine protease espP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	2	0
			2002	1261	344	391	6			
1	B	269	Total	C	N	O	S	0	1	0
			2033	1285	351	391	6			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1301	GLY	-	expression tag	UNP Q7BSW5
A	1302	SER	-	expression tag	UNP Q7BSW5
A	1303	HIS	-	expression tag	UNP Q7BSW5
A	1304	HIS	-	expression tag	UNP Q7BSW5
A	1305	HIS	-	expression tag	UNP Q7BSW5
A	1306	HIS	-	expression tag	UNP Q7BSW5
A	1307	HIS	-	expression tag	UNP Q7BSW5
A	1308	HIS	-	expression tag	UNP Q7BSW5
B	1301	GLY	-	expression tag	UNP Q7BSW5
B	1302	SER	-	expression tag	UNP Q7BSW5
B	1303	HIS	-	expression tag	UNP Q7BSW5
B	1304	HIS	-	expression tag	UNP Q7BSW5
B	1305	HIS	-	expression tag	UNP Q7BSW5
B	1306	HIS	-	expression tag	UNP Q7BSW5
B	1307	HIS	-	expression tag	UNP Q7BSW5
B	1308	HIS	-	expression tag	UNP Q7BSW5


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	O	0	0
			4	4		
2	B	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serine protease espP



• Molecule 1: Serine protease espP



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.95Å 53.30Å 102.42Å 90.00° 103.57° 90.00°	Depositor
Resolution (Å)	15.00 – 2.66 15.00 – 2.66	Depositor EDS
% Data completeness (in resolution range)	95.4 (15.00-2.66) 94.7 (15.00-2.66)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.258 , 0.284 0.276 , 0.309	Depositor DCC
R_{free} test set	1252 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	60.4	Xtriage
Anisotropy	1.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 62.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4041	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	1/2050 (0.0%)	0.85	2/2765 (0.1%)
1	B	0.53	0/2078	0.77	0/2800
All	All	0.59	1/4128 (0.0%)	0.81	2/5565 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1
1	B	0	2
All	All	1	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1032	LEU	N-CA	5.15	1.52	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1247	GLU	N-CA-C	8.05	127.94	110.80
1	A	1068	ASP	N-CA-C	5.15	115.34	108.38

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	1247	GLU	CA

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1129	ASP	Peptide
1	B	1129	ASP	Peptide
1	B	1141	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	1819	26	0
1	B	2033	0	1860	34	0
2	A	4	0	0	0	0
2	B	2	0	0	0	0
All	All	4041	0	3679	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1141:THR:HG23	1:B:1142:ARG:HA	1.52	0.89
1:B:1130:ASN:HD21	1:B:1146:THR:HG22	1.37	0.89
1:B:1130:ASN:ND2	1:B:1146:THR:HG22	1.91	0.84
1:A:1032:LEU:HD22	1:A:1079:PHE:CZ	2.14	0.81
1:B:1293:ASN:C	1:B:1293:ASN:HD22	1.86	0.81
1:A:1032:LEU:HD22	1:A:1079:PHE:CE1	2.20	0.76
1:B:1141:THR:HG23	1:B:1142:ARG:CA	2.16	0.75
1:A:1029:MET:HE2	1:A:1032:LEU:HD11	1.75	0.68
1:B:1209:ASP:OD1	1:B:1226:GLY:HA3	1.94	0.67
1:A:1130:ASN:HB2	1:A:1144:TYR:O	1.99	0.62
1:B:1293:ASN:C	1:B:1293:ASN:ND2	2.56	0.62
1:A:1068:ASP:HB3	1:A:1108:TYR:OH	2.00	0.61
1:A:1032:LEU:HD22	1:A:1079:PHE:CE2	2.36	0.60
1:B:1132:TYR:O	1:B:1142:ARG:N	2.34	0.60
1:A:1243:ASP:OD1	1:A:1245:SER:HB3	2.01	0.60
1:A:1024:ASN:HD22	1:A:1027:LYS:HB2	1.69	0.57
1:A:1032:LEU:HD13	1:A:1079:PHE:CD2	2.40	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1263:GLY:C	1:A:1264:LEU:HD12	2.29	0.57
1:A:1241:LEU:HD12	1:A:1242:ARG:N	2.21	0.56
1:A:1032:LEU:HB3	1:A:1079:PHE:CE2	2.41	0.55
1:B:1029:MET:HE3	1:B:1032:LEU:HD21	1.87	0.55
1:B:1243:ASP:OD1	1:B:1245:SER:OG	2.20	0.55
1:B:1070:LYS:HG3	1:B:1079:PHE:CE1	2.43	0.54
1:B:1141:THR:CG2	1:B:1142:ARG:HA	2.32	0.52
1:A:1161:VAL:HG21	1:B:1206:THR:HG22	1.90	0.51
1:B:1130:ASN:HD21	1:B:1146:THR:CG2	2.18	0.50
1:B:1146:THR:OG1	1:B:1179:SER:O	2.25	0.50
1:B:1141:THR:HG23	1:B:1142:ARG:C	2.37	0.50
1:A:1032:LEU:HD22	1:A:1079:PHE:CD1	2.47	0.50
1:B:1045:ILE:O	1:B:1045:ILE:HG23	2.12	0.50
1:B:1073:LEU:O	1:B:1075:GLY:N	2.46	0.49
1:A:1128:HIS:NE2	1:A:1247:GLU:OE2	2.46	0.48
1:A:1044:ARG:HD2	1:A:1293:ASN:OD1	2.13	0.48
1:A:1161:VAL:HG21	1:B:1206:THR:CG2	2.44	0.48
1:B:1219:TRP:CE3	1:B:1221:VAL:HG23	2.49	0.47
1:B:1293:ASN:HD22	1:B:1294:ALA:N	2.11	0.47
1:A:1148:SER:HB3	1:A:1178:VAL:HG22	1.97	0.47
1:B:1203:ILE:HG12	1:B:1232:ASP:HB3	1.97	0.47
1:A:1068:ASP:OD1	1:A:1068:ASP:C	2.58	0.46
1:B:1173:LEU:HD11	1:B:1202:LEU:HD21	1.97	0.45
1:B:1284:GLY:H	1:B:1287:ASN:HD21	1.65	0.44
1:B:1029:MET:CE	1:B:1032:LEU:HD21	2.48	0.43
1:B:1068:ASP:O	1:B:1068:ASP:OD1	2.37	0.42
1:A:1024:ASN:ND2	1:A:1027:LYS:HB2	2.35	0.42
1:A:1199:TYR:CE1	1:A:1240:VAL:HG11	2.55	0.42
1:A:1266:ALA:CB	1:A:1274:PHE:CE2	3.03	0.42
1:B:1181:LYS:HA	1:B:1196:ASP:HA	2.02	0.42
1:B:1045:ILE:O	1:B:1045:ILE:CG2	2.67	0.41
1:B:1117:ALA:HB2	1:B:1157:TYR:CD1	2.55	0.41
1:A:1203:ILE:HG12	1:A:1232:ASP:HB3	2.02	0.41
1:B:1072:GLU:O	1:B:1073:LEU:HD23	2.19	0.41
1:B:1128:HIS:HB2	1:B:1146:THR:HG23	2.03	0.41
1:B:1216:GLY:N	1:B:1219:TRP:O	2.53	0.41
1:B:1219:TRP:HE3	1:B:1221:VAL:HG23	1.86	0.41
1:A:1169:PRO:HA	1:A:1208:VAL:HG22	2.03	0.41
1:B:1234:LEU:HD13	1:B:1248:LYS:HG3	2.03	0.41
1:A:1052:ALA:HB3	1:A:1056:PHE:HD2	1.85	0.40
1:A:1266:ALA:HB3	1:A:1274:PHE:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/285 (91%)	248 (96%)	11 (4%)	0	100	100
1	B	258/285 (90%)	248 (96%)	10 (4%)	0	100	100
All	All	517/570 (91%)	496 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/224 (85%)	171 (90%)	19 (10%)	7	11
1	B	192/224 (86%)	175 (91%)	17 (9%)	9	15
All	All	382/448 (85%)	346 (91%)	36 (9%)	8	13

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1035	ILE
1	A	1045	ILE
1	A	1059	ASN
1	A	1065	VAL
1	A	1102	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1130	ASN
1	A	1133	THR
1	A	1164	ASP
1	A	1172	GLU
1	A	1203	ILE
1	A	1213	SER
1	A	1215	SER
1	A	1232	ASP
1	A	1241	LEU
1	A	1242	ARG
1	A	1247	GLU
1	A	1250	ILE
1	A	1264	LEU
1	A	1299	SER
1	B	1032	LEU
1	B	1045	ILE
1	B	1065	VAL
1	B	1082[A]	PHE
1	B	1082[B]	PHE
1	B	1089	SER
1	B	1103	VAL
1	B	1133	THR
1	B	1141	THR
1	B	1142	ARG
1	B	1172	GLU
1	B	1182	GLN
1	B	1185	TRP
1	B	1232	ASP
1	B	1241	LEU
1	B	1293	ASN
1	B	1299	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1036	ASN
1	A	1230	GLN
1	A	1271	ASN
1	A	1290	ASN
1	A	1295	ASN
1	B	1230	GLN
1	B	1293	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1295	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	265/285 (92%)	0.29	8 (3%)	52 45	26, 62, 74, 80	2 (0%)
1	B	269/285 (94%)	0.77	21 (7%)	19 14	29, 63, 74, 83	1 (0%)
All	All	534/570 (93%)	0.53	29 (5%)	31 25	26, 63, 74, 83	3 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1039	ALA	3.9
1	B	1082[A]	PHE	3.5
1	B	1298	TYR	3.5
1	A	1235	ALA	3.0
1	B	1141	THR	2.8
1	B	1300	PHE	2.8
1	B	1301	GLY	2.7
1	A	1236	ASN	2.6
1	B	1189	GLY	2.6
1	B	1159	TYR	2.5
1	A	1295	ASN	2.5
1	B	1092	SER	2.5
1	B	1135	THR	2.4
1	A	1298	TYR	2.4
1	B	1237	GLY	2.4
1	B	1259	LEU	2.3
1	B	1297	ARG	2.3
1	B	1053	SER	2.3
1	A	1231	PHE	2.3
1	B	1057	SER	2.2
1	B	1091	ALA	2.2
1	B	1137	ALA	2.2
1	B	1275	GLY	2.1
1	B	1134	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1301	GLY	2.1
1	A	1300	PHE	2.1
1	B	1276	LEU	2.1
1	A	1079	PHE	2.1
1	B	1051	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.