



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:40 PM UTC

PDB ID : 7QOY / pdb_00007qoy
Title : A mutant of the nitrile hydratase from *Geobacillus pallidus* having enhanced thermostability
Authors : Van Wyk, J.C.; Cowan, D.A.; Danson, M.J.; Tsekoka, T.L.; Sayed, M.F.; Sewell, B.T.
Deposited on : 2021-12-29
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	64% 86% 6% • 6%
2	B	229	49% 88% 9% • •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrile hydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	S	0	4	0
			1663	1059	293	304	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	169	ARG	SER	engineered mutation	UNP Q84FS5

- Molecule 2 is a protein called Nitrile hydratase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	8	0
			1906	1230	318	349	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	43	LYS	MET	engineered mutation	UNP Q84FS6
B	150	ALA	THR	engineered mutation	UNP Q84FS6

- Molecule 3 is COBALT (III) ION (CCD ID: 3CO) (formula: Co) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Co	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Cl 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	189	Total 189	O 189	0	0
5	B	238	Total 238	O 238	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nitrile hydratase



• Molecule 2: Nitrile hydratase subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.18Å 106.18Å 82.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.16 – 1.45 75.08 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.5 (75.16-1.45) 99.5 (75.08-1.45)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.97 (at 1.45Å)	Xtriage
Refinement program	PHENIX 1.10.1-2155	Depositor
R, R_{free}	0.142 , 0.184 0.290 , 0.309	Depositor DCC
R_{free} test set	4175 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3998	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, 3CO, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	2/1701 (0.1%)	0.84	0/2310
2	B	1.17	5/1974 (0.3%)	0.94	3/2664 (0.1%)
All	All	1.13	7/3675 (0.2%)	0.90	3/4974 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	163	ARG	CD-NE	-9.49	1.32	1.46
2	B	163	ARG	CZ-NH2	7.23	1.42	1.33
2	B	163	ARG	CG-CD	6.77	1.72	1.52
2	B	25	ILE	CA-CB	6.71	1.62	1.54
2	B	188	ARG	CD-NE	5.72	1.54	1.46
1	A	148	ARG	CZ-NH2	-5.28	1.26	1.33
1	A	61	MET	CG-SD	5.12	1.93	1.80

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	210	LYS	N-CA-C	5.30	119.72	112.88
2	B	163	ARG	NH1-CZ-NH2	5.20	126.06	119.30
2	B	163	ARG	NE-CZ-NH1	-5.14	116.36	121.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1663	0	1657	11	0
2	B	1906	0	1880	30	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	189	0	0	2	0
5	B	238	0	0	6	1
All	All	3998	0	3537	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:158:HIS:H	2:B:187:HIS:HE1	1.08	0.93
2:B:2:ASN:HD21	2:B:166:ARG:HH12	1.25	0.85
2:B:158:HIS:H	2:B:187:HIS:CE1	1.99	0.78
2:B:162:PRO:HG2	2:B:217[A]:ILE:HD11	1.78	0.66
2:B:100:GLU:HG3	2:B:104:LYS:HE2	1.78	0.65
2:B:93:LYS:NZ	2:B:97:GLU:HG2	2.16	0.61
2:B:163:ARG:HD2	5:B:337:HOH:O	2.01	0.60
1:A:41:HIS:HE1	5:B:412:HOH:O	1.83	0.60
2:B:211[B]:GLN:HG3	2:B:213:ASP:OD1	2.02	0.60
2:B:22:GLU:HG2	5:B:453:HOH:O	2.03	0.56
1:A:97[B]:GLN:HG3	1:A:124:TRP:CZ3	2.42	0.55
2:B:162:PRO:HG2	2:B:217[A]:ILE:CD1	2.38	0.54
2:B:60:GLU:OE1	2:B:158:HIS:HD2	1.91	0.54
1:A:16:HIS:H	1:A:16:HIS:CD2	2.24	0.53
2:B:158:HIS:HE1	5:B:352:HOH:O	1.91	0.53
2:B:93:LYS:HE2	2:B:93:LYS:HA	1.89	0.53
2:B:158:HIS:N	2:B:187:HIS:HE1	1.92	0.53
2:B:121:LEU:C	2:B:121:LEU:HD23	2.33	0.53
2:B:5:HIS:HD2	5:B:407:HOH:O	1.92	0.52
2:B:105:LYS:HE3	2:B:107:ASP:HB2	1.92	0.52
1:A:195:LYS:N	1:A:195:LYS:HD2	2.26	0.50
2:B:93:LYS:C	2:B:93:LYS:HD3	2.37	0.49
2:B:93:LYS:HD3	2:B:93:LYS:O	2.12	0.49
2:B:49[A]:MET:HE1	2:B:80:VAL:HG22	1.95	0.49
2:B:162:PRO:CG	2:B:217[A]:ILE:HD11	2.41	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:93:LYS:HZ3	2:B:97:GLU:HG2	1.79	0.48
2:B:105:LYS:C	2:B:105:LYS:HD2	2.39	0.48
1:A:17:HIS:HD2	5:A:562:HOH:O	1.95	0.48
1:A:52[A]:LYS:HB2	1:A:52[A]:LYS:NZ	2.29	0.48
2:B:2:ASN:ND2	2:B:166:ARG:HH12	2.04	0.47
1:A:14:PHE:HB3	1:A:17:HIS:CD2	2.51	0.45
1:A:199:ARG:NE	2:B:21[B]:GLU:HG2	2.32	0.45
1:A:97[B]:GLN:NE2	5:A:405:HOH:O	2.51	0.44
1:A:121:CSD:HB2	1:A:174:ARG:CZ	2.47	0.44
2:B:93:LYS:HZ2	2:B:97:GLU:HG2	1.81	0.43
2:B:93:LYS:O	2:B:97:GLU:HG3	2.19	0.42
2:B:163:ARG:CD	5:B:337:HOH:O	2.64	0.41
2:B:93:LYS:HD3	2:B:97:GLU:HG3	2.03	0.40
1:A:188:MET:HE2	1:A:192:GLU:HB3	2.03	0.40
2:B:5:HIS:CG	2:B:60:GLU:HG2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:462:HOH:O	5:B:495:HOH:O[8_777]	2.02	0.18

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/216 (94%)	199 (98%)	2 (1%)	1 (0%)	24	7
2	B	232/229 (101%)	228 (98%)	4 (2%)	0	100	100
All	All	434/445 (98%)	427 (98%)	6 (1%)	1 (0%)	43	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/190 (95%)	177 (98%)	4 (2%)	45	13
2	B	201/195 (103%)	197 (98%)	4 (2%)	48	16
All	All	382/385 (99%)	374 (98%)	8 (2%)	55	14

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52[A]	LYS
1	A	52[B]	LYS
1	A	61	MET
1	A	78	GLN
2	B	49[A]	MET
2	B	49[B]	MET
2	B	93	LYS
2	B	105	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	41	HIS
1	A	56	HIS
1	A	62	ASN
1	A	78	GLN
1	A	149	GLN
1	A	180	GLN
2	B	2	ASN
2	B	5	HIS
2	B	29	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	152	ASN
2	B	158	HIS
2	B	187	HIS
2	B	192	ASN
2	B	194	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSD	A	119	1	4,7,8	1.00	0	1,8,10	0.23	0
1	CSD	A	121	3,1	4,7,8	2.21	1 (25%)	1,8,10	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	119	1	-	0/2/6/8	-
1	CSD	A	121	3,1	-	0/2/6/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	121	CSD	OD1-SG	3.85	1.51	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	121	CSD	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/216 (92%)	2.82	138 (69%) 0 0	9, 17, 34, 42	4 (2%)
2	B	227/229 (99%)	2.15	112 (49%) 0 0	5, 15, 30, 38	8 (3%)
All	All	427/445 (95%)	2.46	250 (58%) 0 0	5, 16, 32, 42	12 (2%)

All (250) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	117	ALA	7.6
1	A	71	TRP	6.6
1	A	189	THR	6.5
1	A	184	GLY	6.4
1	A	185	THR	6.4
2	B	118	LEU	6.3
1	A	14	PHE	6.2
1	A	75	ALA	6.1
1	A	68	ALA	6.0
1	A	70	ALA	6.0
2	B	101	VAL	5.8
2	B	93	LYS	5.7
1	A	192	GLU	5.7
1	A	197	VAL	5.7
1	A	193	LEU	5.6
1	A	86	THR	5.5
2	B	111	PRO	5.4
1	A	72	THR	5.3
1	A	36	LEU	5.2
2	B	95	LEU	5.2
2	B	130	LEU	5.2
1	A	194	ALA	5.2
1	A	66	VAL	5.1
1	A	37	ILE	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	12	PRO	4.9
1	A	188	MET	4.9
1	A	208	ILE	4.9
2	B	103	LEU	4.8
1	A	24	PHE	4.8
2	B	107	ASP	4.8
2	B	119	VAL	4.8
2	B	116	PRO	4.7
2	B	85	VAL	4.7
1	A	182	PRO	4.7
1	A	64	ALA	4.6
2	B	104	LYS	4.6
1	A	76	PHE	4.6
1	A	190	GLU	4.6
1	A	67	VAL	4.6
2	B	110	ILE	4.6
1	A	74	PRO	4.5
1	A	34[A]	SER	4.5
1	A	87	VAL	4.5
2	B	114	GLU	4.5
2	B	126	LEU	4.4
2	B	42	CYS	4.4
1	A	110	VAL	4.4
2	B	122	VAL	4.4
1	A	81	LEU	4.3
1	A	196	LEU	4.3
1	A	10	ILE	4.3
1	A	206	ALA	4.3
2	B	125	ALA	4.3
1	A	210	PRO	4.2
1	A	42	LEU	4.2
2	B	186	ALA	4.2
1	A	211	PRO	4.2
1	A	91	LEU	4.1
2	B	90	LEU	4.0
1	A	159	PRO	4.0
1	A	89	ARG	4.0
2	B	99	THR	4.0
1	A	187	GLY	4.0
1	A	50	VAL	4.0
1	A	162	VAL	4.0
2	B	81	ALA	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	155	GLY	3.9
1	A	79[A]	ARG	3.9
2	B	113	ARG	3.9
2	B	120	LYS	3.9
2	B	102	PHE	3.9
1	A	109	THR	3.9
1	A	90	GLU	3.9
1	A	191	GLU	3.9
1	A	161	SER	3.8
2	B	46	GLY	3.8
1	A	183	GLU	3.8
1	A	93	TYR	3.7
2	B	82	TYR	3.7
1	A	65	LYS	3.7
1	A	78	GLN	3.7
1	A	92	GLY	3.7
2	B	41	GLY	3.7
1	A	201	SER	3.6
2	B	89	VAL	3.6
1	A	179	PRO	3.6
1	A	80	LEU	3.6
2	B	209	TYR	3.6
1	A	13	ARG	3.6
1	A	186	GLU	3.6
1	A	178	LEU	3.6
2	B	88	GLY	3.6
1	A	11	ASP	3.6
2	B	91	ASP	3.5
2	B	121	LEU	3.5
2	B	22	GLU	3.5
2	B	207	TRP	3.5
2	B	44	ALA	3.5
2	B	87	THR	3.5
2	B	19	VAL	3.5
1	A	40	GLY	3.4
2	B	210	LYS	3.4
1	A	198	THR	3.4
1	A	31	ALA	3.4
2	B	48	GLY	3.4
2	B	20	LYS	3.4
1	A	35	LEU	3.4
1	A	209	GLU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	61	MET	3.4
2	B	43	LYS	3.4
1	A	15	PRO	3.3
1	A	32	LEU	3.3
1	A	158	LEU	3.3
2	B	47[A]	LEU	3.3
2	B	84	LEU	3.3
2	B	185	ALA	3.3
2	B	86	ASP	3.3
2	B	115	ASP	3.3
1	A	47	ILE	3.3
1	A	44	SER	3.2
2	B	80	VAL	3.2
1	A	103	VAL	3.2
2	B	227	VAL	3.2
1	A	126	LEU	3.2
1	A	45	ASP	3.2
2	B	94	GLU	3.2
1	A	63	GLY	3.2
2	B	129	GLY	3.2
1	A	107	THR	3.1
2	B	156	THR	3.1
1	A	88	LEU	3.1
1	A	62	ASN	3.1
1	A	181	ARG	3.1
2	B	208	GLY	3.1
1	A	46	ALA	3.1
1	A	180	GLN	3.1
1	A	150	VAL	3.1
2	B	109	LYS	3.1
2	B	25	ILE	3.1
2	B	127	TYR	3.0
1	A	17	HIS	3.0
1	A	106	ASN	3.0
1	A	104	VAL	3.0
1	A	41	HIS	3.0
1	A	154	PHE	3.0
1	A	38	GLU	3.0
2	B	100	GLU	2.9
2	B	98	ARG	2.9
1	A	111	HIS	2.9
1	A	94	TYR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	211[A]	GLN	2.9
2	B	128	ASP	2.9
2	B	108	THR	2.9
2	B	97	GLU	2.9
1	A	108	ASP	2.9
1	A	164	ILE	2.9
1	A	203	ILE	2.9
2	B	18	TYR	2.9
2	B	61	LEU	2.9
1	A	95	GLY	2.9
1	A	156	LEU	2.8
1	A	205	VAL	2.8
2	B	83	ASN	2.8
1	A	207	LYS	2.8
1	A	58	LEU	2.8
1	A	56	HIS	2.8
1	A	43	SER	2.8
2	B	153	ILE	2.8
1	A	113	VAL	2.7
1	A	149	GLN	2.7
2	B	9	GLY	2.7
2	B	36	PHE	2.7
1	A	53	HIS	2.7
2	B	187	HIS	2.7
1	A	125	PRO	2.7
1	A	127	LEU	2.7
2	B	38	LEU	2.7
2	B	40	ALA	2.6
1	A	73	ASP	2.6
2	B	112	ARG	2.6
2	B	52	PHE	2.6
2	B	175	VAL	2.6
2	B	105	LYS	2.6
1	A	175	PHE	2.6
2	B	204	GLU	2.6
1	A	18	HIS	2.6
1	A	160	ASP	2.6
2	B	50	LYS	2.5
2	B	190	GLY	2.5
1	A	177	VAL	2.5
1	A	152	LYS	2.5
2	B	138	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	60	PRO	2.5
2	B	79	THR	2.5
1	A	204	GLY	2.5
1	A	105	GLU	2.4
2	B	76	TRP	2.4
1	A	39	LYS	2.4
2	B	163	ARG	2.4
1	A	147	PRO	2.4
1	A	96	LEU	2.4
2	B	123	GLU	2.4
2	B	133	LEU	2.4
2	B	13	PHE	2.4
2	B	137	SER	2.4
2	B	206	LEU	2.4
1	A	112	ASN	2.4
2	B	92	GLU	2.4
2	B	55	PHE	2.3
2	B	152	ASN	2.3
1	A	51	ILE	2.3
2	B	169	TYR	2.3
2	B	49[A]	MET	2.3
1	A	84	SER	2.3
2	B	27	PHE	2.3
2	B	34	LEU	2.3
2	B	51	ALA	2.2
1	A	49	ARG	2.2
1	A	98	GLY	2.2
1	A	52[A]	LYS	2.2
1	A	151	LEU	2.2
2	B	106	PRO	2.2
1	A	195	LYS	2.2
1	A	148	ARG	2.2
2	B	131	SER	2.2
2	B	77	ILE	2.1
2	B	216	TYR	2.1
2	B	188	ARG	2.1
1	A	25	TRP	2.1
1	A	133	TRP	2.1
2	B	45	GLN	2.1
1	A	77	LYS	2.1
1	A	166	VAL	2.1
1	A	27	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	78	ALA	2.1
2	B	23	GLU	2.1
1	A	54	TYR	2.1
1	A	157	ASP	2.1
2	B	124	LYS	2.1
2	B	180	VAL	2.1
2	B	165	ALA	2.1
2	B	189	LYS	2.1
1	A	128	GLY	2.1
1	A	123	PRO	2.1
2	B	203	ALA	2.0
1	A	57	GLU	2.0
1	A	83	ASP	2.0
2	B	21[A]	GLU	2.0
2	B	39	VAL	2.0
2	B	65	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	121	8/9	0.95	0.11	12,12,19,26	0
1	CSD	A	119	8/9	0.98	0.07	9,11,12,13	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CL	A	302	1/1	0.73	0.23	17,17,17,17	0
3	3CO	A	301	1/1	0.99	0.03	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 3CO A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.