



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 29, 2026 – 04:24 pm BST

PDB ID : 9QOQ / pdb_00009qoq
Title : Leishmania donovani ISP2 in complex with bovine trypsin
Authors : Freitag-Pohl, S.; Pohl, E.
Deposited on : 2025-03-26
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.2
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.2
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	246	0% 80% 9% 9%
1	D	246	6% 78% 11% 9%
1	F	246	8% 79% 11% 9%
1	H	246	2% 80% 9% 9%
2	A	158	3% 67% 18% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	158	3% 63% 20% • 15%
2	E	158	4% 68% 16% • 15%
2	G	158	5% 64% 18% • 16%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 20989 atoms, of which 10252 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cationic Trypsin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	223	Total	C	H	N	O	S	76	0	0
			3151	997	1550	274	316	14			
1	D	223	Total	C	H	N	O	S	105	0	0
			3013	963	1470	270	296	14			
1	F	223	Total	C	H	N	O	S	102	0	0
			3039	969	1486	269	301	14			
1	H	223	Total	C	H	N	O	S	84	0	0
			3128	992	1541	273	308	14			

- Molecule 2 is a protein called Ecotin family protein.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	135	Total	C	H	N	O	S	44	1	0
			2121	686	1056	180	194	5			
2	A	139	Total	C	H	N	O	S	51	3	0
			2174	702	1078	190	199	5			
2	E	135	Total	C	H	N	O	S	52	0	0
			2040	668	1001	173	193	5			
2	G	132	Total	C	H	N	O	S	58	0	0
			1998	651	988	168	186	5			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	3	0
			14	3	8	3		
3	D	1	Total	C	H	O	3	0
			14	3	8	3		
3	H	1	Total	C	H	O	3	0
			14	3	8	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	2	0
			17	4	10	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	H	O	2	0
			17	4	10	3		
4	H	1	Total	C	H	O	2	0
			17	4	10	3		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		
5	D	1	Total	Ca	0	0
			1	1		

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	H	O	2	0
			24	6	14	4		
6	F	1	Total	C	H	O	2	0
			24	6	14	4		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	43	Total	O	0	0
			43	43		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	22	Total 22	O 22	0	0
7	C	25	Total 25	O 25	0	0
7	A	17	Total 17	O 17	0	0
7	E	15	Total 15	O 15	0	0
7	G	10	Total 10	O 10	0	0
7	F	19	Total 19	O 19	0	0
7	H	31	Total 31	O 31	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cationic Trypsin




- Molecule 1: Cationic Trypsin




- Molecule 1: Cationic Trypsin



- Molecule 1: Cationic Trypsin





• Molecule 2: Ecotin family protein



• Molecule 2: Ecotin family protein



• Molecule 2: Ecotin family protein



• Molecule 2: Ecotin family protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	170.70Å 191.79Å 45.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.94 – 2.59 59.94 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.2 (59.94-2.59) 99.2 (59.94-2.59)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.82)	Depositor
R, R_{free}	0.198 , 0.274 0.199 , 0.275	Depositor DCC
R_{free} test set	750 reflections (1.58%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20989	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PGE, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.75	0/1632	1.15	9/2217 (0.4%)
1	D	0.69	0/1573	1.13	5/2145 (0.2%)
1	F	0.65	0/1583	1.14	9/2154 (0.4%)
1	H	0.67	0/1618	1.10	6/2201 (0.3%)
2	A	0.69	0/1132	1.21	3/1543 (0.2%)
2	C	0.69	0/1095	1.21	5/1489 (0.3%)
2	E	0.66	0/1065	1.18	5/1452 (0.3%)
2	G	0.66	0/1035	1.17	2/1411 (0.1%)
All	All	0.68	0/10733	1.16	44/14612 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	7
2	C	0	3
2	E	0	3
2	G	0	1
All	All	0	14

There are no bond length outliers.

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	76	ASP	CA-CB-CG	7.42	120.02	112.60
2	A	65	ARG	NE-CZ-NH2	6.90	125.41	119.20
1	H	231	LYS	N-CA-CB	6.70	119.96	110.24
1	B	197	GLN	N-CA-CB	6.70	119.94	109.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	231	LYS	CB-CA-C	-6.63	99.84	111.05

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	29[A]	ARG	Sidechain
2	A	29[B]	ARG	Sidechain
2	C	138	ARG	Sidechain
2	C	29	ARG	Sidechain
2	C	54	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1601	1550	1525	10	1
1	D	1543	1470	1406	12	0
1	F	1553	1486	1428	12	0
1	H	1587	1541	1506	12	0
2	A	1096	1078	1054	19	1
2	C	1065	1056	1039	19	0
2	E	1039	1001	973	15	0
2	G	1010	988	956	20	0
3	B	6	8	8	0	0
3	D	6	8	8	0	0
3	H	6	8	8	0	0
4	B	7	10	10	1	0
4	E	7	10	10	0	0
4	H	7	10	10	2	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
6	C	10	14	14	0	0
6	F	10	14	14	2	0
7	A	17	0	0	0	0
7	B	43	0	0	0	0
7	C	25	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	D	22	0	0	0	0
7	E	15	0	0	1	0
7	F	19	0	0	0	0
7	G	10	0	0	1	0
7	H	31	0	0	0	0
All	All	10737	10252	9969	105	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:90:MET:CE	2:C:109:VAL:HG21	2.04	0.88
2:E:90:MET:CE	2:E:109:VAL:HG21	2.06	0.86
2:A:90:MET:CE	2:A:109:VAL:HG21	2.06	0.85
2:G:74:THR:HG22	2:G:81:THR:HG22	1.56	0.85
2:C:65:ARG:NH1	2:C:91:SER:OG	2.14	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:131:SER:HG	2:A:105:ARG:HG3[1_554]	1.22	0.38

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	221/246 (90%)	213 (96%)	7 (3%)	1 (0%)	24	44
1	D	221/246 (90%)	210 (95%)	11 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	221/246 (90%)	213 (96%)	8 (4%)	0	100	100
1	H	221/246 (90%)	214 (97%)	7 (3%)	0	100	100
2	A	140/158 (89%)	138 (99%)	2 (1%)	0	100	100
2	C	134/158 (85%)	131 (98%)	3 (2%)	0	100	100
2	E	131/158 (83%)	130 (99%)	1 (1%)	0	100	100
2	G	128/158 (81%)	125 (98%)	3 (2%)	0	100	100
All	All	1417/1616 (88%)	1374 (97%)	42 (3%)	1 (0%)	48	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	81	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	173/202 (86%)	167 (96%)	6 (4%)	32	57
1	D	151/202 (75%)	140 (93%)	11 (7%)	13	27
1	F	154/202 (76%)	146 (95%)	8 (5%)	21	42
1	H	168/202 (83%)	160 (95%)	8 (5%)	23	45
2	A	110/132 (83%)	106 (96%)	4 (4%)	31	56
2	C	109/132 (83%)	103 (94%)	6 (6%)	19	39
2	E	102/132 (77%)	102 (100%)	0	100	100
2	G	99/132 (75%)	95 (96%)	4 (4%)	28	52
All	All	1066/1336 (80%)	1019 (96%)	47 (4%)	25	48

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	G	70	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	172	LYS
2	G	112	SER
1	F	52	LEU
1	F	206	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	42	ASN
2	E	15	ASN
1	H	42	ASN
1	H	84	ASN
1	H	102	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PGE	C	201	-	9,9,9	0.35	0	8,8,8	0.30	0
3	GOL	H	301	-	5,5,5	0.12	0	5,5,5	0.27	0
3	GOL	D	301	-	5,5,5	0.21	0	5,5,5	0.40	0
4	PEG	H	302	-	6,6,6	0.32	0	5,5,5	0.32	0
3	GOL	B	301	-	5,5,5	0.25	0	5,5,5	0.34	0
4	PEG	B	302	-	6,6,6	0.34	0	5,5,5	0.22	0
4	PEG	E	201	-	6,6,6	0.41	0	5,5,5	0.40	0
6	PGE	F	301	-	9,9,9	0.47	0	8,8,8	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	C	201	-	-	5/7/7/7	-
3	GOL	H	301	-	-	2/4/4/4	-
3	GOL	D	301	-	-	4/4/4/4	-
4	PEG	H	302	-	-	3/4/4/4	-
3	GOL	B	301	-	-	1/4/4/4	-
4	PEG	B	302	-	-	2/4/4/4	-
4	PEG	E	201	-	-	2/4/4/4	-
6	PGE	F	301	-	-	3/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	301	GOL	O1-C1-C2-C3
3	D	301	GOL	C1-C2-C3-O3
3	H	301	GOL	O1-C1-C2-O2
3	H	301	GOL	O1-C1-C2-C3
4	B	302	PEG	O2-C3-C4-O4

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	302	PEG	2	0
4	B	302	PEG	1	0
6	F	301	PGE	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	223/246 (90%)	-0.02	2 (0%) 81 79	41, 56, 80, 94	0
1	D	223/246 (90%)	0.63	15 (6%) 24 20	44, 74, 96, 117	0
1	F	223/246 (90%)	0.75	20 (8%) 15 12	51, 75, 102, 125	0
1	H	223/246 (90%)	0.32	5 (2%) 62 58	47, 67, 89, 104	0
2	A	139/158 (87%)	0.28	5 (3%) 46 41	29, 64, 89, 110	3 (2%)
2	C	135/158 (85%)	0.17	4 (2%) 52 48	38, 62, 90, 108	1 (0%)
2	E	135/158 (85%)	0.48	6 (4%) 39 34	47, 69, 95, 123	0
2	G	132/158 (83%)	0.67	8 (6%) 27 22	49, 73, 102, 124	0
All	All	1433/1616 (88%)	0.41	65 (4%) 38 34	29, 68, 95, 125	4 (0%)

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	113	THR	5.6
2	A	12	THR	5.4
2	G	113	THR	5.1
2	G	99	GLY	5.0
2	E	115	ASN	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	D	302	1/1	0.79	0.15	126,126,126,126	0
4	PEG	E	201	7/7	0.86	0.16	85,91,104,109	2
5	CA	B	303	1/1	0.89	0.10	77,77,77,77	1
6	PGE	C	201	10/10	0.89	0.16	80,108,115,121	2
3	GOL	D	301	6/6	0.90	0.13	84,97,102,107	3
6	PGE	F	301	10/10	0.90	0.13	78,91,106,112	2
3	GOL	H	301	6/6	0.92	0.14	79,84,88,91	3
4	PEG	H	302	7/7	0.92	0.11	76,90,94,94	2
4	PEG	B	302	7/7	0.95	0.14	73,86,89,90	2
3	GOL	B	301	6/6	0.96	0.10	69,81,90,91	3

6.5 Other polymers [i](#)

There are no such residues in this entry.