



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:16 PM UTC

PDB ID : 2QPM / pdb_00002qpm
Title : Leu492Ala mutant of Maize cytokinin oxidase/dehydrogenase complexed with benzylurea inhibitor CPBU
Authors : Briozzo, P.
Deposited on : 2007-07-24
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	19% 76% 20% .
2	B	2	100%
2	C	2	100%
2	D	2	50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	D	2	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4302 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytokinin dehydrogenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	0	0
			3781	2399	665	706	11			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ALA	GLY	variant	UNP Q9T0N8
A	168	THR	ASN	variant	UNP Q9T0N8
A	254	LEU	PHE	variant	UNP Q9T0N8
A	492	ALA	LEU	engineered mutation	UNP Q9T0N8

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			


- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
5	A	1	18	13	1	3	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	338	Total	O	0	0
			338	338		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytokinin dehydrogenase 1



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	248.50Å 49.70Å 50.50Å 90.00° 93.70° 90.00°	Depositor
Resolution (Å)	25.00 – 1.85 25.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.6 (25.00-1.85) 87.7 (25.00-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.79Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.221 , 0.246 0.237 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.732	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4302	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 246, FAD, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3870	0.89	7/5273 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	454	ALA	N-CA-C	-6.69	98.35	109.46
1	A	168	THR	N-CA-C	-6.60	99.42	109.85
1	A	250	ARG	N-CA-C	-6.58	99.18	109.25
1	A	257	THR	N-CA-C	-6.10	105.64	114.12
1	A	227	GLY	N-CA-C	-5.80	107.24	115.30
1	A	305	GLY	N-CA-C	-5.28	108.34	115.36
1	A	240	ILE	N-CA-C	5.08	115.63	108.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3781	0	3696	78	0
2	B	28	0	25	3	0
2	C	28	0	25	0	0
2	D	28	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	28	0	26	0	0
4	A	53	0	29	0	0
5	A	18	0	12	3	0
6	A	338	0	0	5	0
All	All	4302	0	3838	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:441:GLY:HA2	1:A:512:ARG:HH12	1.36	0.90
1:A:141:TYR:CE1	1:A:239:ARG:HD2	2.19	0.78
5:A:1536:246:H4	5:A:1536:246:O1	1.83	0.76
1:A:398:LEU:HG	1:A:400:MET:HE3	1.74	0.68
1:A:141:TYR:CD1	1:A:239:ARG:HD2	2.30	0.66
1:A:136:SER:HB3	1:A:141:TYR:CE1	2.31	0.65
1:A:400:MET:HE1	1:A:478:LEU:HD21	1.79	0.65
1:A:143:ASP:OD1	1:A:239:ARG:HD3	1.97	0.63
1:A:90:SER:O	1:A:92:PRO:HD3	2.00	0.62
1:A:382:GLU:HG3	1:A:391:TRP:CE2	2.37	0.60
1:A:143:ASP:HB3	2:B:1:NAG:H81	1.84	0.59
1:A:524:ARG:HG3	1:A:533:PHE:HB3	1.84	0.59
1:A:252:VAL:HG12	1:A:366:GLN:HG2	1.85	0.58
1:A:48:ARG:HD3	1:A:74:TYR:CE2	2.38	0.57
1:A:47:LEU:HD23	1:A:73:LEU:HD13	1.85	0.57
1:A:398:LEU:CG	1:A:400:MET:HE3	2.35	0.57
1:A:441:GLY:CA	1:A:512:ARG:HH12	2.15	0.55
1:A:342:ALA:O	1:A:346:VAL:HG23	2.06	0.55
1:A:136:SER:HB3	1:A:141:TYR:CD1	2.43	0.53
1:A:463:ALA:O	1:A:464:PRO:O	2.26	0.53
1:A:65:THR:HG23	1:A:112:PHE:HE2	1.74	0.53
1:A:134:ASN:ND2	2:B:1:NAG:H82	2.25	0.52
1:A:190:ARG:O	1:A:437:MET:HE2	2.09	0.52
1:A:515:GLU:HG2	1:A:516:MET:HE2	1.91	0.52
1:A:254:LEU:HD22	1:A:254:LEU:N	2.25	0.51
1:A:412:ARG:HD2	6:A:1706:HOH:O	2.10	0.51
2:D:1:NAG:O3	2:D:1:NAG:H83	2.10	0.50
1:A:398:LEU:HD11	1:A:400:MET:CE	2.42	0.49
1:A:427:PRO:HG2	1:A:458:LEU:HD12	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:400:MET:HE1	1:A:478:LEU:CD2	2.41	0.49
1:A:509:LYS:HE3	6:A:1797:HOH:O	2.12	0.49
1:A:199:LEU:N	1:A:199:LEU:HD12	2.27	0.49
1:A:57:ALA:HB1	1:A:118:VAL:HG11	1.95	0.49
1:A:512:ARG:NE	6:A:1933:HOH:O	2.46	0.48
1:A:65:THR:HG22	6:A:1657:HOH:O	2.13	0.48
1:A:130:PRO:HB3	2:B:2:NAG:O7	2.13	0.47
1:A:338:ASN:O	1:A:339:ALA:C	2.57	0.47
1:A:382:GLU:HG3	1:A:391:TRP:CD2	2.48	0.47
1:A:463:ALA:HB3	1:A:466:ASP:HB2	1.96	0.47
5:A:1536:246:O1	5:A:1536:246:C4	2.49	0.47
1:A:58:SER:HB3	1:A:68:LEU:HD23	1.96	0.47
1:A:398:LEU:CD1	1:A:400:MET:HE3	2.45	0.47
1:A:127:ALA:C	1:A:129:ALA:H	2.22	0.46
1:A:460:SER:HB3	1:A:462:VAL:HG13	1.97	0.46
1:A:400:MET:CE	1:A:478:LEU:HD21	2.44	0.45
1:A:472:GLU:HA	1:A:475:ARG:NH1	2.32	0.45
1:A:462:VAL:HG23	1:A:463:ALA:N	2.32	0.45
1:A:88:ALA:HA	1:A:94:TRP:CD2	2.52	0.44
1:A:424:ILE:HD13	1:A:428:LEU:HD11	2.00	0.44
1:A:425:VAL:HG21	1:A:462:VAL:HG21	1.98	0.44
1:A:209:GLU:HG2	1:A:211:VAL:HG23	1.99	0.44
1:A:411:ASP:OD2	1:A:416:LYS:NZ	2.50	0.44
1:A:463:ALA:O	1:A:464:PRO:C	2.61	0.44
1:A:123:SER:HA	1:A:126:ASP:OD2	2.18	0.44
1:A:284:MET:HE2	1:A:334:LEU:HD21	2.00	0.44
1:A:391:TRP:O	1:A:395:HIS:HE1	2.01	0.43
1:A:164:PRO:HD2	1:A:167:TRP:CH2	2.53	0.43
1:A:165:ARG:HG2	1:A:243:GLU:HB3	1.99	0.43
1:A:387:LYS:HD2	1:A:387:LYS:HA	1.86	0.43
1:A:73:LEU:HD22	1:A:84:LEU:HD22	2.00	0.43
1:A:350:LEU:O	1:A:354:LEU:HG	2.19	0.43
1:A:425:VAL:HG11	1:A:462:VAL:HG11	2.01	0.42
1:A:127:ALA:C	1:A:129:ALA:N	2.76	0.42
1:A:141:TYR:CD1	1:A:141:TYR:C	2.98	0.42
1:A:271:ALA:HA	1:A:272:PRO:HD3	1.86	0.42
1:A:100:PHE:CZ	1:A:234:VAL:HB	2.54	0.42
1:A:67:ALA:HB1	1:A:114:PRO:HG3	2.02	0.42
1:A:170:TYR:OH	1:A:382:GLU:OE1	2.30	0.42
5:A:1536:246:H14	6:A:1877:HOH:O	2.20	0.41
1:A:48:ARG:HB3	1:A:50:ASP:OD1	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:441:GLY:HA2	1:A:512:ARG:NH1	2.18	0.41
1:A:400:MET:HE2	1:A:400:MET:HB3	1.89	0.41
1:A:190:ARG:C	1:A:437:MET:HE2	2.46	0.41
1:A:286:TYR:HB3	1:A:333:THR:OG1	2.21	0.41
1:A:170:TYR:CZ	1:A:172:TYR:HB2	2.56	0.41
1:A:286:TYR:CZ	1:A:288:GLU:HG3	2.56	0.41
1:A:70:ALA:N	1:A:116:GLY:O	2.55	0.40
1:A:186:GLY:O	1:A:286:TYR:HA	2.21	0.40
1:A:288:GLU:HA	1:A:431:TYR:HB3	2.04	0.40
1:A:253:ARG:HA	1:A:330:ILE:O	2.21	0.40
1:A:133:ILE:HA	1:A:143:ASP:O	2.21	0.40
1:A:209:GLU:HG2	1:A:211:VAL:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	492/516 (95%)	473 (96%)	17 (4%)	2 (0%)	30 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	PRO
1	A	182	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	385/394 (98%)	380 (99%)	5 (1%)	61 51

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LEU
1	A	95	PRO
1	A	220	ASP
1	A	388	LEU
1	A	464	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	218	ASN
1	A	379	HIS
1	A	504	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.55	0	17,19,21	0.79	1 (5%)
2	NAG	B	2	2	14,14,15	0.48	0	17,19,21	0.76	1 (5%)
2	NAG	C	1	2,1	14,14,15	0.46	0	17,19,21	0.70	1 (5%)
2	NAG	C	2	2	14,14,15	0.49	0	17,19,21	0.76	1 (5%)
2	NAG	D	1	2,1	14,14,15	0.68	0	17,19,21	0.89	1 (5%)
2	NAG	D	2	2	14,14,15	0.61	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	4/6/23/26	0/1/1/1
2	NAG	B	2	2	-	5/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	1/1/5/7	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	C2-N2-C7	-2.38	119.71	122.90
2	C	2	NAG	C2-N2-C7	-2.27	119.86	122.90
2	B	1	NAG	C2-N2-C7	-2.23	119.91	122.90
2	B	2	NAG	C2-N2-C7	-2.23	119.92	122.90
2	C	1	NAG	C2-N2-C7	-2.03	120.18	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	D	2	NAG	C1

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
2	B	2	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	B	2	NAG	C1-C2-N2-C7


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0
2	B	1	NAG	2	0
2	B	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1601	1	14,14,15	0.47	0	17,19,21	0.75	1 (5%)
3	NAG	A	1608	1	14,14,15	0.49	0	17,19,21	0.73	1 (5%)
4	FAD	A	1535	1	58,58,58	1.57	11 (18%)	85,89,89	1.77	16 (18%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	246	A	1536	-	19,19,19	0.98	2 (10%)	23,24,24	1.91	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1601	1	-	1/6/23/26	0/1/1/1
3	NAG	A	1608	1	-	4/6/23/26	0/1/1/1
4	FAD	A	1535	1	-	3/34/50/50	0/6/6/6
5	246	A	1536	-	-	0/9/9/9	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1535	FAD	C4X-N5	5.16	1.41	1.30
4	A	1535	FAD	C9-C8	4.61	1.45	1.39
4	A	1535	FAD	C8M-C8	3.97	1.58	1.51
4	A	1535	FAD	C4A-N3A	2.92	1.39	1.34
4	A	1535	FAD	C9A-C5X	2.85	1.45	1.41
4	A	1535	FAD	C9A-N10	2.80	1.46	1.41
5	A	1536	246	C1-N3	-2.61	1.31	1.37
4	A	1535	FAD	C10-N1	2.57	1.38	1.33
5	A	1536	246	C3-N3	-2.35	1.36	1.41
4	A	1535	FAD	P-O3P	2.10	1.61	1.59
4	A	1535	FAD	C6-C5X	2.05	1.43	1.40
4	A	1535	FAD	C5A-C4A	-2.05	1.35	1.39
4	A	1535	FAD	C9-C9A	2.03	1.42	1.39

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1535	FAD	N3A-C2A-N1A	-6.22	119.16	128.58
5	A	1536	246	C4-C5-N6	-6.17	120.22	125.59
4	A	1535	FAD	C4A-C5A-N7A	4.45	115.67	110.58
4	A	1535	FAD	C5A-C6A-N6A	-4.38	112.44	123.29
5	A	1536	246	C7-N6-C5	4.29	121.76	115.58
4	A	1535	FAD	C6A-C5A-C4A	-4.23	111.40	117.18
4	A	1535	FAD	C5'-C4'-C3'	4.02	119.81	112.22
4	A	1535	FAD	O2A-PA-O5B	3.64	124.06	107.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1535	FAD	C5A-N7A-C8A	-3.26	98.32	103.45
4	A	1535	FAD	N6A-C6A-N1A	3.15	125.39	118.38
4	A	1535	FAD	O3P-PA-O1A	-3.10	101.37	110.70
4	A	1535	FAD	O2P-P-O3P	2.93	115.20	107.27
4	A	1535	FAD	C1B-N9A-C8A	2.90	133.54	127.09
5	A	1536	246	C8-C7-N6	-2.73	120.62	123.97
4	A	1535	FAD	N9A-C8A-N7A	2.62	117.66	113.94
5	A	1536	246	C3-N3-C1	-2.61	121.28	126.61
4	A	1535	FAD	C4X-C10-N10	2.57	120.16	116.48
3	A	1601	NAG	C2-N2-C7	-2.50	119.55	122.90
4	A	1535	FAD	O2B-C2B-C3B	2.46	119.69	111.82
4	A	1535	FAD	C2A-N3A-C4A	2.31	117.47	111.83
3	A	1608	NAG	C2-N2-C7	-2.23	119.92	122.90
4	A	1535	FAD	O4B-C1B-N9A	-2.07	104.11	108.09

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1608	NAG	C8-C7-N2-C2
3	A	1608	NAG	O7-C7-N2-C2
3	A	1608	NAG	O5-C5-C6-O6
3	A	1608	NAG	C4-C5-C6-O6
4	A	1535	FAD	C2'-C3'-C4'-C5'
4	A	1535	FAD	PA-O3P-P-O1P
3	A	1601	NAG	C1-C2-N2-C7
4	A	1535	FAD	PA-O3P-P-O2P


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1536	246	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/516 (96%)	1.23	96 (19%) 3 2	15, 28, 49, 62	13 (2%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	ALA	7.1
1	A	465	ASN	6.9
1	A	464	PRO	5.7
1	A	47	LEU	4.9
1	A	425	VAL	4.9
1	A	128	ALA	4.6
1	A	51	SER	4.1
1	A	211	VAL	4.0
1	A	92	PRO	3.9
1	A	130	PRO	3.9
1	A	462	VAL	3.8
1	A	498	ARG	3.8
1	A	463	ALA	3.7
1	A	384	ALA	3.7
1	A	39	ALA	3.6
1	A	129	ALA	3.6
1	A	124	LEU	3.5
1	A	210	MET	3.5
1	A	512	ARG	3.5
1	A	497	ASP	3.5
1	A	388	LEU	3.5
1	A	280	SER	3.4
1	A	138	ASP	3.4
1	A	34	TRP	3.4
1	A	86	SER	3.3
1	A	282	GLY	3.3
1	A	220	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	136	SER	3.3
1	A	216	GLN	3.2
1	A	141	TYR	3.2
1	A	40	ALA	3.2
1	A	156	ALA	3.0
1	A	219	ALA	3.0
1	A	281	PHE	3.0
1	A	90	SER	2.9
1	A	273	ARG	2.9
1	A	41	LEU	2.9
1	A	91	THR	2.9
1	A	116	GLY	2.9
1	A	96	TYR	2.9
1	A	85	LEU	2.9
1	A	45	GLY	2.8
1	A	87	ALA	2.8
1	A	519	LYS	2.8
1	A	49	THR	2.8
1	A	389	GLY	2.7
1	A	337	ASP	2.7
1	A	339	ALA	2.7
1	A	154	LEU	2.6
1	A	378	VAL	2.6
1	A	35	PRO	2.6
1	A	65	THR	2.6
1	A	466	ASP	2.5
1	A	506	GLY	2.5
1	A	493	ALA	2.5
1	A	272	PRO	2.5
1	A	217	LEU	2.5
1	A	392	ARG	2.5
1	A	94	TRP	2.5
1	A	383	VAL	2.5
1	A	213	CYS	2.4
1	A	52	ASN	2.4
1	A	367	ARG	2.4
1	A	54	THR	2.4
1	A	82	VAL	2.4
1	A	387	LYS	2.4
1	A	499	SER	2.4
1	A	385	LEU	2.3
1	A	36	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	67	ALA	2.3
1	A	345	ALA	2.3
1	A	131	PRO	2.3
1	A	214	SER	2.3
1	A	137	ALA	2.2
1	A	212	THR	2.2
1	A	246	PRO	2.2
1	A	296	SER	2.2
1	A	336	TYR	2.2
1	A	518	ASN	2.2
1	A	514	VAL	2.2
1	A	43	LEU	2.2
1	A	84	LEU	2.2
1	A	33	PRO	2.2
1	A	158	LEU	2.1
1	A	44	ASP	2.1
1	A	69	PRO	2.1
1	A	476	ARG	2.1
1	A	172	TYR	2.0
1	A	125	GLY	2.0
1	A	334	LEU	2.0
1	A	501	TRP	2.0
1	A	48	ARG	2.0
1	A	140	ARG	2.0
1	A	93	GLY	2.0
1	A	134	ASN	2.0
1	A	393	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	D	1	14/15	0.52	0.20	35,38,41,45	14

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	2	14/15	0.64	0.20	46,48,48,49	14
2	NAG	C	2	14/15	0.65	0.21	41,42,43,45	14
2	NAG	B	1	14/15	0.65	0.17	39,41,42,44	14
2	NAG	D	2	14/15	0.65	0.22	49,50,55,55	14
2	NAG	C	1	14/15	0.76	0.15	31,34,39,40	14

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	1608	14/15	0.61	0.24	55,56,57,57	14
3	NAG	A	1601	14/15	0.77	0.16	27,35,40,41	14
5	246	A	1536	18/18	0.78	0.19	15,33,44,44	0
4	FAD	A	1535	53/53	0.93	0.09	15,21,25,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.