



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 02:04 PM EDT

PDB ID : 3QPR / pdb_00003qpr
Title : HK97 Prohead I encapsidating inactive virally encoded protease
Authors : Huang, R.K.; Khayat, R.; Lee, K.K.; Gertsman, I.; Duda, R.L.; Hendrix, R.W.; Johnson, J.E.
Deposited on : 2011-02-14
Resolution : 5.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 5.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1066 (6.38-4.00)
Ramachandran outliers	187476	1004 (6.46-3.94)
RSRZ outliers	180081	1006 (6.38-4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	48% 16% • 34%
1	B	385	2% 42% 19% • 35%
1	C	385	44% 17% • 36%
1	D	385	50% 13% • 34%
1	E	385	42% 21% • 34%
1	F	385	39% 21% 5% 36%
1	G	385	42% 17% 5% 36%

2 Entry composition


There is only 1 type of molecule in this entry. The entry contains 8654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Major capsid protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	0	0	0
			1260	748	256	256			
1	B	250	Total	C	N	O	0	0	0
			1231	731	250	250			
1	C	248	Total	C	N	O	0	0	0
			1221	725	248	248			
1	D	254	Total	C	N	O	0	0	0
			1250	742	254	254			
1	E	255	Total	C	N	O	0	0	0
			1255	745	255	255			
1	F	248	Total	C	N	O	0	0	0
			1221	725	248	248			
1	G	247	Total	C	N	O	0	0	0
			1216	722	247	247			


- Molecule 1: Major capsid protein



- Molecule 1: Major capsid protein



- Molecule 1: Major capsid protein



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	560.12Å 560.12Å 560.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 5.20 45.00 – 5.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (45.00-5.20) 77.3 (45.00-5.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 5.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.461 , 0.467 0.390 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	175.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 417.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.32$, $\langle L^2 \rangle = 0.16$	Xtriage
Estimated twinning fraction	0.125 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.60	EDS
Total number of atoms	8654	wwPDB-VP
Average B, all atoms (Å ²)	149.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	1/1259 (0.1%)	1.43	34/1749 (1.9%)
1	B	0.67	1/1230 (0.1%)	1.40	21/1709 (1.2%)
1	C	0.54	1/1220 (0.1%)	1.27	16/1695 (0.9%)
1	D	0.54	0/1249	1.37	18/1735 (1.0%)
1	E	0.51	0/1254	1.40	20/1742 (1.1%)
1	F	0.72	1/1220 (0.1%)	1.54	37/1695 (2.2%)
1	G	0.62	0/1215	1.71	44/1688 (2.6%)
All	All	0.61	4/8647 (0.0%)	1.45	190/12013 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	212	MET	C-N	-9.41	1.22	1.33
1	C	288	LEU	C-N	7.19	1.42	1.33
1	B	190	VAL	C-N	-6.66	1.24	1.33
1	F	134	ILE	CA-CB	-5.10	1.47	1.54

The worst 5 of 190 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	125	ILE	N-CA-C	19.33	129.05	110.30
1	G	297	PHE	N-CA-C	-19.00	86.10	112.13
1	G	207	ILE	N-CA-C	15.30	125.11	110.42
1	A	298	GLY	N-CA-C	14.98	135.37	115.59
1	D	250	THR	N-CA-C	14.43	127.05	111.03

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	362	CYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1260	0	595	46	0
1	B	1231	0	583	84	0
1	C	1221	0	580	54	0
1	D	1250	0	591	44	0
1	E	1255	0	593	76	0
1	F	1221	0	580	67	0
1	G	1216	0	578	55	0
All	All	8654	0	4100	389	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 389 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:ASN:CB	1:B:172:SER:HA	1.21	1.57
1:B:158:ASN:CB	1:B:172:SER:CA	2.08	1.31
1:E:202:MET:HA	1:F:142:ARG:O	1.26	1.26
1:B:193:SER:CB	1:C:149:GLU:CB	2.14	1.26
1:A:293:GLY:O	1:C:298:GLY:CA	1.84	1.24

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/385 (66%)	194 (76%)	54 (21%)	6 (2%)	4	26
1	B	248/385 (64%)	194 (78%)	47 (19%)	7 (3%)	4	24
1	C	246/385 (64%)	200 (81%)	36 (15%)	10 (4%)	2	17
1	D	252/385 (66%)	205 (81%)	39 (16%)	8 (3%)	3	21
1	E	253/385 (66%)	215 (85%)	33 (13%)	5 (2%)	6	30
1	F	246/385 (64%)	201 (82%)	36 (15%)	9 (4%)	2	19
1	G	245/385 (64%)	201 (82%)	38 (16%)	6 (2%)	4	26
All	All	1744/2695 (65%)	1410 (81%)	283 (16%)	51 (3%)	3	23

5 of 51 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	ILE
1	A	170	PRO
1	A	208	ASN
1	B	170	PRO
1	B	350	ARG

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/385 (66%)	-0.24	4 (1%) 70 57	118, 145, 169, 195	0
1	B	250/385 (64%)	-0.29	6 (2%) 59 49	118, 149, 175, 187	0
1	C	248/385 (64%)	-0.27	1 (0%) 88 77	120, 150, 174, 189	0
1	D	254/385 (65%)	-0.26	2 (0%) 82 69	118, 149, 170, 188	0
1	E	255/385 (66%)	-0.31	4 (1%) 70 57	120, 152, 173, 203	0
1	F	248/385 (64%)	-0.30	2 (0%) 82 69	123, 147, 170, 191	0
1	G	247/385 (64%)	-0.25	0 100 100	122, 149, 172, 193	0
All	All	1758/2695 (65%)	-0.27	19 (1%) 78 63	118, 149, 173, 203	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	198	ASP	4.0
1	E	121	ILE	3.9
1	B	241	THR	3.8
1	C	298	GLY	3.7
1	A	241	THR	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.