



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:22 PM UTC

PDB ID : 5QPQ / pdb_00005qpq
Title : PanDDA analysis group deposition – Crystal Structure of T. cruzi FPPS in complex with FMOPL000631a
Authors : Petrick, J.K.; Nelson, E.R.; Muenzker, L.; Krojer, T.; Douangamath, A.; Brandao-Neto, J.; von Delft, F.; Dekker, C.; Jahnke, W.
Deposited on : 2019-03-12
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	403	-	-	X	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Farnesyl diphosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	360	2883	1853	471	536	23	0	1	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q8WS26
A	0	PRO	-	expression tag	UNP Q8WS26

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is {N}-[2-(aminocarbamoyl)phenyl]ethanamide (CCD ID: LV7) (formula: $C_9H_{11}N_3O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	9	3	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	333	Total	O	0	0
			333	333		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Farnesyl diphosphate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	57.91Å 57.91Å 396.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.02 – 1.49 66.02 – 1.49	Depositor EDS
% Data completeness (in resolution range)	98.1 (66.02-1.49) 98.1 (66.02-1.49)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 1.49Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.200 , 0.235 0.208 , 0.244	Depositor DCC
R_{free} test set	3433 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3246	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACT, LV7, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.79	41/2948 (1.4%)	1.51	18/3995 (0.5%)

All (41) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	101	MET	SD-CE	-15.55	1.40	1.79
1	A	242	ILE	N-CA	-9.32	1.35	1.46
1	A	338	GLN	CA-C	7.99	1.63	1.52
1	A	105	VAL	CA-CB	7.87	1.63	1.55
1	A	339	LEU	N-CA	7.57	1.56	1.46
1	A	222	LEU	C-O	7.55	1.32	1.24
1	A	230	VAL	CA-C	-7.17	1.44	1.52
1	A	31	ASN	CA-C	7.16	1.62	1.52
1	A	34	ARG	N-CA	-6.88	1.38	1.46
1	A	102	ASP	CA-C	-6.43	1.43	1.52
1	A	110	LYS	CA-C	6.31	1.58	1.52
1	A	226	ALA	CA-CB	-6.29	1.44	1.53
1	A	340	LYS	N-CA	6.27	1.54	1.46
1	A	23	GLN	N-CA	6.25	1.54	1.46
1	A	34	ARG	CA-C	6.13	1.60	1.52
1	A	77	LEU	C-O	6.12	1.31	1.24
1	A	180	LEU	CA-C	-6.11	1.44	1.52
1	A	113	TRP	N-CA	-6.05	1.39	1.46
1	A	28	ILE	CA-C	-6.05	1.45	1.52
1	A	146	ARG	CD-NE	-5.97	1.37	1.46
1	A	17	PHE	N-CA	-5.93	1.39	1.46
1	A	218	VAL	CA-CB	-5.86	1.46	1.54
1	A	323	ALA	N-CA	5.51	1.53	1.46
1	A	337	GLU	CA-C	-5.48	1.45	1.52
1	A	33	ALA	N-CA	5.40	1.52	1.46
1	A	165	VAL	C-O	5.40	1.30	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	ILE	C-O	-5.26	1.18	1.24
1	A	204	VAL	C-O	5.26	1.30	1.24
1	A	166	GLY	C-O	5.21	1.30	1.23
1	A	139	ALA	CA-C	5.20	1.59	1.52
1	A	204	VAL	N-CA	-5.20	1.40	1.46
1	A	169	TYR	N-CA	5.14	1.52	1.46
1	A	144	ALA	C-N	-5.11	1.27	1.33
1	A	165	VAL	N-CA	5.09	1.52	1.46
1	A	38	ILE	CA-C	5.08	1.59	1.52
1	A	102	ASP	C-O	5.07	1.30	1.24
1	A	36	LEU	N-CA	5.07	1.52	1.46
1	A	74	GLU	C-O	5.07	1.30	1.24
1	A	77	LEU	N-CA	5.07	1.52	1.46
1	A	31	ASN	N-CA	-5.06	1.40	1.46
1	A	160	ASP	C-O	5.03	1.29	1.24

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	MET	CG-SD-CE	-7.78	83.78	100.90
1	A	101	MET	CG-SD-CE	-7.50	84.40	100.90
1	A	144	ALA	CA-C-N	-7.43	111.01	122.60
1	A	144	ALA	C-N-CA	-7.43	111.01	122.60
1	A	340	LYS	N-CA-C	7.35	120.16	111.71
1	A	138	MET	CG-SD-CE	-6.95	85.60	100.90
1	A	125	ILE	N-CA-CB	6.26	117.46	110.51
1	A	338	GLN	N-CA-C	-6.05	104.69	111.28
1	A	203	ILE	CB-CA-C	-5.97	104.24	111.88
1	A	204	VAL	N-CA-C	5.83	116.61	110.72
1	A	91	GLN	N-CA-C	-5.82	104.85	111.07
1	A	208	THR	CB-CA-C	-5.52	100.52	110.23
1	A	220	GLY	O-C-N	5.43	127.41	122.19
1	A	2	ALA	N-CA-CB	5.32	117.86	110.04
1	A	51	ARG	NE-CZ-NH2	5.27	123.94	119.20
1	A	347	ALA	O-C-N	-5.09	116.72	122.12
1	A	84	GLY	N-CA-C	-5.02	106.70	112.73
1	A	213	TYR	N-CA-C	5.01	119.13	112.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2883	0	2849	20	1
2	A	10	0	0	1	0
3	A	4	0	3	2	0
4	A	2	0	0	0	0
5	A	14	0	0	1	0
6	A	333	0	0	10	4
All	All	3246	0	2852	24	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:GLU:HG2	6:A:743:HOH:O	1.46	1.14
2:A:402:SO4:O3	6:A:501:HOH:O	2.05	0.74
1:A:355:GLU:O	1:A:359:LYS:HG3	1.91	0.69
1:A:201:LYS:HG2	1:A:244:GLU:OE2	1.98	0.63
1:A:318:GLN:O	6:A:502:HOH:O	2.16	0.63
3:A:403:ACT:H2	6:A:523:HOH:O	2.01	0.61
1:A:3:SER:OG	1:A:60:GLU:OE2	2.16	0.60
1:A:208:THR:HG23	6:A:724:HOH:O	2.05	0.56
1:A:209:THR:HG22	1:A:243:GLY:HA3	1.88	0.55
1:A:250:ASP:HB2	6:A:536:HOH:O	2.06	0.54
5:A:406:LV7:N	5:A:406:LV7:O1	2.42	0.52
1:A:94[B]:TYR:CE2	1:A:163:THR:HG21	2.47	0.49
1:A:94[B]:TYR:CZ	1:A:163:THR:HG21	2.47	0.49
1:A:3:SER:HB3	1:A:53:MET:HE1	1.95	0.48
1:A:356:LYS:NZ	6:A:510:HOH:O	2.43	0.47
3:A:403:ACT:CH3	6:A:523:HOH:O	2.59	0.46
1:A:340:LYS:O	1:A:341:VAL:C	2.61	0.44
1:A:353:VAL:HG22	6:A:731:HOH:O	2.18	0.42
1:A:272:ALA:HA	1:A:297:TYR:CE2	2.55	0.42
1:A:204:VAL:HG22	1:A:247:GLN:HG2	2.02	0.41
1:A:250:ASP:CB	6:A:536:HOH:O	2.67	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:TYR:HB2	1:A:85:TRP:CZ2	2.55	0.41
1:A:105:VAL:HG12	1:A:106:MET:HG2	2.03	0.41
1:A:101:MET:HB2	1:A:101:MET:HE3	1.73	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:807:HOH:O	6:A:807:HOH:O[10_554]	1.78	0.42
1:A:137:ILE:CD1	6:A:736:HOH:O[10_444]	2.01	0.19
6:A:689:HOH:O	6:A:703:HOH:O[8_545]	2.09	0.11
6:A:769:HOH:O	6:A:811:HOH:O[8_445]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/364 (99%)	354 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	308/310 (99%)	305 (99%)	3 (1%)	68 45

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	260	GLU
1	A	300	LYS
1	A	359	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	316	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	A	403	-	3,3,3	0.66	0	3,3,3	0.70	0
2	SO4	A	402	-	4,4,4	0.47	0	6,6,6	0.15	0
5	LV7	A	406	-	14,14,14	2.26	7 (50%)	18,18,18	2.65	6 (33%)
2	SO4	A	401	-	4,4,4	1.63	1 (25%)	6,6,6	1.83	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	LV7	A	406	-	-	2/10/10/10	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	406	LV7	C1-N	4.02	1.43	1.36
5	A	406	LV7	C8-N1	3.78	1.37	1.33
5	A	406	LV7	C7-C8	-3.75	1.42	1.50
5	A	406	LV7	N2-N1	-2.66	1.37	1.41
5	A	406	LV7	O1-C8	2.46	1.29	1.23
5	A	406	LV7	C5-C6	-2.43	1.34	1.38
2	A	401	SO4	O2-S	2.41	1.59	1.44
5	A	406	LV7	C2-N	2.33	1.46	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	406	LV7	C-C1-N	5.92	123.84	114.95
5	A	406	LV7	C8-N1-N2	-5.73	112.53	121.50
5	A	406	LV7	O-C1-N	-4.03	117.54	123.06
2	A	401	SO4	O4-S-O1	-3.70	90.21	109.56
5	A	406	LV7	C6-C7-C2	3.17	122.38	118.81
5	A	406	LV7	O1-C8-N1	-3.14	118.69	122.46
5	A	406	LV7	C2-C7-C8	-2.10	118.26	120.96

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	406	LV7	C-C1-N-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	406	LV7	O-C1-N-C2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	ACT	2	0
2	A	402	SO4	1	0
5	A	406	LV7	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/364 (98%)	0.68	25 (6%) 23 25	3, 24, 42, 76	8 (2%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	94[A]	TYR	16.5
1	A	133	SER	8.3
1	A	150	LYS	7.7
1	A	157	GLN	7.6
1	A	98	ASP	7.0
1	A	341	VAL	3.5
1	A	359	LYS	3.5
1	A	1	MET	3.3
1	A	260	GLU	3.1
1	A	329	VAL	3.1
1	A	310	ARG	3.0
1	A	344	PRO	2.9
1	A	265	VAL	2.9
1	A	261	GLN	2.8
1	A	308	VAL	2.7
1	A	263	GLY	2.6
1	A	332	VAL	2.6
1	A	338	GLN	2.5
1	A	345	THR	2.5
1	A	336	ILE	2.4
1	A	2	ALA	2.3
1	A	360	ARG	2.2
1	A	145	ASP	2.1
1	A	240	HIS	2.1
1	A	258	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ACT	A	403	4/4	0.41	0.40	32,35,36,36	4
2	SO4	A	402	5/5	0.74	0.31	37,39,42,43	5
5	LV7	A	406	14/14	0.78	0.17	38,43,60,60	0
2	SO4	A	401	5/5	0.84	0.13	27,31,34,34	0
4	ZN	A	404	1/1	0.98	0.05	24,24,24,24	1
4	ZN	A	405	1/1	0.99	0.06	29,29,29,29	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LV7 A 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.