



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:57 PM UTC

PDB ID : 7QPJ / pdb_00007qpj
Title : Crystal structure of engineered TCR (756) complexed to HLA-A*02:01 presenting MAGE-A10 9-mer peptide
Authors : Simister, P.C.; Border, E.C.; Vieira, J.F.; Pumphrey, N.J.
Deposited on : 2022-01-04
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	277	7% 90% 10%
2	D	100	2% 89% 10%
3	E	9	89% 11%
4	A	206	10% 89% 8%
5	B	241	12% 89% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	C	303	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 13302 atoms, of which 6204 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	C	276	Total	C	H	N	O	S	91	8	0
			4370	1414	2115	406	426	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	MET	-	initiating methionine	UNP Q861F7

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	100	Total	C	H	N	O	S	37	1	0
			1575	517	765	136	153	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Melanoma-associated antigen 10.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	E	9	Total	C	H	N	O	S	3	0	0
			138	45	66	11	15	1			

- Molecule 4 is a protein called T-cell receptor alpha chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	A	200	Total	C	H	N	O	S	57	3	0
			2990	955	1449	261	317	8			

- Molecule 5 is a protein called T-cell receptor beta chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
5	B	238	Total	C	H	N	O	S	64	3	0
			3645	1189	1777	315	355	9			

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	H	O	2	0
			14	3	8	3		
6	C	1	Total	C	H	O	2	0
			14	3	8	3		
6	C	1	Total	C	H	O	2	0
			14	3	8	3		
6	C	1	Total	C	H	O	2	0
			14	3	8	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	192	Total	O	0	0
			192	192		
7	D	77	Total	O	0	0
			77	77		
7	E	4	Total	O	0	0
			4	4		
7	A	136	Total	O	0	0
			136	136		
7	B	119	Total	O	0	0
			119	119		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Melanoma-associated antigen 10




- Molecule 4: T-cell receptor alpha chain



- Molecule 5: T-cell receptor beta chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.20Å 77.45Å 116.68Å 90.00° 102.70° 90.00°	Depositor
Resolution (Å)	64.11 – 1.54 64.11 – 1.54	Depositor EDS
% Data completeness (in resolution range)	96.7 (64.11-1.54) 89.1 (64.11-1.54)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.54Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC 5.8.0267	Depositor
R, R_{free}	0.202 , 0.236 0.203 , 0.239	Depositor DCC
R_{free} test set	6620 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.423	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 32.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13302	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	1.11	2/2350 (0.1%)	1.27	2/3195 (0.1%)
2	D	1.00	0/836	1.17	2/1137 (0.2%)
3	E	1.46	0/73	1.25	0/95
4	A	1.13	1/1578 (0.1%)	1.27	0/2141
5	B	1.05	0/1929	1.27	0/2633
All	All	1.09	3/6766 (0.0%)	1.26	4/9201 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	68	VAL	C-O	6.02	1.31	1.24
4	A	36	TRP	C-O	5.92	1.31	1.24
1	C	46	MET	C-O	5.29	1.30	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	153	VAL	CA-C-O	-5.25	115.28	120.85
2	D	97	ASP	CA-C-N	5.03	127.72	120.38
2	D	97	ASP	C-N-CA	5.03	127.72	120.38
1	C	142	GLN	N-CA-C	-5.01	106.00	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2255	2115	2083	24	0
2	D	810	765	744	5	0
3	E	72	66	65	1	0
4	A	1541	1449	1435	12	0
5	B	1868	1777	1755	15	0
6	C	24	32	32	6	0
7	A	136	0	0	6	0
7	B	119	0	0	3	0
7	C	192	0	0	4	0
7	D	77	0	0	1	0
7	E	4	0	0	0	0
All	All	7098	6204	6114	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:98:ARG:HE	1:C:115:HIS:HE1	1.21	0.87
2:D:4:ARG:NH1	7:D:201:HOH:O	2.10	0.84
1:C:66:ARG:HH22	6:C:303:GOL:H2	1.43	0.83
4:A:196:GLU:OE1	7:A:301:HOH:O	1.97	0.82
1:C:231:LEU:O	1:C:231:LEU:HD12	1.88	0.72
5:B:201:PRO:O	7:B:501:HOH:O	2.13	0.66
5:B:206:ARG:NH2	5:B:208:GLN:OE1	2.29	0.66
1:C:229:THR:HG21	7:C:502:HOH:O	1.95	0.66
1:C:55:GLN:NE2	7:C:402:HOH:O	2.29	0.65
1:C:264:HIS:CD2	1:C:266:GLY:H	2.16	0.63
4:A:54:ASN:ND2	7:A:304:HOH:O	2.22	0.62
1:C:36:ARG:NH2	7:C:403:HOH:O	2.33	0.60
4:A:111:GLN:OE1	7:A:303:HOH:O	2.16	0.60
4:A:151:ASP:OD1	4:A:152:SER:N	2.34	0.60
4:A:4:GLU:HG3	7:A:365:HOH:O	2.03	0.58
1:C:98:ARG:HE	1:C:115:HIS:CE1	2.13	0.56
1:C:232:VAL:HG13	1:C:245:TRP:CZ2	2.41	0.56
5:B:31:ASN:HB3	7:B:541:HOH:O	2.05	0.55
5:B:124:VAL:HG23	5:B:234:ALA:HB3	1.90	0.53
5:B:14:LEU:HD11	5:B:20:MET:HG2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:193:ILE:HG23	7:A:341:HOH:O	2.09	0.52
1:C:168:TRP:CZ2	6:C:301:GOL:H31	2.45	0.51
1:C:66:ARG:HH12	6:C:303:GOL:H31	1.75	0.50
4:A:8:ASN:N	7:A:302:HOH:O	2.10	0.50
5:B:16:ILE:HD12	5:B:112:GLU:HG2	1.92	0.50
1:C:71[A]:HIS:HE1	1:C:100:TYR:OH	1.96	0.48
1:C:66:ARG:HH12	6:C:303:GOL:C3	2.27	0.48
1:C:231:LEU:HD12	1:C:231:LEU:C	2.39	0.48
2:D:25:ASN:HB3	2:D:66:LEU:HD11	1.95	0.47
2:D:16:ALA:HB3	2:D:98:ARG:HG3	1.97	0.46
3:E:6:MET:HG2	4:A:98:ARG:CD	2.45	0.46
1:C:264:HIS:HD2	1:C:266:GLY:H	1.61	0.46
5:B:199:GLN:HA	5:B:239:ARG:O	2.17	0.45
1:C:66:ARG:NH2	6:C:303:GOL:H2	2.21	0.45
1:C:76[B]:ARG:HG3	7:C:464:HOH:O	2.16	0.45
1:C:118:ALA:HB2	2:D:61:TRP:CE2	2.51	0.45
4:A:104:SER:HA	5:B:43:GLY:HA3	1.98	0.45
5:B:107[B]:ARG:HG2	5:B:151:HIS:CD2	2.52	0.44
4:A:157:THR:HG21	5:B:188:SER:OG	2.17	0.44
1:C:128:LYS:HD2	1:C:133:SER:OG	2.18	0.43
1:C:188:THR:HA	1:C:205:TRP:O	2.18	0.43
1:C:225:GLN:O	1:C:229:THR:HG23	2.19	0.43
5:B:208:GLN:HG3	5:B:231:ILE:HG23	2.01	0.43
4:A:155:TYR:O	4:A:176:ALA:HA	2.20	0.42
5:B:201:PRO:HD3	7:B:597:HOH:O	2.19	0.42
5:B:204:HIS:NE2	5:B:235:GLU:OE1	2.47	0.41
5:B:172:GLN:HE21	5:B:172:GLN:HB3	1.69	0.41
1:C:211:PRO:O	1:C:264:HIS:HE1	2.03	0.41
1:C:168:TRP:CE2	6:C:301:GOL:H31	2.56	0.41
1:C:232:VAL:CG1	1:C:245:TRP:CZ2	3.03	0.41
4:A:111:GLN:HB3	4:A:142:SER:HB3	2.02	0.41
2:D:24:LEU:O	2:D:68:TYR:HA	2.21	0.40
5:B:206:ARG:HG3	5:B:235:GLU:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	282/277 (102%)	278 (99%)	4 (1%)	0	100	100
2	D	99/100 (99%)	98 (99%)	1 (1%)	0	100	100
3	E	7/9 (78%)	7 (100%)	0	0	100	100
4	A	201/206 (98%)	195 (97%)	6 (3%)	0	100	100
5	B	239/241 (99%)	230 (96%)	8 (3%)	1 (0%)	30	12
All	All	828/833 (99%)	808 (98%)	19 (2%)	1 (0%)	48	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	B	97	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	230/233 (99%)	227 (99%)	3 (1%)	61	31
2	D	87/95 (92%)	84 (97%)	3 (3%)	32	7
3	E	7/7 (100%)	7 (100%)	0	100	100
4	A	172/182 (94%)	169 (98%)	3 (2%)	53	23
5	B	197/207 (95%)	193 (98%)	4 (2%)	48	18
All	All	693/724 (96%)	680 (98%)	13 (2%)	55	20

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	20[A]	GLU
1	C	20[B]	GLU
1	C	226	THR
2	D	2	ILE
2	D	90	GLN
2	D	98	ARG
4	A	181	SER
4	A	182	ASP
4	A	193	ILE
5	B	10	LYS
5	B	66[A]	ASN
5	B	66[B]	ASN
5	B	128	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	33	GLN
1	C	44	GLN
1	C	115	HIS
1	C	116	GLN
1	C	142	GLN
1	C	264	HIS
4	A	65	GLN
4	A	111	GLN
4	A	123	GLN
4	A	191	ASN
5	B	23	GLN
5	B	31	ASN
5	B	172	GLN
5	B	177	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	C	302	-	5,5,5	0.11	0	5,5,5	0.30	0
6	GOL	C	304	-	5,5,5	0.10	0	5,5,5	0.46	0
6	GOL	C	303	-	5,5,5	0.18	0	5,5,5	0.62	0
6	GOL	C	301	-	5,5,5	0.19	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	C	302	-	-	2/4/4/4	-
6	GOL	C	304	-	-	2/4/4/4	-
6	GOL	C	303	-	-	1/4/4/4	-
6	GOL	C	301	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	301	GOL	O1-C1-C2-C3
6	C	301	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	C	301	GOL	O2-C2-C3-O3
6	C	302	GOL	C1-C2-C3-O3
6	C	303	GOL	C1-C2-C3-O3
6	C	304	GOL	C1-C2-C3-O3
6	C	301	GOL	O1-C1-C2-O2
6	C	302	GOL	O2-C2-C3-O3
6	C	304	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	303	GOL	4	0
6	C	301	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	276/277 (99%)	0.32	20 (7%) 21 27	7, 22, 52, 73	9 (3%)
2	D	100/100 (100%)	0.37	2 (2%) 65 74	12, 26, 43, 53	1 (1%)
3	E	9/9 (100%)	-0.37	0 100 100	11, 13, 15, 16	0
4	A	200/206 (97%)	0.56	20 (10%) 12 17	9, 24, 55, 101	3 (1%)
5	B	238/241 (98%)	0.68	28 (11%) 9 12	11, 27, 50, 84	3 (1%)
All	All	823/833 (98%)	0.48	70 (8%) 16 21	7, 24, 51, 101	16 (1%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	B	3	ALA	5.5
1	C	277	PRO	4.9
5	B	240	ALA	4.7
4	A	129	SER	4.7
1	C	231	LEU	4.4
1	C	195	VAL	4.2
4	A	193	ILE	4.1
4	A	130	SER	4.0
4	A	131	ASP	3.6
4	A	128	LYS	3.6
4	A	149	SER	3.5
5	B	223	ASP	3.4
4	A	150	LYS	3.3
5	B	197	PHE	3.2
5	B	61	VAL	3.2
5	B	217	ASN	3.1
4	A	186	ALA	3.1
1	C	226	THR	3.0
5	B	64	GLY	3.0
4	A	179	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	275	TRP	3.0
1	C	196	SER	2.9
4	A	202	SER	2.9
1	C	268	PRO	2.9
5	B	237	TRP	2.8
1	C	273	LEU	2.8
5	B	239	ARG	2.8
4	A	183	PHE	2.8
1	C	267	LEU	2.7
5	B	158	VAL	2.7
1	C	222	GLY	2.6
5	B	157	TRP	2.6
5	B	55	ILE	2.6
5	B	4	GLY	2.6
5	B	238	GLY	2.6
1	C	52	TRP	2.5
2	D	2	ILE	2.5
4	A	151	ASP	2.5
5	B	79	LEU	2.4
5	B	221	THR	2.4
1	C	249	VAL	2.4
1	C	253	GLY	2.4
4	A	132	LYS	2.4
1	C	194	ALA	2.4
5	B	81	LEU	2.4
5	B	202	ARG	2.4
4	A	3	LYS	2.3
5	B	63	ASN	2.3
4	A	10	GLY	2.3
5	B	203	ASN	2.3
4	A	133	SER	2.3
1	C	138	ASP	2.3
1	C	19	GLY	2.3
5	B	225	ALA	2.2
2	D	44	GLY	2.2
4	A	143	GLN	2.2
4	A	185	CYS	2.1
5	B	70	SER	2.1
5	B	215	SER	2.1
5	B	163	VAL	2.1
4	A	184	ALA	2.1
1	C	198	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
5	B	164	HIS	2.1
1	C	2	GLY	2.1
1	C	223	GLU	2.1
1	C	137	ALA	2.1
5	B	201	PRO	2.1
5	B	113	ASP	2.0
5	B	216	GLU	2.0
4	A	187	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	C	303	6/6	0.82	0.14	31,38,51,51	2
6	GOL	C	302	6/6	0.87	0.14	35,42,48,48	2
6	GOL	C	301	6/6	0.91	0.12	32,39,42,43	2
6	GOL	C	304	6/6	0.91	0.11	31,33,34,35	2

6.5 Other polymers [i](#)

There are no such residues in this entry.