



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:44 AM UTC

PDB ID : 7QPT / pdb_00007qpt
Title : Botulinum neurotoxin A4 cell binding domain in complex with GD1a oligosaccharide
Authors : Gregory, K.S.; Acharya, K.R.; Liu, S.M.; Mojanaga, O.O.
Deposited on : 2022-01-05
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	3% 89% 8% .
1	B	433	12% 81% 8% . 10%
2	C	5	100%
2	D	5	100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neurotoxin type A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	420	Total	C	N	O	S	0	0	0
			3458	2211	587	646	14			
1	B	390	Total	C	N	O	S	0	1	0
			3238	2082	543	599	14			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	870	MET	-	initiating methionine	UNP Q3LRX8
A	871	HIS	-	expression tag	UNP Q3LRX8
A	872	HIS	-	expression tag	UNP Q3LRX8
A	873	HIS	-	expression tag	UNP Q3LRX8
A	874	HIS	-	expression tag	UNP Q3LRX8
A	875	HIS	-	expression tag	UNP Q3LRX8
A	876	HIS	-	expression tag	UNP Q3LRX8
B	870	MET	-	initiating methionine	UNP Q3LRX8
B	871	HIS	-	expression tag	UNP Q3LRX8
B	872	HIS	-	expression tag	UNP Q3LRX8
B	873	HIS	-	expression tag	UNP Q3LRX8
B	874	HIS	-	expression tag	UNP Q3LRX8
B	875	HIS	-	expression tag	UNP Q3LRX8
B	876	HIS	-	expression tag	UNP Q3LRX8

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	5	Total 68	C 37	N 2	O 29	0	0	0
2	D	5	Total 68	C 37	N 2	O 29	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	163	Total 163	O 163	0	0
3	B	74	Total 74	O 74	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neurotoxin type A



- Molecule 1: Neurotoxin type A



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



BGC1
GAL2
NGA3
GAL4
SIA5

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	94.68Å 94.68Å 181.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	81.99 – 2.30 81.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (81.99-2.30) 99.6 (81.99-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.29Å)	Xtriage
Refinement program	REFMAC 1.19.1_4122, PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.203 , 0.247 (Not available) , 0.230	Depositor DCC
R_{free} test set	2024 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	35.6	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 32.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.048 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7069	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NGA, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/3527	0.38	0/4764
1	B	0.12	0/3307	0.37	0/4463
All	All	0.13	0/6834	0.37	0/9227

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3434	14	0
1	B	3238	0	3198	18	0
2	C	68	0	58	0	0
2	D	68	0	58	0	0
3	A	163	0	0	0	0
3	B	74	0	0	1	0
All	All	7069	0	6748	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:965:ASN:ND2	3:B:1401:HOH:O	2.27	0.67
1:B:978:SER:HG	1:B:1054:HIS:N	2.00	0.59
1:B:1103:LEU:HD23	1:B:1242:LYS:HD3	1.86	0.57
1:B:905:TYR:HB2	1:B:931:GLU:HG3	1.88	0.54
1:B:1003:VAL:HG22	1:B:1043:ASP:HB3	1.91	0.52
1:A:991:TRP:HB3	1:A:1003:VAL:HG23	1.97	0.47
1:A:995:ASP:HA	1:A:1053:ILE:HG13	1.97	0.47
1:A:1005:PHE:CG	1:A:1037:ILE:HG13	2.49	0.47
1:B:956:PRO:O	1:B:1071:ARG:NH2	2.49	0.46
1:B:1034:LYS:HG2	1:B:1044:GLN:HG3	1.98	0.46
1:A:1301:PRO:HB2	1:A:1302:LEU:HD12	1.97	0.46
1:A:930:ILE:HB	1:A:1061:PHE:HB2	1.98	0.46
1:A:1247:ASP:HB3	1:A:1253:ILE:HD11	1.99	0.45
1:B:930:ILE:HB	1:B:1061:PHE:HB2	1.98	0.45
1:B:935:LYS:O	1:B:939:VAL:HG23	2.17	0.45
1:B:1185:ARG:HD3	1:B:1228:VAL:HG22	1.99	0.45
1:A:889:TYR:CE2	1:A:917:LYS:HD3	2.53	0.44
1:B:959[A]:PHE:HZ	1:B:1298:ARG:H	1.64	0.44
1:B:1009:GLN:HA	1:B:1017:ILE:HD11	2.01	0.43
1:A:1185:ARG:HG2	1:A:1228:VAL:HG22	2.01	0.42
1:A:1141:TYR:CE2	1:A:1143:LYS:HB2	2.54	0.42
1:B:1025:ILE:HA	1:B:1034:LYS:O	2.20	0.42
1:A:934:LEU:HB3	1:A:938:ILE:HD11	2.02	0.41
1:B:1275:ARG:HA	1:B:1275:ARG:HD3	1.90	0.41
1:B:1024:THR:HB	1:B:1036:TYR:HB2	2.02	0.41
1:A:991:TRP:CD2	1:A:1025:ILE:HG21	2.55	0.41
1:A:1241:CYS:HA	1:A:1286:CYS:O	2.21	0.41
1:B:1042:ILE:HG22	1:B:1043:ASP:N	2.34	0.41
1:B:1246:GLN:NE2	1:B:1252:ASP:OD1	2.54	0.41
1:A:1230:LYS:HE3	1:A:1230:LYS:HB3	1.88	0.40
1:B:1274:ASN:N	1:B:1274:ASN:OD1	2.54	0.40
1:A:886:SER:HB3	1:A:897:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/433 (96%)	394 (95%)	20 (5%)	0	100	100
1	B	377/433 (87%)	361 (96%)	16 (4%)	0	100	100
All	All	791/866 (91%)	755 (95%)	36 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/400 (97%)	379 (98%)	8 (2%)	47	66
1	B	362/400 (90%)	350 (97%)	12 (3%)	33	50
All	All	749/800 (94%)	729 (97%)	20 (3%)	39	58

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	880	THR
1	A	893	ASP
1	A	895	ILE
1	A	946	ASN
1	A	993	PHE
1	A	998	GLU
1	A	1071	ARG
1	A	1208	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	879	ILE
1	B	904	ILE
1	B	931	GLU
1	B	961	SER
1	B	983	SER
1	B	1071	ARG
1	B	1087	GLU
1	B	1143	LYS
1	B	1150	MET
1	B	1246	GLN
1	B	1275	ARG
1	B	1277	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1044	GLN
1	A	1175	ASN
1	B	941	ASN
1	B	1070	HIS
1	B	1148	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1	2	12,12,12	0.59	0	17,17,17	1.20	2 (11%)
2	GAL	C	2	2	11,11,12	0.80	0	15,15,17	1.38	3 (20%)
2	NGA	C	3	2	14,14,15	0.89	1 (7%)	17,19,21	1.60	2 (11%)
2	GAL	C	4	2	11,11,12	0.80	1 (9%)	15,15,17	1.23	1 (6%)
2	SIA	C	5	2	20,20,21	1.58	2 (10%)	21,28,31	1.58	3 (14%)
2	BGC	D	1	2	12,12,12	0.57	0	17,17,17	1.88	2 (11%)
2	GAL	D	2	2	11,11,12	1.24	1 (9%)	15,15,17	1.30	2 (13%)
2	NGA	D	3	2	14,14,15	0.75	1 (7%)	17,19,21	1.67	4 (23%)
2	GAL	D	4	2	11,11,12	0.74	1 (9%)	15,15,17	1.20	2 (13%)
2	SIA	D	5	2	20,20,21	1.69	3 (15%)	21,28,31	1.39	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	GAL	C	2	2	-	1/2/19/22	0/1/1/1
2	NGA	C	3	2	-	1/6/23/26	0/1/1/1
2	GAL	C	4	2	-	1/2/19/22	0/1/1/1
2	SIA	C	5	2	-	0/18/34/38	0/1/1/1
2	BGC	D	1	2	-	2/2/22/22	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	NGA	D	3	2	-	2/6/23/26	0/1/1/1
2	GAL	D	4	2	-	0/2/19/22	0/1/1/1
2	SIA	D	5	2	-	5/18/34/38	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	5	SIA	C2-C1	5.55	1.59	1.52
2	C	5	SIA	C2-C1	5.26	1.58	1.52
2	D	5	SIA	O6-C2	2.43	1.48	1.43
2	D	5	SIA	C7-C6	2.41	1.55	1.52
2	C	5	SIA	O6-C2	2.16	1.47	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	NGA	O3-C3	-2.14	1.37	1.43
2	D	4	GAL	O3-C3	-2.07	1.37	1.43
2	C	4	GAL	O3-C3	-2.04	1.37	1.43
2	C	3	NGA	O3-C3	-2.03	1.37	1.43
2	D	2	GAL	O5-C5	-2.03	1.39	1.43

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	BGC	O4-C4-C3	6.02	124.56	110.38
2	C	3	NGA	O3-C3-C2	5.15	120.09	109.40
2	C	5	SIA	O1A-C1-C2	-5.12	111.78	122.85
2	C	1	BGC	O4-C4-C3	-3.72	101.62	110.38
2	C	4	GAL	O3-C3-C2	3.69	117.58	110.05
2	D	1	BGC	O4-C4-C5	-3.69	100.24	109.32
2	D	3	NGA	O5-C1-C2	-3.67	105.61	111.29
2	D	3	NGA	O3-C3-C4	3.51	118.66	110.38
2	D	4	GAL	O3-C3-C4	-3.25	102.71	110.38
2	C	1	BGC	O4-C4-C5	3.15	117.09	109.32
2	D	5	SIA	C5-N5-C10	2.91	129.92	123.11
2	C	2	GAL	C2-C3-C4	-2.77	105.99	110.86
2	C	3	NGA	O3-C3-C4	-2.52	104.44	110.38
2	C	5	SIA	O1B-C1-O1A	2.51	129.78	124.08
2	D	5	SIA	O1B-C1-O1A	2.50	129.76	124.08
2	D	5	SIA	O1A-C1-C2	-2.43	117.60	122.85
2	D	3	NGA	O3-C3-C2	2.36	114.31	109.40
2	C	5	SIA	O1B-C1-C2	2.20	118.44	112.71
2	D	3	NGA	C4-C3-C2	2.20	114.24	111.02
2	D	2	GAL	C2-C3-C4	-2.17	107.04	110.86
2	C	2	GAL	C1-O5-C5	2.08	114.97	112.19
2	C	2	GAL	O2-C2-C1	2.06	113.94	109.22
2	D	2	GAL	C1-O5-C5	2.06	114.95	112.19
2	D	4	GAL	C1-O5-C5	2.02	114.89	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	5	SIA	C7-C8-C9-O9
2	D	5	SIA	O8-C8-C9-O9
2	D	3	NGA	O5-C5-C6-O6
2	D	3	NGA	C4-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	1	BGC	C4-C5-C6-O6
2	D	5	SIA	C4-C5-N5-C10
2	C	3	NGA	O5-C5-C6-O6
2	D	1	BGC	O5-C5-C6-O6
2	C	2	GAL	O5-C5-C6-O6
2	D	5	SIA	C6-C7-C8-C9
2	C	4	GAL	C4-C5-C6-O6
2	D	5	SIA	C6-C5-N5-C10

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	420/433 (96%)	-0.05	11 (2%) 57 59	18, 32, 59, 93	0
1	B	390/433 (90%)	0.71	53 (13%) 7 7	26, 46, 80, 99	1 (0%)
All	All	810/866 (93%)	0.32	64 (7%) 18 20	18, 40, 73, 99	1 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1046	PRO	6.4
1	B	1277	ILE	5.2
1	B	879	ILE	4.4
1	B	942	SER	4.0
1	A	1174	GLY	4.0
1	B	993	PHE	3.8
1	B	940	TYR	3.7
1	B	991	TRP	3.7
1	B	1042	ILE	3.5
1	B	1054	HIS	3.3
1	B	1033	SER	3.3
1	B	1045	LYS	3.2
1	B	1276	GLN	3.2
1	B	992	THR	3.2
1	B	944	TYR	3.1
1	B	1003	VAL	3.1
1	B	979	GLY	3.0
1	A	877	LYS	3.0
1	B	1222	ASN	3.0
1	A	879	ILE	2.9
1	B	1041	LEU	2.9
1	B	911	TYR	2.8
1	B	1027	ASN	2.7
1	A	1173	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1087	GLU	2.6
1	B	1029	ARG	2.6
1	A	1236	GLY	2.6
1	B	1098	SER	2.6
1	B	941	ASN	2.5
1	A	1302	LEU	2.5
1	B	1025	ILE	2.5
1	B	1284	LEU	2.5
1	B	937	ALA	2.5
1	B	1171	TYR	2.4
1	B	1300	ARG	2.4
1	B	1035	ILE	2.4
1	A	959	PHE	2.4
1	B	960	ASN	2.4
1	B	976	ASN	2.4
1	B	1231	SER	2.3
1	B	990	ILE	2.3
1	B	945	GLU	2.3
1	B	880	THR	2.3
1	A	1194	ASN	2.3
1	B	1026	THR	2.3
1	B	977	ASN	2.2
1	B	978	SER	2.2
1	B	1302	LEU	2.2
1	B	1090	ILE	2.2
1	B	959[A]	PHE	2.2
1	B	936	LYS	2.2
1	B	1088	LYS	2.2
1	A	1237	ILE	2.1
1	B	1175	ASN	2.1
1	B	943	MET	2.1
1	B	1094	TYR	2.1
1	B	939	VAL	2.1
1	B	1044	GLN	2.1
1	B	1239	ASN	2.1
1	B	974	MET	2.1
1	A	878	ASN	2.1
1	A	891	ASP	2.1
1	B	980	TRP	2.0
1	B	1091	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	D	1	12/12	0.45	0.18	95,107,113,128	0
2	BGC	C	1	12/12	0.47	0.16	88,105,107,108	0
2	GAL	C	2	11/12	0.53	0.18	90,93,102,103	0
2	GAL	D	2	11/12	0.60	0.18	86,97,107,113	0
2	SIA	C	5	20/21	0.82	0.13	60,66,72,73	0
2	SIA	D	5	20/21	0.82	0.13	33,48,69,72	0
2	NGA	C	3	14/15	0.83	0.14	61,73,85,91	0
2	GAL	C	4	11/12	0.87	0.12	51,57,63,64	0
2	NGA	D	3	14/15	0.90	0.10	40,51,63,64	0
2	GAL	D	4	11/12	0.92	0.07	29,38,44,46	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.