



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:53 PM UTC

PDB ID : 1QQJ / pdb_00001qqj
Title : CRYSTAL STRUCTURE OF MOUSE FUMARYLACETOACETATE HYDROLASE REFINED AT 1.55 ANGSTROM RESOLUTION
Authors : Timm, D.E.; Mueller, H.A.; Bhanumoorthy, P.; Harp, J.M.; Bunick, G.J.
Deposited on : 1999-06-07
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	
1	B	419	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	B	2000	-	X	-	-
4	CAC	A	2004	-	X	-	-
4	CAC	B	2003	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7362 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FUMARYLACETOACETATE HYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	0	0
			3224	2050	562	590	22			
1	B	416	Total	C	N	O	S	0	0	0
			3224	2050	562	590	22			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is CACODYLATE ION (CCD ID: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	As	C	O	0	0
			5	1	2	2		
4	B	1	Total	As	C	O	0	0
			5	1	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	469	Total	O	0	0
			469	469		
5	B	425	Total	O	0	0
			425	425		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FUMARYLACETOACETATE HYDROLASE



• Molecule 1: FUMARYLACETOACETATE HYDROLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.00Å 109.02Å 65.24Å 90.00° 95.68° 90.00°	Depositor
Resolution (Å)	29.10 – 1.55 29.10 – 1.55	Depositor EDS
% Data completeness (in resolution range)	92.5 (29.10-1.55) 92.1 (29.10-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.55Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.183 , 0.211 0.176 , 0.202	Depositor DCC
R_{free} test set	5952 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7362	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CAC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/3313	0.98	6/4499 (0.1%)
1	B	0.43	0/3313	1.00	5/4499 (0.1%)
All	All	0.42	0/6626	0.99	11/8998 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	379	GLN	CB-CG-CD	9.79	129.24	112.60
1	A	126	ASP	CA-CB-CG	-6.07	106.53	112.60
1	B	201	GLU	CG-CD-OE2	-6.04	104.51	118.40
1	A	238	ASP	CA-CB-CG	6.00	118.60	112.60
1	B	183	ASP	CA-C-O	5.64	127.77	121.40
1	B	126	ASP	CA-CB-CG	-5.45	107.15	112.60
1	A	95	ARG	CB-CA-C	-5.34	102.42	110.92
1	A	199	GLU	CG-CD-OE2	5.23	130.43	118.40
1	A	201	GLU	CG-CD-OE2	-5.22	106.40	118.40
1	B	204	PHE	CA-CB-CG	5.13	118.93	113.80
1	A	95	ARG	N-CA-C	5.10	117.24	111.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3224	0	3167	19	0
1	B	3224	0	3167	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	469	0	0	5	0
5	B	425	0	0	6	0
All	All	7362	0	6340	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:274:VAL:HB	1:B:275:PRO:HD2	1.67	0.76
1:A:322:MET:HE2	5:A:2063:HOH:O	1.93	0.67
1:A:125:THR:OG1	1:A:333:HIS:HE1	1.76	0.67
1:B:387:GLY:HA2	1:B:411:LYS:HE3	1.82	0.62
1:A:274:VAL:HB	1:A:275:PRO:HD2	1.81	0.62
1:A:157:VAL:HG12	1:B:247:LEU:HD12	1.84	0.58
1:A:247:LEU:HD12	1:B:157:VAL:HG12	1.84	0.58
1:B:144:LYS:H	1:B:144:LYS:HD2	1.71	0.55
1:A:154:HIS:HD2	5:A:2007:HOH:O	1.91	0.53
1:B:130:SER:HB2	1:B:322:MET:SD	2.50	0.52
1:B:144:LYS:H	1:B:144:LYS:CD	2.23	0.51
1:B:83:ALA:HA	1:B:86:GLN:HE21	1.76	0.51
1:B:416:LEU:HD12	5:B:2052:HOH:O	2.10	0.49
1:A:322:MET:HE3	1:A:324:TRP:O	2.14	0.48
1:A:309:SER:H	1:A:377:GLN:HE22	1.61	0.48
1:A:1:MET:HG2	5:A:2273:HOH:O	2.12	0.48
1:A:60:HIS:HD2	5:A:2412:HOH:O	1.97	0.47
1:B:182:PRO:HG3	5:B:2017:HOH:O	2.14	0.47
1:B:135:THR:HG23	1:B:147:ALA:HB1	1.97	0.46
1:A:138:GLY:HA3	1:A:147:ALA:O	2.16	0.45
1:A:79:LYS:HG3	5:A:2099:HOH:O	2.15	0.45
1:A:280:ASP:HA	1:A:281:PRO:C	2.41	0.44
1:B:280:ASP:HA	1:B:281:PRO:C	2.43	0.44
1:B:245:VAL:HA	1:B:246:PRO:HA	1.76	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:VAL:HA	1:A:246:PRO:HA	1.74	0.43
1:B:117:LEU:HD11	5:B:2148:HOH:O	2.17	0.43
1:A:237:ARG:HH11	1:A:237:ARG:HD3	1.62	0.42
1:B:139:ILE:HG23	1:B:144:LYS:HG3	2.02	0.41
1:B:322:MET:HE2	1:B:348:SER:HB2	2.01	0.41
1:B:278:LYS:HE2	5:B:2379:HOH:O	2.20	0.41
1:A:181:ARG:HD2	1:A:242:TRP:CZ3	2.56	0.41
1:A:282:LYS:HA	1:A:282:LYS:HD3	1.82	0.41
1:B:182:PRO:HD2	5:B:2012:HOH:O	2.19	0.41
1:B:237:ARG:HH11	1:B:237:ARG:HD3	1.67	0.41
1:B:274:VAL:HB	1:B:275:PRO:CD	2.45	0.41
1:B:321:HIS:HE1	5:B:2072:HOH:O	2.04	0.41
1:A:174:ARG:HD3	1:A:416:LEU:HB3	2.02	0.40
1:A:149:LEU:HD12	1:B:246:PRO:HD3	2.03	0.40
1:B:179:GLN:HG2	1:B:242:TRP:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/419 (99%)	400 (97%)	14 (3%)	0	100	100
1	B	414/419 (99%)	403 (97%)	11 (3%)	0	100	100
All	All	828/838 (99%)	803 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	351/353 (99%)	346 (99%)	5 (1%)	59	33
1	B	351/353 (99%)	347 (99%)	4 (1%)	65	43
All	All	702/706 (99%)	693 (99%)	9 (1%)	61	36

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	ASP
1	A	101	GLU
1	A	198	MET
1	A	330	LEU
1	A	416	LEU
1	B	63	ASP
1	B	144	LYS
1	B	183	ASP
1	B	184	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	HIS
1	A	154	HIS
1	A	241	GLN
1	A	291	GLN
1	A	333	HIS
1	A	377	GLN
1	A	407	GLN
1	B	27	ASN
1	B	86	GLN
1	B	184	ASN
1	B	291	GLN
1	B	321	HIS
1	B	379	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	CAC	A	2004	-	2,4,4	3.50	2 (100%)	4,6,6	4.78	4 (100%)
3	ACT	B	2000	-	3,3,3	2.49	2 (66%)	3,3,3	1.35	1 (33%)
4	CAC	B	2003	-	2,4,4	3.67	2 (100%)	4,6,6	4.26	3 (75%)
3	ACT	A	2001	-	3,3,3	2.53	2 (66%)	3,3,3	1.31	0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	2003	CAC	AS-C1	4.18	2.00	1.90
4	A	2004	CAC	AS-C1	3.94	1.99	1.90
3	B	2000	ACT	CH3-C	3.59	1.63	1.49
3	A	2001	ACT	CH3-C	3.54	1.63	1.49
4	B	2003	CAC	AS-C2	3.09	1.97	1.90
4	A	2004	CAC	AS-C2	3.00	1.97	1.90
3	A	2001	ACT	OXT-C	-2.58	1.18	1.30
3	B	2000	ACT	OXT-C	-2.38	1.19	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2004	CAC	O1-AS-C1	-8.11	101.42	111.50
4	B	2003	CAC	O1-AS-C1	-7.51	102.16	111.50
4	A	2004	CAC	O2-AS-C1	3.16	113.50	105.84
4	A	2004	CAC	O1-AS-C2	-2.79	108.02	111.50
4	A	2004	CAC	O2-AS-C2	2.77	112.55	105.84
4	B	2003	CAC	O2-AS-C2	2.72	112.43	105.84
4	B	2003	CAC	O2-AS-C1	2.24	111.28	105.84
3	B	2000	ACT	OXT-C-O	-2.22	113.80	122.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/419 (99%)	-0.06	15 (3%) 46 54	9, 14, 27, 37	0
1	B	416/419 (99%)	0.10	17 (4%) 41 49	9, 16, 30, 41	0
All	All	832/838 (99%)	0.02	32 (3%) 44 52	9, 15, 29, 41	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	416	LEU	5.4
1	A	1	MET	5.1
1	A	307	GLY	4.9
1	A	8	GLU	4.5
1	B	183	ASP	4.0
1	A	144	LYS	3.5
1	B	147	ALA	3.4
1	B	180	MET	3.2
1	B	146	ASN	3.1
1	B	100	LYS	3.0
1	B	104	GLN	2.9
1	B	194	ARG	2.8
1	A	91	ALA	2.6
1	A	93	GLN	2.5
1	A	310	GLN	2.5
1	B	310	GLN	2.5
1	B	60	HIS	2.5
1	A	141	PHE	2.5
1	A	94	ALA	2.5
1	B	416	LEU	2.5
1	B	141	PHE	2.4
1	A	87	ASN	2.4
1	A	92	SER	2.3
1	B	1	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	182	PRO	2.3
1	A	25	GLN	2.2
1	A	282	LYS	2.2
1	B	184	ASN	2.1
1	A	90	SER	2.0
1	B	56	SER	2.0
1	B	309	SER	2.0
1	B	52	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CAC	A	2004	5/5	0.84	0.24	33,35,36,37	0
4	CAC	B	2003	5/5	0.87	0.22	34,35,36,37	0
2	CA	B	420	1/1	0.93	0.21	18,18,18,18	0
3	ACT	B	2000	4/4	0.93	0.10	20,20,21,21	0
2	CA	A	420	1/1	0.96	0.16	15,15,15,15	0
3	ACT	A	2001	4/4	0.97	0.06	19,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.