



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 09:40 AM UTC

PDB ID : 4QQF / pdb_00004qqf
Title : Crystal structure of mitochondrial import inner membrane translocase subunit TIM50
Authors : Li, J.Z.
Deposited on : 2014-06-27
Resolution : 2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	202	
1	B	202	
1	C	202	
1	D	202	
1	E	202	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	202	18% 63% 26% 5% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9432 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitochondrial import inner membrane translocase subunit TIM50.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	187	Total	C	N	O	S	0	0	0
			1550	1011	249	286	4			
1	A	198	Total	C	N	O	S	0	0	0
			1641	1070	263	302	6			
1	B	181	Total	C	N	O	S	0	0	0
			1496	973	242	277	4			
1	C	193	Total	C	N	O	S	0	0	0
			1602	1045	256	296	5			
1	E	182	Total	C	N	O	S	0	0	0
			1504	981	241	278	4			
1	F	191	Total	C	N	O	S	0	0	0
			1584	1035	251	293	5			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	160	GLY	-	expression tag	UNP Q02776
D	161	SER	-	expression tag	UNP Q02776
D	162	HIS	-	expression tag	UNP Q02776
D	163	MET	-	expression tag	UNP Q02776
A	160	GLY	-	expression tag	UNP Q02776
A	161	SER	-	expression tag	UNP Q02776
A	162	HIS	-	expression tag	UNP Q02776
A	163	MET	-	expression tag	UNP Q02776
B	160	GLY	-	expression tag	UNP Q02776
B	161	SER	-	expression tag	UNP Q02776
B	162	HIS	-	expression tag	UNP Q02776
B	163	MET	-	expression tag	UNP Q02776
C	160	GLY	-	expression tag	UNP Q02776
C	161	SER	-	expression tag	UNP Q02776
C	162	HIS	-	expression tag	UNP Q02776
C	163	MET	-	expression tag	UNP Q02776

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	160	GLY	-	expression tag	UNP Q02776
E	161	SER	-	expression tag	UNP Q02776
E	162	HIS	-	expression tag	UNP Q02776
E	163	MET	-	expression tag	UNP Q02776
F	160	GLY	-	expression tag	UNP Q02776
F	161	SER	-	expression tag	UNP Q02776
F	162	HIS	-	expression tag	UNP Q02776
F	163	MET	-	expression tag	UNP Q02776

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	E	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	12	Total O 12 12	0	0
3	A	17	Total O 17 17	0	0
3	B	6	Total O 6 6	0	0
3	C	6	Total O 6 6	0	0
3	E	6	Total O 6 6	0	0
3	F	2	Total O 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitochondrial import inner membrane translocase subunit TIM50




- Molecule 1: Mitochondrial import inner membrane translocase subunit TIM50




- Molecule 1: Mitochondrial import inner membrane translocase subunit TIM50





- Molecule 1: Mitochondrial import inner membrane translocase subunit TIM50






- Molecule 1: Mitochondrial import inner membrane translocase subunit TIM50



- Molecule 1: Mitochondrial import inner membrane translocase subunit TIM50



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	164.29Å 164.29Å 150.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.52 – 2.67 38.52 – 2.67	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.52-2.67) 95.6 (38.52-2.67)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.69Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.212 , 0.252 0.238 , 0.268	Depositor DCC
R_{free} test set	2937 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	69.7	Xtriage
Anisotropy	0.244	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9432	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	1/1696 (0.1%)	1.19	11/2309 (0.5%)
1	B	0.64	0/1543	1.02	4/2101 (0.2%)
1	C	0.63	1/1655 (0.1%)	1.10	10/2255 (0.4%)
1	D	0.68	3/1601 (0.2%)	1.13	9/2182 (0.4%)
1	E	0.65	2/1552 (0.1%)	0.98	5/2114 (0.2%)
1	F	0.64	3/1636 (0.2%)	1.08	9/2229 (0.4%)
All	All	0.66	10/9683 (0.1%)	1.09	48/13190 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	173	GLU	C-N	6.61	1.41	1.33
1	F	173	GLU	C-N	6.55	1.41	1.33
1	D	177	PRO	N-CD	5.82	1.55	1.47
1	D	174	PRO	N-CD	5.61	1.55	1.47
1	F	174	PRO	N-CD	5.34	1.55	1.47

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	185	PRO	CA-C-N	12.67	133.43	120.38
1	C	185	PRO	C-N-CA	12.67	133.43	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	ARG	CA-C-N	-11.36	108.09	119.56
1	A	339	ARG	C-N-CA	-11.36	108.09	119.56
1	C	186	PRO	CA-C-N	-10.34	108.87	120.04

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	212	GLY	Peptide
1	B	356	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1641	0	1590	41	0
1	B	1496	0	1467	16	0
1	C	1602	0	1557	14	0
1	D	1550	0	1512	48	0
1	E	1504	0	1476	30	0
1	F	1584	0	1544	77	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	17	0	0	0	1
3	B	6	0	0	3	0
3	C	6	0	0	0	0
3	D	12	0	0	1	0
3	E	6	0	0	0	0
3	F	2	0	0	0	0
All	All	9432	0	9146	214	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 214 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:207:TRP:CD1	1:F:213:TRP:HZ3	1.48	1.31
1:F:207:TRP:CE2	1:F:213:TRP:CH2	2.20	1.30
1:F:207:TRP:CD1	1:F:213:TRP:CZ3	2.20	1.30
1:A:337:ASP:OD1	1:A:339:ARG:NE	1.72	1.20
1:F:207:TRP:CE2	1:F:213:TRP:CZ3	2.36	1.13

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:511:HOH:O	3:A:511:HOH:O[8_554]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/202 (97%)	183 (93%)	8 (4%)	5 (3%)	4	9
1	B	179/202 (89%)	176 (98%)	2 (1%)	1 (1%)	21	41
1	C	191/202 (95%)	181 (95%)	8 (4%)	2 (1%)	12	28
1	D	185/202 (92%)	177 (96%)	6 (3%)	2 (1%)	11	26
1	E	180/202 (89%)	176 (98%)	3 (2%)	1 (1%)	21	41
1	F	189/202 (94%)	171 (90%)	11 (6%)	7 (4%)	2	5
All	All	1120/1212 (92%)	1064 (95%)	38 (3%)	18 (2%)	7	18

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	334	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	186	PRO
1	A	187	PRO
1	A	337	ASP
1	A	338	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/188 (98%)	175 (95%)	9 (5%)	22	46
1	B	168/188 (89%)	157 (94%)	11 (6%)	15	34
1	C	180/188 (96%)	170 (94%)	10 (6%)	19	40
1	D	174/188 (93%)	167 (96%)	7 (4%)	28	53
1	E	169/188 (90%)	163 (96%)	6 (4%)	31	57
1	F	178/188 (95%)	169 (95%)	9 (5%)	21	44
All	All	1053/1128 (93%)	1001 (95%)	52 (5%)	22	46

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	205	SER
1	C	353	GLU
1	F	323	LEU
1	C	209	GLN
1	C	279	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	275	HIS
1	F	343	ASN
1	E	189	GLN
1	E	204	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	172	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	198/202 (98%)	0.25	15 (7%)	20 17	46, 60, 111, 192	0
1	B	181/202 (89%)	0.23	10 (5%)	30 26	48, 69, 104, 167	0
1	C	193/202 (95%)	0.22	11 (5%)	29 25	48, 69, 118, 141	0
1	D	187/202 (92%)	0.62	14 (7%)	20 17	24, 71, 117, 156	0
1	E	182/202 (90%)	0.31	9 (4%)	35 30	24, 68, 107, 131	0
1	F	191/202 (94%)	1.04	37 (19%)	3 2	57, 92, 148, 263	0
All	All	1132/1212 (93%)	0.44	96 (8%)	16 14	24, 70, 121, 263	0

The worst 5 of 96 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	176	PHE	8.9
1	D	176	PHE	8.7
1	A	336	LYS	7.9
1	D	174	PRO	7.8
1	A	187	PRO	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	D	401	1/1	0.81	0.42	68,68,68,68	0
2	MG	F	401	1/1	0.90	0.28	55,55,55,55	0
2	MG	B	401	1/1	0.95	0.33	47,47,47,47	0
2	MG	C	401	1/1	0.97	0.39	41,41,41,41	0
2	MG	E	401	1/1	0.98	0.40	39,39,39,39	0
2	MG	A	401	1/1	0.98	0.43	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.