



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 01:59 PM UTC

PDB ID : 6QQN / pdb_00006qqn
Title : Tubulin-TH588 complex
Authors : Patterson, J.C.; Joughin, B.A.; Prota, A.E.; Muehlethaler, T.; Jonas, O.H.; Whitman, M.A.; Varmeh, S.; Chen, S.; Balk, S.P.; Steinmetz, M.O.; Lauffenburger, D.A.; Yaffe, M.B.
Deposited on : 2019-02-18
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	0% 74% 23% •
1	C	451	0% 75% 23% •
2	B	445	3% 65% 30% 5%
2	D	445	2% 68% 26% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	143	
4	F	384	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	2GE	B	504	-	-	X	-

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 17568 atoms, of which 18 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	0	0
			3409	2159	580	648	22			
1	C	440	Total	C	N	O	S	0	1	0
			3444	2180	585	657	22			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	423	Total	C	N	O	S	0	1	0
			3344	2102	572	643	27			
2	D	421	Total	C	N	O	S	0	2	0
			3326	2089	567	642	28			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	0	0
			1000	617	181	197	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	-	initiating methionine	UNP P63043
E	4	ALA	-	expression tag	UNP P63043


- Molecule 4 is a protein called Tubulin-Tyrosine Ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	314	Total	C	N	O	S	0	0	0
			2573	1659	437	464	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	B	2	Total	Ca	0	0
			2	2		
7	C	1	Total	Ca	0	0
			1	1		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total 28	C 10	N 5	O 11	P 2	0	0
9	D	1	Total 28	C 10	N 5	O 11	P 2	0	0


- Molecule 10 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- 2GE
-
- Clc1cc(Cl)ccc1-c1ccn(CN2CC2)c1


- Molecule 12 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $\text{C}_{11}\text{H}_{18}\text{N}_5\text{O}_{12}\text{P}_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	F	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 13 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	50	Total 50	O 50	0	0
13	B	41	Total 41	O 41	0	0
13	C	117	Total 117	O 117	0	0
13	D	27	Total 27	O 27	0	0
13	E	11	Total 11	O 11	0	0
13	F	13	Total 13	O 13	0	0



• Molecule 2: Tubulin beta-2B chain




• Molecule 3: Stathmin-4



• Molecule 4: Tubulin-Tyrosine Ligase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.83Å 156.19Å 179.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.48 – 2.30 47.48 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.48-2.30) 99.9 (47.48-2.30)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.29Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.196 , 0.244 0.197 , 0.244	Depositor DCC
R_{free} test set	6477 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	56.9	Xtriage
Anisotropy	0.164	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17568	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACP, CA, GDP, MG, MES, EDO, 2GE, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/3486	0.30	0/4730
1	C	0.13	0/3522	0.31	0/4782
2	B	0.12	0/3418	0.32	0/4628
2	D	0.10	0/3399	0.28	0/4603
3	E	0.11	0/1008	0.24	0/1337
4	F	0.09	0/2630	0.28	0/3551
All	All	0.11	0/17463	0.30	0/23631

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3409	0	3322	96	0
1	C	3444	0	3356	87	0
2	B	3344	0	3227	126	0
2	D	3326	0	3205	100	0
3	E	1000	0	1018	29	0
4	F	2573	0	2550	95	0
5	A	32	0	12	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
7	C	1	0	0	0	0
8	A	4	6	6	0	0
9	B	28	0	12	1	0
9	D	28	0	12	3	0
10	B	12	0	12	0	0
11	B	19	12	12	7	0
12	F	31	0	14	3	0
13	A	50	0	0	5	0
13	B	41	0	0	0	0
13	C	117	0	0	1	0
13	D	27	0	0	1	0
13	E	11	0	0	1	0
13	F	13	0	0	1	0
All	All	17550	18	16770	498	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 498 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:173:ILE:HD12	4:F:180:HIS:HB2	1.32	1.07
1:C:270:ALA:HB3	1:C:302:MET:HE2	1.42	1.02
2:B:62:VAL:HG11	2:B:88:ARG:HG3	1.41	1.00
4:F:292:ARG:HB3	4:F:296:MET:HE3	1.43	0.96
2:B:176:LYS:HE2	2:B:207:GLU:HG3	1.54	0.90

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	432/451 (96%)	423 (98%)	9 (2%)	0	100	100
1	C	439/451 (97%)	426 (97%)	13 (3%)	0	100	100
2	B	420/445 (94%)	407 (97%)	13 (3%)	0	100	100
2	D	419/445 (94%)	404 (96%)	15 (4%)	0	100	100
3	E	117/143 (82%)	116 (99%)	1 (1%)	0	100	100
4	F	298/384 (78%)	284 (95%)	14 (5%)	0	100	100
All	All	2125/2319 (92%)	2060 (97%)	65 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/379 (97%)	366 (100%)	1 (0%)	86	93
1	C	372/379 (98%)	371 (100%)	1 (0%)	86	93
2	B	367/383 (96%)	364 (99%)	3 (1%)	73	86
2	D	366/383 (96%)	362 (99%)	4 (1%)	65	81
3	E	109/127 (86%)	109 (100%)	0	100	100
4	F	283/342 (83%)	281 (99%)	2 (1%)	76	87
All	All	1864/1993 (94%)	1853 (99%)	11 (1%)	78	89

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	389	LYS
2	D	402	LYS
4	F	348	GLN
4	F	331	GLU
1	C	241	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	282	GLN
3	E	103	GLN
2	B	436	GLN
3	E	115	HIS
1	C	300	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 9 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	GDP	B	501	6	29,30,30	1.17	4 (13%)	45,47,47	1.75	7 (15%)
5	GTP	C	501	6	33,34,34	1.00	3 (9%)	50,54,54	1.54	10 (20%)
8	EDO	A	504	-	3,3,3	0.43	0	2,2,2	0.37	0
12	ACP	F	701	6	31,33,33	1.36	5 (16%)	47,52,52	1.99	9 (19%)
9	GDP	D	501	6	29,30,30	1.17	3 (10%)	45,47,47	1.71	6 (13%)
10	MES	B	503	-	12,12,12	2.26	1 (8%)	15,16,16	1.79	2 (13%)
5	GTP	A	501	6	33,34,34	0.98	3 (9%)	50,54,54	1.53	10 (20%)
11	2GE	B	504	-	21,21,21	1.31	2 (9%)	28,30,30	2.16	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GDP	B	501	6	-	3/16/32/32	0/3/3/3
5	GTP	C	501	6	-	9/22/38/38	0/3/3/3
8	EDO	A	504	-	-	0/1/1/1	-
12	ACP	F	701	6	-	5/19/38/38	0/3/3/3
9	GDP	D	501	6	-	4/16/32/32	0/3/3/3
10	MES	B	503	-	-	0/6/14/14	0/1/1/1
5	GTP	A	501	6	-	9/22/38/38	0/3/3/3
11	2GE	B	504	-	-	4/8/10/10	0/3/3/3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	503	MES	C8-S	-7.54	1.67	1.77
12	F	701	ACP	PB-O3A	3.33	1.62	1.58
11	B	504	2GE	C02-N01	3.22	1.40	1.33
9	D	501	GDP	C5-C4	3.07	1.47	1.38
9	B	501	GDP	C5-C4	3.01	1.47	1.38

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	F	701	ACP	PB-O3A-PA	-6.05	112.64	132.37
9	B	501	GDP	C5-C4-N3	-5.98	118.88	128.39
9	D	501	GDP	C5-C4-N3	-5.84	119.10	128.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	B	504	2GE	C11-C09-CL1	-5.63	115.34	120.51
12	F	701	ACP	C5-C4-N3	-5.60	119.00	126.72

There are no chirality outliers.

5 of 34 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O3G
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	501	GTP	PB-O3B-PG-O2G

There are no ring outliers.


5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	B	501	GDP	1	0
12	F	701	ACP	3	0
9	D	501	GDP	3	0
5	A	501	GTP	1	0
11	B	504	2GE	7	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand ACP F 701



Ligand GDP D 501







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/451 (96%)	0.01	6 (1%) 73 75	44, 64, 101, 142	0
1	C	440/451 (97%)	-0.19	5 (1%) 78 79	24, 52, 79, 100	1 (0%)
2	B	423/445 (95%)	0.15	14 (3%) 49 51	24, 62, 104, 124	1 (0%)
2	D	421/445 (94%)	0.26	11 (2%) 57 59	30, 75, 106, 126	2 (0%)
3	E	121/143 (84%)	0.39	4 (3%) 49 51	54, 81, 119, 124	0
4	F	314/384 (81%)	0.63	30 (9%) 13 15	57, 91, 145, 164	0
All	All	2155/2319 (92%)	0.16	70 (3%) 50 52	24, 67, 117, 164	4 (0%)

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	248	LEU	6.2
2	D	286	LEU	4.9
2	B	249	ASN	4.9
2	D	172	MET	4.1
2	B	202	TYR	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	CA	B	505	1/1	0.79	0.16	96,96,96,96	0
7	CA	B	506	1/1	0.79	0.12	124,124,124,124	0
6	MG	F	702	1/1	0.80	0.12	83,83,83,83	0
12	ACP	F	701	31/31	0.80	0.13	89,102,159,161	0
11	2GE	B	504	19/19	0.90	0.17	73,88,107,108	31
6	MG	D	502	1/1	0.92	0.09	71,71,71,71	0
9	GDP	D	501	28/28	0.92	0.10	64,68,78,86	0
10	MES	B	503	12/12	0.93	0.10	51,60,77,82	0
8	EDO	A	504	4/4	0.93	0.10	71,86,94,98	0
7	CA	A	503	1/1	0.93	0.07	89,89,89,89	0
7	CA	C	503	1/1	0.97	0.04	72,72,72,72	0
5	GTP	A	501	32/32	0.98	0.05	37,47,53,59	0
9	GDP	B	501	28/28	0.98	0.05	33,46,53,63	0
5	GTP	C	501	32/32	0.98	0.05	37,42,50,52	0
6	MG	A	502	1/1	0.99	0.03	46,46,46,46	0
6	MG	B	502	1/1	0.99	0.04	45,45,45,45	0
6	MG	C	502	1/1	0.99	0.02	46,46,46,46	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP F 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2GE B 504:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP A 501:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.