



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:11 AM UTC

PDB ID : 7QQV / pdb_00007qqv
Title : SpCas9 bound to FANCF off-target3 DNA substrate
Authors : Pacesa, M.; Jinek, M.
Deposited on : 2022-01-10
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)
RNA backbone	3983	1005 (2.56-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	85	
2	B	1368	
3	C	28	
4	D	12	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called FANCF sgRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	85	Total	C	N	O	P	0	0	1
			1778	794	322	578	84			

- Molecule 2 is a protein called CRISPR-associated endonuclease Cas9/Csn1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1331	Total	C	N	O	S	0	0	0
			10878	6931	1887	2038	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	ALA	ASP	engineered mutation	UNP Q99ZW2
B	840	ALA	HIS	engineered mutation	UNP Q99ZW2

- Molecule 3 is a DNA chain called FANCF off-target3 target strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	28	Total	C	N	O	P	0	3	0
			641	302	127	182	30			

- Molecule 4 is a DNA chain called FANCF off-target3 non-target strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	10	Total	C	N	O	P	0	0	0
			201	97	35	60	9			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total K 2 2	0	0
6	B	7	Total K 7 7	0	0
6	C	1	Total K 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	148	Total O 148 148	0	0
7	B	347	Total O 347 347	0	0
7	C	23	Total O 23 23	0	0
7	D	6	Total O 6 6	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FANCF sgRNA

Chain A: 




• Molecule 2: CRISPR-associated endonuclease Cas9/Csn1

Chain B: 




• Molecule 3: FANCF off-target3 target strand

Chain C: 



- Molecule 4: FANCF off-target3 non-target strand

Chain D:  58% 25% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	179.02Å 67.14Å 187.77Å 90.00° 111.14° 90.00°	Depositor
Resolution (Å)	47.81 – 2.25 47.81 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.81-2.25) 86.2 (47.81-2.25)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.195 , 0.230 (Not available) , 0.221	Depositor DCC
R_{free} test set	5276 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtriage
Anisotropy	0.384	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14034	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.18	0/1990	0.37	0/3101
2	B	0.17	0/11070	0.36	0/14873
3	C	0.24	0/721	0.42	0/1113
4	D	0.27	0/224	0.54	0/344
All	All	0.18	0/14005	0.37	0/19431

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1778	0	892	26	0
2	B	10878	0	11053	91	0
3	C	641	0	347	10	0
4	D	201	0	115	3	0
5	A	2	0	0	0	0
6	A	2	0	0	0	0
6	B	7	0	0	0	0
6	C	1	0	0	0	0
7	A	148	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	347	0	0	4	0
7	C	23	0	0	2	0
7	D	6	0	0	0	0
All	All	14034	0	12407	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (117) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:178:ASN:HA	2:B:184:LEU:HD11	1.68	0.75
2:B:80:CYS:SG	7:B:1801:HOH:O	2.44	0.75
2:B:1212:ARG:NH2	2:B:1280:VAL:O	2.23	0.71
2:B:895:ARG:NH1	3:C:14[A]:DG:OP1	2.26	0.68
3:C:15[B]:DG:N7	7:C:202:HOH:O	2.28	0.67
2:B:763:MET:HE1	2:B:931:VAL:HG21	1.76	0.66
2:B:1004:LEU:HD11	2:B:1042:ILE:HD11	1.78	0.66
2:B:898:ASP:O	2:B:905:ARG:NH2	2.29	0.65
2:B:1335:ARG:NH2	4:D:2:DG:O6	2.33	0.61
2:B:844:GLN:NE2	2:B:850:ASP:OD1	2.36	0.59
2:B:623:LEU:HD13	2:B:654:ARG:HG3	1.85	0.59
1:A:5:U:H3	3:C:15[A]:DG:H22	1.51	0.59
1:A:69:A:H2'	1:A:70:C:H6	1.68	0.59
1:A:49:A:N3	2:B:1122:ARG:NH2	2.48	0.58
1:A:20:C:OP2	2:B:403:ARG:NH1	2.37	0.57
2:B:1047:LYS:O	2:B:1076:LYS:NZ	2.30	0.57
1:A:46:A:H2'	1:A:47:A:C8	2.40	0.56
2:B:1210:ARG:HA	2:B:1280:VAL:HG22	1.87	0.56
2:B:215:ARG:NH2	2:B:304:ASP:O	2.38	0.56
2:B:790:GLU:HG2	2:B:889:ALA:HA	1.87	0.56
2:B:184:LEU:HD12	2:B:299:ALA:HB2	1.87	0.55
2:B:654:ARG:HD2	2:B:656:TYR:CZ	2.40	0.55
2:B:1042:ILE:HG23	2:B:1043:MET:HG2	1.89	0.55
1:A:22:U:H2'	1:A:23:U:C6	2.42	0.55
2:B:849:ASP:OD2	2:B:895:ARG:NH2	2.40	0.55
2:B:1110:ILE:HG23	2:B:1122:ARG:HD2	1.88	0.54
2:B:489:GLN:HG3	2:B:625:LEU:HD21	1.88	0.54
1:A:59:U:OP1	2:B:467:ARG:NH2	2.38	0.54
2:B:1334:LYS:NZ	3:C:-5:DA:OP2	2.31	0.54
2:B:1215:ALA:HB2	2:B:1221:GLN:HG3	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:846:PHE:O	2:B:920:GLN:NE2	2.41	0.53
1:A:69:A:H2'	1:A:70:C:C6	2.44	0.53
2:B:391:VAL:HG12	2:B:395:ARG:HD3	1.91	0.52
2:B:788:ILE:HG23	2:B:793:SER:HB3	1.91	0.52
1:A:19:C:H2'	1:A:20:C:O4'	2.10	0.52
2:B:1143:VAL:HG13	2:B:1195:ILE:HG23	1.91	0.52
2:B:262:ALA:HB1	2:B:278:LEU:HD12	1.92	0.52
3:C:15[A]:DG:N7	7:C:202:HOH:O	2.41	0.52
2:B:1241:HIS:CD2	2:B:1245:LEU:HD21	2.45	0.51
1:A:0:G:OP1	2:B:918:LYS:NZ	2.43	0.51
2:B:137:HIS:HA	2:B:322:ILE:HD11	1.92	0.50
2:B:526:LYS:NZ	7:B:1521:HOH:O	2.44	0.50
2:B:1163:LEU:HG	2:B:1343:LEU:HD21	1.94	0.50
2:B:1205:GLU:OE1	2:B:1359:ARG:NH2	2.44	0.50
2:B:814:TYR:CZ	2:B:830:ILE:HG12	2.47	0.49
3:C:13[B]:DG:H8	3:C:13[B]:DG:H5''	1.77	0.49
2:B:308:VAL:HG11	2:B:319:ALA:HB3	1.93	0.49
1:A:77:A:H2'	1:A:78:A:C8	2.48	0.49
2:B:525:THR:HG22	2:B:545:LYS:NZ	2.28	0.48
4:D:-2:DC:H2'	4:D:-1:DC:C6	2.48	0.48
1:A:4:A:H2'	1:A:5:U:H6	1.79	0.48
2:B:557:ARG:NH2	2:B:596:ASP:OD1	2.42	0.48
2:B:704:PHE:O	2:B:708:ILE:HG12	2.14	0.48
2:B:830:ILE:O	2:B:833:LEU:HG	2.13	0.48
2:B:821:ASP:HB3	2:B:824:VAL:O	2.13	0.48
2:B:323:LYS:HE2	2:B:327:GLU:OE2	2.14	0.47
2:B:982:HIS:HD2	7:B:1505:HOH:O	1.98	0.47
2:B:518:PHE:CD1	2:B:667:ILE:HD12	2.51	0.46
2:B:654:ARG:HD3	2:B:655:ARG:N	2.30	0.46
2:B:633:GLU:HB2	2:B:652:LYS:HD2	1.97	0.46
2:B:756:PRO:HD2	2:B:939:MET:HE2	1.97	0.46
2:B:15:SER:HA	2:B:51:LEU:O	2.16	0.46
2:B:956:ILE:HG23	2:B:1008:PHE:HB3	1.98	0.46
2:B:985:HIS:CG	2:B:1087:LEU:HD22	2.51	0.46
1:A:28:A:O4'	2:B:122:ILE:HD11	2.16	0.46
2:B:870:VAL:HG13	2:B:871:PRO:HD2	1.97	0.46
1:A:70:C:H2'	1:A:71:U:C6	2.51	0.45
1:A:81:G:N1	2:B:1356:TYR:HB3	2.32	0.45
2:B:841:ILE:HD13	2:B:900:LEU:HG	1.98	0.45
2:B:787:GLY:HA3	2:B:891:LEU:HD21	1.98	0.45
2:B:781:MET:HG2	2:B:803:ASN:HA	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1306:ALA:O	2:B:1310:ILE:HG12	2.15	0.45
2:B:349:GLU:OE2	2:B:356:LYS:HE2	2.17	0.45
2:B:882:TYR:CZ	2:B:886:LEU:HD11	2.52	0.45
2:B:273:ASP:O	2:B:277:ASN:N	2.49	0.45
2:B:278:LEU:O	2:B:282:ILE:HG12	2.17	0.44
2:B:348:LYS:O	2:B:352:PHE:HB2	2.17	0.44
2:B:501:ASN:HB3	2:B:708:ILE:HD12	2.00	0.44
2:B:654:ARG:HD2	2:B:656:TYR:CE2	2.53	0.44
1:A:73:G:H5'	1:A:74:A:OP2	2.17	0.44
2:B:507:VAL:HG11	2:B:660:GLY:O	2.18	0.44
1:A:5:U:H3	3:C:15[B]:DG:H1	1.65	0.43
2:B:680:LEU:HG	2:B:684:LYS:HE3	2.00	0.43
1:A:46:A:H2'	1:A:47:A:H8	1.81	0.43
1:A:70:C:H2'	1:A:71:U:H6	1.83	0.43
2:B:387:GLU:O	2:B:391:VAL:HG23	2.19	0.43
2:B:780:ARG:NH1	2:B:806:LEU:O	2.46	0.43
2:B:1308:ASN:HB3	2:B:1326:TYR:CD1	2.54	0.43
2:B:345:GLU:H	2:B:345:GLU:CD	2.27	0.43
2:B:174:LEU:HA	2:B:174:LEU:HD12	1.77	0.43
2:B:972:PHE:HE1	2:B:1084:ARG:HD3	1.84	0.43
3:C:16:DC:H2'	3:C:17:DT:C6	2.53	0.43
1:A:74:A:OP1	1:A:74:A:H8	2.01	0.43
2:B:784:ILE:HD12	2:B:806:LEU:HD13	2.00	0.43
2:B:258:LEU:HD21	2:B:282:ILE:HD11	2.00	0.43
1:A:54:G:H2'	1:A:55:C:C6	2.54	0.42
2:B:314:LYS:HB3	2:B:314:LYS:HE2	1.80	0.42
2:B:478:PHE:CE1	2:B:482:VAL:HG21	2.53	0.42
2:B:962:LEU:HD23	2:B:962:LEU:HA	1.92	0.42
1:A:18:A:N7	7:A:201:HOH:O	2.37	0.42
2:B:335:LEU:HD21	2:B:376:ILE:HD11	2.02	0.42
2:B:1136:SER:HA	4:D:2:DG:O3'	2.20	0.42
1:A:30:C:H2'	1:A:31:U:C6	2.55	0.42
2:B:830:ILE:C	2:B:832:ARG:H	2.28	0.42
2:B:185:PHE:HE2	2:B:203:ALA:HB1	1.85	0.41
2:B:258:LEU:C	2:B:260:GLU:H	2.27	0.41
2:B:901:THR:O	2:B:904:GLU:HG2	2.20	0.41
2:B:733:ILE:HD12	2:B:733:ILE:HA	1.85	0.41
2:B:1122:ARG:HD3	7:B:1569:HOH:O	2.20	0.41
3:C:9:DG:H2'	3:C:10:DA:C8	2.56	0.41
1:A:7:C:H2'	1:A:8:C:O4'	2.21	0.41
3:C:13[B]:DG:H2'	3:C:14[B]:DG:O4'	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:U:H1'	2:B:104:SER:O	2.21	0.40
1:A:70:C:C2	1:A:71:U:C5	3.08	0.40
2:B:526:LYS:HD3	2:B:526:LYS:HA	1.86	0.40
2:B:528:LYS:HE3	2:B:539:PHE:CE1	2.56	0.40
2:B:1236:LEU:HD23	2:B:1236:LEU:HA	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	1323/1368 (97%)	1283 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	1194/1225 (98%)	1177 (99%)	17 (1%)	59	70

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	75	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	174	LEU
2	B	175	ASN
2	B	305	ILE
2	B	419	LEU
2	B	532	GLU
2	B	766	GLU
2	B	801	VAL
2	B	830	ILE
2	B	832	ARG
2	B	870	VAL
2	B	1007	GLU
2	B	1084	ARG
2	B	1123	LYS
2	B	1212	ARG
2	B	1246	LYS
2	B	1280	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	265	GLN
2	B	277	ASN
2	B	459	ASN
2	B	501	ASN
2	B	690	ASN
2	B	723	HIS
2	B	758	ASN
2	B	826	GLN
2	B	885	GLN
2	B	985	HIS
2	B	1101	GLN
2	B	1221	GLN
2	B	1241	HIS
2	B	1262	HIS
2	B	1297	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	82/85 (96%)	11 (13%)	0

All (11) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	8	C
1	A	20	C
1	A	28	A
1	A	29	G
1	A	36	A
1	A	51	A
1	A	56	U
1	A	59	U
1	A	68	A
1	A	74	A
1	A	77	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	85/85 (100%)	-0.49	0 100 100	39, 58, 132, 159	0
2	B	1331/1368 (97%)	0.24	29 (2%) 62 63	34, 59, 100, 159	0
3	C	28/28 (100%)	-0.12	0 100 100	35, 58, 101, 129	3 (10%)
4	D	10/12 (83%)	0.09	0 100 100	50, 69, 111, 136	0
All	All	1454/1493 (97%)	0.18	29 (1%) 65 65	34, 59, 105, 159	3 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1245	LEU	5.2
2	B	713	VAL	3.4
2	B	1039	TYR	3.3
2	B	1366	GLY	2.9
2	B	1249	PRO	2.8
2	B	1037	PHE	2.8
2	B	775	LYS	2.7
2	B	802	GLU	2.7
2	B	176	PRO	2.6
2	B	259	ALA	2.5
2	B	1051	THR	2.5
2	B	174	LEU	2.5
2	B	195	LEU	2.5
2	B	203	ALA	2.4
2	B	1032	ALA	2.4
2	B	312	ILE	2.4
2	B	1034	ALA	2.3
2	B	655	ARG	2.3
2	B	1257	LEU	2.3
2	B	714	SER	2.2
2	B	801	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	1259	VAL	2.2
2	B	293	ALA	2.2
2	B	830	ILE	2.1
2	B	3	LYS	2.1
2	B	1145	VAL	2.1
2	B	868	ASP	2.1
2	B	921	LEU	2.0
2	B	766	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	K	B	1401	1/1	0.85	0.15	109,109,109,109	0
6	K	C	101	1/1	0.90	0.16	111,111,111,111	0
6	K	B	1402	1/1	0.92	0.08	60,60,60,60	0
6	K	B	1404	1/1	0.94	0.07	82,82,82,82	0
6	K	B	1405	1/1	0.95	0.06	72,72,72,72	0
6	K	B	1406	1/1	0.95	0.07	94,94,94,94	0
6	K	B	1403	1/1	0.95	0.09	72,72,72,72	0
6	K	B	1407	1/1	0.96	0.08	64,64,64,64	0
5	MG	A	101	1/1	0.96	0.12	82,82,82,82	0
6	K	A	103	1/1	0.97	0.14	56,56,56,56	0
6	K	A	104	1/1	0.98	0.07	53,53,53,53	0
5	MG	A	102	1/1	0.99	0.03	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.