



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:50 PM UTC

PDB ID : 8QQG / pdb_00008qqg
Title : Structure of BRAF in Complex With Exarafenib (KIN-2787).
Authors : Schmitt, A.; Costanzi, E.; Kania, R.; Chen, Y.K.
Deposited on : 2023-10-04
Resolution : 2.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3580 (3.00-2.96)
Clashscore	190562	3904 (3.00-2.96)
Ramachandran outliers	187476	3761 (3.00-2.96)
Sidechain outliers	187428	3764 (3.00-2.96)
RSRZ outliers	180081	3579 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	11% 95% • •
1	B	275	8% 93% • 5%
1	C	275	12% 93% • •

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6597 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	56	0	0
			2165	1388	379	385	13			
1	B	262	Total	C	N	O	S	26	1	0
			2102	1347	367	375	13			
1	C	266	Total	C	N	O	S	57	0	0
			2132	1366	372	381	13			

- Molecule 2 is Exarafenib (CCD ID: WJ9) (formula: $C_{26}H_{34}F_3N_5O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			37	26	3	5	3		
2	B	1	Total	C	F	N	O	0	0
			37	26	3	5	3		
2	C	1	Total	C	F	N	O	0	0
			37	26	3	5	3		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	Cl 3	0	0
3	B	2	Total 2	Cl 2	0	0
3	C	2	Total 2	Cl 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	28	Total 28	O 28	0	0
4	B	33	Total 33	O 33	0	0
4	C	19	Total 19	O 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	206.78Å 206.78Å 149.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	146.21 – 2.98 146.21 – 2.98	Depositor EDS
% Data completeness (in resolution range)	88.6 (146.21-2.98) 88.6 (146.21-2.98)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.224 , 0.261 0.226 , 0.258	Depositor DCC
R_{free} test set	522 reflections (1.77%)	wwPDB-VP
Wilson B-factor (Å ²)	57.0	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6597	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: WJ9, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	0/2213	1.50	2/2983 (0.1%)
1	B	1.03	0/2146	1.50	1/2896 (0.0%)
1	C	1.04	0/2176	1.51	5/2932 (0.2%)
All	All	1.03	0/6535	1.51	8/8811 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	628	ASP	CA-CB-CG	6.66	119.26	112.60
1	B	448	ASP	CA-CB-CG	6.43	119.03	112.60
1	A	604	SER	CA-C-N	5.38	125.08	119.92
1	A	604	SER	C-N-CA	5.38	125.08	119.92
1	C	668	VAL	CA-C-N	5.22	125.73	119.94
1	C	668	VAL	C-N-CA	5.22	125.73	119.94
1	C	594	PHE	CA-C-N	5.03	125.12	120.34
1	C	594	PHE	C-N-CA	5.03	125.12	120.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2165	0	2195	3	0
1	B	2102	0	2143	2	0
1	C	2132	0	2168	4	0
2	A	37	0	0	0	0
2	B	37	0	0	0	0
2	C	37	0	0	0	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	28	0	0	1	0
4	B	33	0	0	0	0
4	C	19	0	0	0	0
All	All	6597	0	6506	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:619:MET:HE3	1:C:623:VAL:HG12	1.91	0.53
1:C:581:ILE:HG23	1:C:589:VAL:HG13	1.93	0.51
1:A:581:ILE:HG23	1:A:589:VAL:HG13	1.94	0.49
1:B:581:ILE:HG23	1:B:589:VAL:HG13	1.95	0.48
1:C:612:LEU:HD22	1:C:619:MET:HE1	1.97	0.47
1:B:484:LEU:HD22	1:B:497:PHE:HB2	2.01	0.41
1:A:708:GLN:HB2	4:A:913:HOH:O	2.20	0.41
1:C:464:SER:HB3	1:C:469:THR:HG23	2.03	0.41
1:A:658:ILE:CD1	1:A:673:LEU:HD11	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/275 (96%)	251 (95%)	12 (5%)	0	100	100
1	B	259/275 (94%)	250 (96%)	9 (4%)	0	100	100
1	C	260/275 (94%)	251 (96%)	9 (4%)	0	100	100
All	All	782/825 (95%)	752 (96%)	30 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/242 (99%)	238 (100%)	1 (0%)	84	90
1	B	231/242 (96%)	229 (99%)	2 (1%)	70	85
1	C	234/242 (97%)	233 (100%)	1 (0%)	84	90
All	All	704/726 (97%)	700 (99%)	4 (1%)	78	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	523	GLN
1	B	523	GLN
1	B	626	MET
1	C	720	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	511	ASN
1	A	538	HIS
1	A	539	HIS
1	A	561	GLN
1	B	455	GLN
1	B	539	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	652	GLN
1	B	657	ASN
1	C	511	ASN
1	C	538	HIS
1	C	539	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 7 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	WJ9	B	801	-	40,40,40	0.83	0	56,57,57	1.79	8 (14%)
2	WJ9	C	801	-	40,40,40	0.93	1 (2%)	56,57,57	1.83	11 (19%)
2	WJ9	A	801	-	40,40,40	0.93	1 (2%)	56,57,57	1.89	12 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	WJ9	B	801	-	-	4/27/44/44	0/4/4/4
2	WJ9	C	801	-	-	11/27/44/44	0/4/4/4
2	WJ9	A	801	-	-	6/27/44/44	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	WJ9	C4-C2	2.71	1.56	1.51
2	C	801	WJ9	C4-C2	2.59	1.56	1.51

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	WJ9	C7-N12-C11	7.29	125.11	117.62
2	C	801	WJ9	C7-N12-C11	6.32	124.11	117.62
2	B	801	WJ9	C7-N12-C11	6.11	123.90	117.62
2	B	801	WJ9	C30-N29-C34	-4.79	106.79	112.38
2	B	801	WJ9	C31-C30-N29	4.73	109.01	102.93
2	C	801	WJ9	C30-N29-C34	-4.67	106.92	112.38
2	A	801	WJ9	N26-C27-N29	4.66	121.38	115.91
2	A	801	WJ9	C31-C30-N29	4.66	108.92	102.93
2	C	801	WJ9	N26-C27-N29	4.38	121.05	115.91
2	C	801	WJ9	C31-C30-N29	4.35	108.52	102.93
2	B	801	WJ9	N26-C27-N29	3.53	120.06	115.91
2	A	801	WJ9	C30-N29-C34	-3.36	108.45	112.38
2	B	801	WJ9	C9-C10-C11	3.36	119.35	118.17
2	A	801	WJ9	O28-C27-N29	-3.23	117.16	121.67
2	A	801	WJ9	C10-C11-N13	2.90	125.86	122.35
2	C	801	WJ9	O28-C27-N29	-2.84	117.70	121.67
2	A	801	WJ9	C9-C10-C11	2.80	119.15	118.17
2	A	801	WJ9	C10-C11-N12	-2.55	118.42	123.34
2	C	801	WJ9	C9-C10-C11	2.48	119.04	118.17
2	B	801	WJ9	C10-C11-N12	-2.48	118.56	123.34
2	B	801	WJ9	N6-C7-N12	2.44	120.56	117.03
2	C	801	WJ9	C4-C2-N6	2.37	113.18	109.57
2	B	801	WJ9	C7-N6-C2	-2.29	120.40	123.98
2	A	801	WJ9	F37-C36-C35	-2.28	106.77	112.89
2	C	801	WJ9	C18-N13-C11	-2.26	114.40	120.41
2	A	801	WJ9	C7-N6-C2	-2.16	120.61	123.98
2	A	801	WJ9	C35-C32-C31	-2.14	107.37	115.45
2	C	801	WJ9	N6-C7-N12	2.13	120.11	117.03
2	C	801	WJ9	C10-C11-N12	-2.06	119.37	123.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	WJ9	C4-C2-N6	2.04	112.68	109.57
2	C	801	WJ9	C36-C35-C32	-2.01	111.34	115.32

There are no chirality outliers.

All (21) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	WJ9	C1-C2-C4-O5
2	A	801	WJ9	N6-C2-C4-O5
2	B	801	WJ9	C1-C2-C4-O5
2	C	801	WJ9	C1-C2-C4-O5
2	C	801	WJ9	N6-C2-C4-O5
2	C	801	WJ9	C1-C2-N6-C7
2	C	801	WJ9	C4-C2-N6-C7
2	C	801	WJ9	N12-C11-N13-C18
2	C	801	WJ9	C10-C11-N13-C18
2	A	801	WJ9	C10-C11-N13-C18
2	A	801	WJ9	N12-C11-N13-C18
2	B	801	WJ9	C10-C11-N13-C18
2	B	801	WJ9	N12-C11-N13-C18
2	B	801	WJ9	N6-C2-C4-O5
2	C	801	WJ9	C24-C19-C9-C8
2	A	801	WJ9	C22-C21-N26-C27
2	C	801	WJ9	C20-C19-C9-C8
2	C	801	WJ9	C24-C19-C9-C10
2	C	801	WJ9	C22-C21-N26-C27
2	A	801	WJ9	C20-C21-N26-C27
2	C	801	WJ9	C20-C21-N26-C27


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	269/275 (97%)	0.80	30 (11%) 10 7	41, 61, 102, 135	22 (8%)
1	B	262/275 (95%)	0.52	22 (8%) 17 10	29, 47, 85, 124	12 (4%)
1	C	266/275 (96%)	0.76	34 (12%) 7 5	36, 59, 107, 148	21 (7%)
All	All	797/825 (96%)	0.69	86 (10%) 11 7	29, 57, 102, 148	55 (6%)

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	628	ASP	6.6
1	C	628	ASP	6.4
1	A	486	VAL	5.7
1	B	612	LEU	5.6
1	B	598	THR	5.6
1	A	599	VAL	5.3
1	B	627	GLN	5.2
1	B	599	VAL	4.9
1	C	488	ALA	4.3
1	A	627	GLN	4.3
1	B	720	LEU	4.2
1	C	586	ASP	4.1
1	C	610	GLU	3.9
1	A	484	LEU	3.9
1	B	485	ASN	3.9
1	B	629	LYS	3.7
1	C	600	LYS	3.7
1	A	522	PRO	3.7
1	A	720	LEU	3.6
1	C	523	GLN	3.6
1	A	467	PHE	3.6
1	C	602	ARG	3.5
1	B	492	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	485	ASN	3.4
1	A	598	THR	3.4
1	C	489	PRO	3.3
1	A	721	PRO	3.2
1	B	586	ASP	3.2
1	B	597	ALA	3.0
1	C	522	PRO	3.0
1	A	466	SER	3.0
1	C	720	LEU	3.0
1	C	715	LEU	3.0
1	B	613	SER	3.0
1	C	613	SER	3.0
1	C	719	SER	3.0
1	A	453	ASP	2.9
1	C	520	THR	2.9
1	C	492	GLN	2.9
1	A	465	GLY	2.9
1	A	489	PRO	2.9
1	A	600	LYS	2.9
1	A	492	GLN	2.9
1	C	601	SER	2.9
1	B	614	GLY	2.8
1	C	494	LEU	2.8
1	A	485	ASN	2.8
1	A	523	GLN	2.8
1	B	453	ASP	2.8
1	C	484	LEU	2.7
1	B	628	ASP	2.7
1	C	493	GLN	2.7
1	A	521	LYS	2.7
1	C	495	GLN	2.7
1	C	521	LYS	2.6
1	C	524	LEU	2.6
1	C	609	PHE	2.6
1	B	484	LEU	2.6
1	A	605	GLY	2.6
1	B	447	ASP	2.5
1	C	701	ASP	2.4
1	A	626	MET	2.4
1	C	550	ILE	2.4
1	C	447	ASP	2.3
1	B	486	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	701	ASP	2.3
1	C	490	THR	2.3
1	A	700	ARG	2.2
1	C	717	ALA	2.2
1	A	695	CYS	2.2
1	A	461	ARG	2.2
1	A	483	MET	2.2
1	C	467	PHE	2.2
1	B	718	ARG	2.2
1	C	491	PRO	2.2
1	C	461	ARG	2.1
1	A	669	GLY	2.1
1	A	497	PHE	2.1
1	C	714	GLU	2.1
1	B	719	SER	2.1
1	B	717	ALA	2.1
1	A	464	SER	2.0
1	A	683	ASN	2.0
1	C	587	LEU	2.0
1	A	686	LYS	2.0
1	B	615	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	A	802	1/1	0.90	0.07	83,83,83,83	0
2	WJ9	A	801	37/37	0.93	0.11	40,42,47,47	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	C	802	1/1	0.94	0.28	67,67,67,67	0
3	CL	C	803	1/1	0.94	0.11	61,61,61,61	0
2	WJ9	C	801	37/37	0.95	0.10	44,48,54,55	0
3	CL	B	802	1/1	0.95	0.14	62,62,62,62	0
3	CL	A	803	1/1	0.96	0.13	67,67,67,67	0
3	CL	A	804	1/1	0.97	0.15	66,66,66,66	0
2	WJ9	B	801	37/37	0.97	0.08	26,27,37,43	0
3	CL	B	803	1/1	0.99	0.06	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around WJ9 A 801:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around WJ9 C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around WJ9 B 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.