



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 07:57 AM UTC

PDB ID : 9QQT / pdb_00009qqt
Title : Methyl-coenzyme M reductase of ANME-2d Candidatus Methanoperedens
Vercelli Strain 1 from a bioreactor enrichment culture
Authors : Mueller, M.-C.; Wagner, T.
Deposited on : 2025-04-02
Resolution : 0.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1146 (1.02-0.94)
Clashscore	190562	1197 (1.02-0.94)
Ramachandran outliers	187476	1136 (1.02-0.94)
Sidechain outliers	187428	1137 (1.02-0.94)
RSRZ outliers	180081	1144 (1.02-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	566	98%
2	B	434	97%
2	E	434	98%
3	C	249	98%
3	F	249	98%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	566	98% ..

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 43046 atoms, of which 19722 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methyl-coenzyme M reductase subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	564	Total	C	H	N	O	S	0	29	0
			8801	2828	4335	761	847	30			

- Molecule 2 is a protein called Methyl-coenzyme M reductase subunit beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	433	Total	C	H	N	O	S	0	43	0
			6868	2131	3499	575	644	19			
2	E	433	Total	C	H	N	O	S	0	35	0
			6759	2101	3434	567	639	18			

- Molecule 3 is a protein called Methyl-coenzyme M reductase subunit gamma.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	248	Total	C	H	N	O	S	0	15	0
			4053	1281	2001	366	392	13			
3	F	248	Total	C	H	N	O	S	0	13	0
			4016	1270	1980	366	387	13			

- Molecule 4 is a protein called Methyl-coenzyme M reductase subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	563	Total	C	H	N	O	S	0	31	0
			8799	2825	4341	757	845	31			

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	K	0	0
			1	1		

- Molecule 6 is FACTOR 430 (CCD ID: F43) (formula: C₄₂H₅₁N₆NiO₁₃) (labeled as "Ligand

F43

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 62	C 42	N 6	Ni 1	O 13	0	0
6	D	1	Total 62	C 42	N 6	Ni 1	O 13	0	0

-
- EDO
- Chemical structure of EDO (Ethane-1,2-diol) showing a zigzag conformation. The carbon atoms are labeled C1 and C2 in green. The hydroxyl groups are labeled HO and OH in red, with their respective oxygen atoms labeled O1 and O2 in green.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		



Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	C	1	Total	C	H	O	0	0
			10	2	6	2		
7	C	1	Total	C	H	O	0	0
			10	2	6	2		
7	D	1	Total	C	H	O	0	0
			10	2	6	2		
7	D	1	Total	C	H	O	0	0
			10	2	6	2		
7	D	1	Total	C	H	O	0	0
			10	2	6	2		
7	D	1	Total	C	H	O	0	0
			10	2	6	2		
7	D	1	Total	C	H	O	0	0
			10	2	6	2		
7	D	1	Total	C	H	O	0	0
			10	2	6	2		
7	E	1	Total	C	H	O	0	0
			10	2	6	2		
7	E	1	Total	C	H	O	0	0
			10	2	6	2		
7	F	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 8 is Coenzyme B (CCD ID: TP7) (formula: $C_{11}H_{22}NO_7PS$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total	C	N	O	P	S	0	1
			21	11	1	7	1	1		
8	D	1	Total	C	N	O	P	S	0	1
			21	11	1	7	1	1		

- Molecule 9 is 1-THIOETHANESULFONIC ACID (CCD ID: COM) (formula: $C_2H_6O_3S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	O	S	0	1
			7	2	3	2		
9	D	1	Total	C	O	S	0	1
			7	2	3	2		

- Molecule 10 is O-PHOSPHONO-N-{(2E)-7-[(2-SULFOETHYL)DITHIO]HEPT-2-ENOYL}-L-THREONINE (CCD ID: SHT) (formula: $C_{13}H_{24}NO_{10}PS_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
10	A	1	Total 28	C 13	N 1	O 10	P 1	S 3	0	1
10	D	1	Total 28	C 13	N 1	O 10	P 1	S 3	0	1

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	699	Total	O	0	71
			699	699		
11	B	587	Total	O	0	96
			587	587		
11	C	391	Total	O	0	36
			391	391		
11	D	687	Total	O	0	76
			687	687		
11	E	582	Total	O	0	98
			582	582		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	F	347	Total 347	O 347	0	18

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methyl-coenzyme M reductase subunit alpha

Chain A:  98%





- Molecule 2: Methyl-coenzyme M reductase subunit beta

Chain B:  97%





- Molecule 2: Methyl-coenzyme M reductase subunit beta

Chain E:  98%





- Molecule 3: Methyl-coenzyme M reductase subunit gamma

Chain C:  98%




- Molecule 3: Methyl-coenzyme M reductase subunit gamma

Chain F:  98%



- Molecule 4: Methyl-coenzyme M reductase subunit alpha

Chain D:  98% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.62Å 189.82Å 83.93Å 90.00° 114.36° 90.00°	Depositor
Resolution (Å)	41.71 – 0.98 41.71 – 0.98	Depositor EDS
% Data completeness (in resolution range)	74.0 (41.71-0.98) 75.3 (41.71-0.98)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 0.98Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.128 , 0.148 (Not available) , 0.133	Depositor DCC
R_{free} test set	49938 reflections (3.74%)	wwPDB-VP
Wilson B-factor (Å ²)	6.6	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for l,-k,h	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	43046	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRX, EDO, TP7, CSO, GL3, A1I9G, K, SMC, AGM, F43, COM, MHS, SHT, DYA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	4/4607 (0.1%)	0.77	4/6230 (0.1%)
2	B	0.52	0/3573	0.71	2/4834 (0.0%)
2	E	0.54	2/3487 (0.1%)	0.72	0/4721
3	C	0.54	0/2131	0.72	0/2876
3	F	0.50	0/2125	0.72	0/2868
4	D	0.53	0/4600	0.76	0/6217
All	All	0.54	6/20523 (0.0%)	0.74	6/27746 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	3
2	E	0	2
3	C	0	1
3	F	0	1
4	D	0	1
All	All	0	9

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	78[A]	PRO	C-O	7.19	1.32	1.23
1	A	78[B]	PRO	C-O	7.19	1.32	1.23
1	A	160[A]	ILE	C-O	5.84	1.31	1.24
1	A	160[B]	ILE	C-O	5.84	1.31	1.24
2	E	193[B]	MET	C-O	5.40	1.30	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160[A]	ILE	CA-C-N	-5.94	116.90	122.66
1	A	160[A]	ILE	C-N-CA	-5.94	116.90	122.66
1	A	160[B]	ILE	CA-C-N	-5.94	116.90	122.66
1	A	160[B]	ILE	C-N-CA	-5.94	116.90	122.66
2	B	127[A]	SER	CA-C-O	5.36	126.23	120.55

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	456	ARG	Sidechain
2	B	191[A]	ASN	Mainchain
2	B	191[B]	ASN	Mainchain
2	B	331	TYR	Sidechain
3	C	121	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4466	4335	4292	3	0
2	B	3369	3499	3404	6	0
2	E	3325	3434	3366	5	0
3	C	2052	2001	1977	2	0
3	F	2036	1980	1947	2	0
4	D	4458	4341	4299	4	0
5	A	1	0	0	0	0
6	A	62	0	43	1	0
6	D	62	0	43	1	0
7	A	12	18	18	0	0
7	B	28	42	42	0	0
7	C	8	12	12	0	0
7	D	28	42	42	0	0
7	E	8	12	12	0	0
7	F	4	6	6	0	0
8	A	21	0	19	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	D	21	0	19	0	0
9	A	7	0	4	0	0
9	D	7	0	5	0	0
10	A	28	0	20	0	0
10	D	28	0	20	0	0
11	A	699	0	0	0	0
11	B	587	0	0	1	0
11	C	391	0	0	0	0
11	D	687	0	0	1	0
11	E	582	0	0	0	0
11	F	347	0	0	0	0
All	All	23324	19722	19590	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:124[B]:TYR:CE1	2:B:188:THR:HG21	2.21	0.76
2:E:124[B]:TYR:CE1	2:E:188:THR:HG21	2.26	0.70
4:D:234:ASP:CG	4:D:236:PRO:HD2	2.34	0.52
3:C:85:TYR:CE1	4:D:257:CYS:HB2	2.46	0.51
4:D:343:VAL:HB	6:D:603:F43:H9A1	1.93	0.50

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	584/566 (103%)	564 (97%)	19 (3%)	1 (0%)	43 16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	476/434 (110%)	466 (98%)	10 (2%)	0	100	100
2	E	466/434 (107%)	457 (98%)	9 (2%)	0	100	100
3	C	260/249 (104%)	254 (98%)	6 (2%)	0	100	100
3	F	258/249 (104%)	251 (97%)	7 (3%)	0	100	100
4	D	586/566 (104%)	566 (97%)	19 (3%)	1 (0%)	43	16
All	All	2630/2498 (105%)	2558 (97%)	70 (3%)	2 (0%)	48	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	340	SER
4	D	340	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	461/435 (106%)	461 (100%)	0	100	100
2	B	377/335 (112%)	373 (99%)	4 (1%)	65	32
2	E	367/335 (110%)	365 (100%)	2 (0%)	81	55
3	C	222/210 (106%)	222 (100%)	0	100	100
3	F	221/210 (105%)	221 (100%)	0	100	100
4	D	463/436 (106%)	462 (100%)	1 (0%)	87	63
All	All	2111/1961 (108%)	2104 (100%)	7 (0%)	87	62

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	179[B]	PRO
4	D	437	GLN
2	E	123[B]	GLU
2	E	123[A]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	179[A]	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	465	GLN
4	D	472	ASN
2	E	207	ASN
4	D	493	ASN
2	B	112	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

15 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MHS	D	272	4	10,11,12	0.66	0	5,14,16	0.85	0
3	A1I9G	C	159	3	9,11,12	1.53	1 (11%)	7,14,16	1.13	0
1	MHS	A	272	1	10,11,12	0.67	0	5,14,16	1.82	3 (60%)
4	SMC	D	468	4	5,6,7	0.65	0	3,6,8	1.11	0
3	A1I9G	F	159	3	9,11,12	1.20	1 (11%)	7,14,16	0.79	0
1	AGM	A	286	1	10,11,12	0.89	0	7,13,15	0.68	0
1	SMC	A	468	1	5,6,7	0.41	0	3,6,8	1.00	0
4	GL3	D	461	4	2,3,4	2.68	1 (50%)	1,2,4	0.49	0
4	DYA	D	466	4	6,7,8	0.99	0	6,8,10	2.07	3 (50%)
4	AGM	D	286	4	10,11,12	0.81	0	7,13,15	0.72	0
4	TRX	D	443	4	15,16,17	0.67	0	17,22,24	0.75	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TRX	A	443	1	15,16,17	0.66	0	17,22,24	0.60	0
1	GL3	A	461	1	2,3,4	1.84	1 (50%)	1,2,4	0.17	0
1	DYA	A	466	1	6,7,8	1.22	1 (16%)	6,8,10	2.40	3 (50%)
1	CSO	A	55	1	3,6,7	0.85	0	1,6,8	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MHS	D	272	4	-	0/5/6/8	0/1/1/1
3	A1I9G	C	159	3	-	2/9/10/12	0/1/1/1
1	MHS	A	272	1	-	0/5/6/8	0/1/1/1
4	SMC	D	468	4	-	1/3/5/7	-
3	A1I9G	F	159	3	-	2/9/10/12	0/1/1/1
1	AGM	A	286	1	-	2/10/11/13	-
1	SMC	A	468	1	-	1/3/5/7	-
4	GL3	D	461	4	-	0/1/1/2	-
4	DYA	D	466	4	-	3/4/6/8	-
4	AGM	D	286	4	-	2/10/11/13	-
4	TRX	D	443	4	-	0/5/6/8	0/2/2/2
1	TRX	A	443	1	-	0/5/6/8	0/2/2/2
1	GL3	A	461	1	-	1/1/1/2	-
1	DYA	A	466	1	-	3/4/6/8	-
1	CSO	A	55	1	-	0/1/5/7	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	159	A1I9G	CG-ND1	-4.05	1.33	1.37
4	D	461	GL3	C-S	-3.79	1.65	1.80
3	F	159	A1I9G	CG-ND1	-2.79	1.34	1.37
1	A	461	GL3	C-S	-2.60	1.69	1.80
1	A	466	DYA	OD1-CG	-2.00	1.25	1.30

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	466	DYA	O-C-CA	-3.66	120.80	125.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	466	DYA	CB-CA-N	-3.34	117.19	123.25
4	D	466	DYA	O-C-CA	-3.09	121.52	125.39
1	A	272	MHS	CM-ND1-CG	-2.78	123.74	126.69
4	D	466	DYA	CB-CA-N	-2.70	118.34	123.25

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	466	DYA	O-C-CA-CB
1	A	466	DYA	CA-CB-CG-OD1
1	A	468	SMC	CA-CB-SG-CS
4	D	466	DYA	O-C-CA-CB
4	D	466	DYA	CA-CB-CG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 1 is monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	TP7	D	604[A]	-	19,20,20	0.97	0	24,26,26	0.64	0
7	EDO	B	503	-	3,3,3	0.48	0	2,2,2	0.42	0
7	EDO	A	606	-	3,3,3	0.16	0	2,2,2	0.08	0
9	COM	D	601[A]	6	6,6,6	1.20	1 (16%)	8,8,8	1.72	3 (37%)
7	EDO	D	605	-	3,3,3	0.50	0	2,2,2	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	E	501	-	3,3,3	0.50	0	2,2,2	0.26	0
7	EDO	B	507	-	3,3,3	0.08	0	2,2,2	0.19	0
9	COM	A	607[A]	6	6,6,6	2.93	1 (16%)	8,8,8	1.27	1 (12%)
7	EDO	B	504	-	3,3,3	0.16	0	2,2,2	0.71	0
7	EDO	D	607	-	3,3,3	0.48	0	2,2,2	0.36	0
7	EDO	E	502	-	3,3,3	0.13	0	2,2,2	0.22	0
7	EDO	B	502	-	3,3,3	0.56	0	2,2,2	0.27	0
7	EDO	D	609	-	3,3,3	0.07	0	2,2,2	0.11	0
7	EDO	C	301	-	3,3,3	0.85	0	2,2,2	0.49	0
10	SHT	A	608[C]	6	26,27,27	1.54	1 (3%)	32,36,36	1.56	4 (12%)
6	F43	D	603	10,1,9	63,71,71	1.92	5 (7%)	73,118,118	1.21	6 (8%)
7	EDO	B	506	-	3,3,3	0.07	0	2,2,2	0.18	0
7	EDO	D	608	-	3,3,3	0.14	0	2,2,2	0.43	0
7	EDO	F	301	-	3,3,3	0.36	0	2,2,2	0.48	0
7	EDO	D	602	-	3,3,3	0.30	0	2,2,2	0.26	0
7	EDO	A	603	-	3,3,3	0.31	0	2,2,2	0.34	0
7	EDO	A	605	-	3,3,3	0.50	0	2,2,2	0.39	0
7	EDO	C	302	-	3,3,3	0.41	0	2,2,2	0.52	0
8	TP7	A	604[A]	-	19,20,20	0.85	0	24,26,26	0.78	0
10	SHT	D	610[C]	6	26,27,27	1.53	1 (3%)	32,36,36	1.50	2 (6%)
6	F43	A	602	10,9,4	63,71,71	1.96	6 (9%)	73,118,118	1.17	7 (9%)
7	EDO	D	606	-	3,3,3	0.45	0	2,2,2	0.44	0
7	EDO	B	501	-	3,3,3	0.43	0	2,2,2	0.16	0
7	EDO	D	611	-	3,3,3	0.12	0	2,2,2	0.19	0
7	EDO	B	505	-	3,3,3	0.26	0	2,2,2	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	TP7	D	604[A]	-	-	1/24/24/24	-
7	EDO	B	503	-	-	0/1/1/1	-
7	EDO	A	606	-	-	0/1/1/1	-
9	COM	D	601[A]	6	-	0/4/4/4	-
7	EDO	D	605	-	-	0/1/1/1	-
7	EDO	E	501	-	-	0/1/1/1	-
7	EDO	B	507	-	-	1/1/1/1	-
9	COM	A	607[A]	6	-	0/4/4/4	-
7	EDO	B	504	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	D	607	-	-	0/1/1/1	-
7	EDO	E	502	-	-	1/1/1/1	-
7	EDO	B	502	-	-	0/1/1/1	-
7	EDO	D	609	-	-	0/1/1/1	-
7	EDO	C	301	-	-	0/1/1/1	-
10	SHT	A	608[C]	6	-	4/31/31/31	-
6	F43	D	603	10,1,9	-	8/28/185/185	-
7	EDO	B	506	-	-	0/1/1/1	-
7	EDO	D	608	-	-	0/1/1/1	-
7	EDO	F	301	-	-	0/1/1/1	-
7	EDO	D	602	-	-	0/1/1/1	-
7	EDO	A	603	-	-	0/1/1/1	-
7	EDO	A	605	-	-	0/1/1/1	-
7	EDO	C	302	-	-	0/1/1/1	-
8	TP7	A	604[A]	-	-	0/24/24/24	-
10	SHT	D	610[C]	6	-	4/31/31/31	-
6	F43	A	602	10,9,4	-	7/28/185/185	-
7	EDO	D	606	-	-	0/1/1/1	-
7	EDO	B	501	-	-	0/1/1/1	-
7	EDO	D	611	-	-	1/1/1/1	-
7	EDO	B	505	-	-	0/1/1/1	-

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	602	F43	NI-NA	8.02	2.08	1.89
6	A	602	F43	NI-NB	7.63	2.07	1.89
6	D	603	F43	NI-NA	7.60	2.07	1.89
6	D	603	F43	NI-NB	7.35	2.07	1.89
10	A	608[C]	SHT	CK1-CK2	7.13	1.51	1.32

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	608[C]	SHT	CK3-CK2-CK1	-6.03	113.35	125.92
10	D	610[C]	SHT	CK3-CK2-CK1	-5.95	113.52	125.92
10	D	610[C]	SHT	CK2-CK1-CK	-4.11	112.48	122.94
10	A	608[C]	SHT	CK2-CK1-CK	-4.00	112.77	122.94
6	D	603	F43	O8D-C7D-C6D	-3.22	115.59	120.87

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	A	608[C]	SHT	S1-CC-CD-SG2
10	D	610[C]	SHT	S1-CC-CD-SG2
7	B	507	EDO	O1-C1-C2-O2
10	D	610[C]	SHT	CK1-CK2-CK3-CK4
10	A	608[C]	SHT	CK1-CK2-CK3-CK4


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	603	F43	1	0
6	A	602	F43	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand TP7 D 604 (A)




Ligand COM D 601 (A)







Ligand F43 D 603




Bond lengths



Bond angles




Torsions





Rings

Ligand TP7 A 604 (A)



Ligand SHT D 610 (C)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	557/566 (98%)	-0.69	3 (0%) 87 88	3, 8, 16, 30	20 (3%)
2	B	433/434 (99%)	-0.74	2 (0%) 87 88	3, 7, 14, 22	30 (6%)
2	E	433/434 (99%)	-0.68	0 100 100	3, 8, 15, 23	26 (6%)
3	C	247/249 (99%)	-0.68	1 (0%) 88 89	5, 8, 15, 24	12 (4%)
3	F	247/249 (99%)	-0.42	3 (1%) 76 81	6, 11, 19, 28	9 (3%)
4	D	557/566 (98%)	-0.77	1 (0%) 91 90	3, 7, 15, 26	22 (3%)
All	All	2474/2498 (99%)	-0.68	10 (0%) 88 89	3, 8, 16, 30	119 (4%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	7.7
3	F	2	ALA	6.0
3	C	2	ALA	4.2
3	F	60	ILE	3.5
4	D	565	ALA	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MHS	D	272	11/12	0.98	0.06	4,5,10,12	0
1	MHS	A	272	11/12	0.99	0.05	5,6,10,12	0
1	AGM	A	286	12/13	0.99	0.04	3,4,5,6	0
1	TRX	A	443	15/16	0.99	0.04	4,5,5,6	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	DYA	A	466	8/9	0.99	0.03	4,5,6,6	0
1	CSO	A	55	7/8	0.99	0.05	7,7,8,13	0
4	AGM	D	286	12/13	0.99	0.04	4,4,5,6	0
4	TRX	D	443	15/16	0.99	0.03	4,4,5,5	0
4	DYA	D	466	8/9	0.99	0.03	4,5,6,6	0
3	A1I9G	C	159	11/12	0.99	0.04	5,5,6,6	0
3	A1I9G	F	159	11/12	0.99	0.04	6,6,7,7	0
1	SMC	A	468	7/8	1.00	0.04	4,4,6,6	0
4	SMC	D	468	7/8	1.00	0.03	4,4,6,7	0
1	GL3	A	461	4/5	1.00	0.03	4,4,4,4	0
4	GL3	D	461	4/5	1.00	0.04	4,4,4,5	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	EDO	E	502	4/4	0.75	0.17	23,32,43,44	0
7	EDO	B	507	4/4	0.80	0.15	26,31,36,43	0
7	EDO	D	611	4/4	0.81	0.17	18,24,29,29	10
7	EDO	B	504	4/4	0.82	0.17	14,20,24,29	0
7	EDO	B	506	4/4	0.83	0.12	28,34,35,36	0
7	EDO	D	607	4/4	0.85	0.12	29,35,36,37	0
7	EDO	D	609	4/4	0.85	0.13	26,32,32,33	0
7	EDO	B	503	4/4	0.88	0.12	21,26,28,30	0
7	EDO	B	505	4/4	0.91	0.12	12,21,25,27	0
7	EDO	A	605	4/4	0.92	0.10	20,24,26,27	0
7	EDO	D	608	4/4	0.93	0.10	21,25,28,29	0
7	EDO	D	605	4/4	0.93	0.09	20,24,26,27	0
7	EDO	B	501	4/4	0.94	0.09	16,22,25,27	0
7	EDO	D	606	4/4	0.95	0.08	14,17,18,18	0
7	EDO	A	606	4/4	0.96	0.07	16,20,23,23	0
7	EDO	E	501	4/4	0.96	0.08	13,16,16,17	0
7	EDO	B	502	4/4	0.96	0.07	12,14,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	EDO	C	301	4/4	0.97	0.06	12,14,16,16	0
10	SHT	A	608[C]	28/28	0.97	0.06	5,5,9,10	28
7	EDO	F	301	4/4	0.98	0.05	14,17,18,21	0
7	EDO	C	302	4/4	0.98	0.04	13,16,18,21	0
10	SHT	D	610[C]	28/28	0.98	0.06	4,5,8,8	28
7	EDO	D	602	4/4	0.99	0.04	8,10,12,12	0
7	EDO	A	603	4/4	0.99	0.04	7,9,11,11	0
9	COM	A	607[A]	7/7	0.99	0.07	3,4,6,6	7
9	COM	D	601[A]	7/7	0.99	0.06	5,6,8,8	7
6	F43	A	602	62/62	0.99	0.04	4,5,8,10	0
6	F43	D	603	62/62	0.99	0.05	5,6,10,11	0
8	TP7	A	604[A]	21/21	1.00	0.03	2,3,4,6	21
8	TP7	D	604[A]	21/21	1.00	0.03	3,4,5,6	21
5	K	A	601	1/1	1.00	0.04	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.