



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:53 PM UTC

PDB ID : 1QR3 / pdb_00001qr3
Title : Structure of porcine pancreatic elastase in complex with FR901277, a novel macrocyclic inhibitor of elastases at 1.6 angstrom resolution
Authors : Nakanishi, I.; Kinoshita, T.; Sato, A.; Tada, T.
Deposited on : 1999-06-18
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	240	
2	I	8	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chymotrypsin-like elastase family member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	240	Total	C	N	O	S	0	0	0
			1822	1135	330	347	10			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	77	ASN	ASP	variant	UNP P00772

- Molecule 2 is a protein called FR901277 Inhibitor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	8	Total	C	N	O	0	0	0
			69	47	9	13			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	1	Total	Ca	0	0
			1	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	333	Total	O	0	0
			333	333		
5	I	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chymotrypsin-like elastase family member 1



- Molecule 2: FR901277 Inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.67Å 57.84Å 75.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.60 6.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-1.60) 33.1 (6.00-1.60)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 1.58Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.197 , (Not available) 0.200 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.387	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2234	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLJ, ALQ, CIR, SO4, TYJ, CA, DBU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	1.18	7/1862 (0.4%)	2.02	55/2543 (2.2%)
2	I	1.04	0/23	1.45	0/27
All	All	1.18	7/1885 (0.4%)	2.02	55/2570 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	7

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	200	HIS	CE1-NE2	6.74	1.39	1.32
1	E	200	HIS	ND1-CE1	5.69	1.38	1.32
1	E	91	HIS	ND1-CE1	5.29	1.37	1.32
1	E	65(A)	ARG	NE-CZ	5.20	1.38	1.33
1	E	57	HIS	CG-CD2	5.14	1.41	1.35
1	E	91	HIS	CE1-NE2	5.05	1.37	1.32
1	E	71	HIS	CE1-NE2	5.01	1.37	1.32

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	241	VAL	N-CA-C	9.17	119.15	110.53
1	E	65(A)	ARG	NE-CZ-NH2	8.96	127.27	119.20
1	E	37	SER	N-CA-C	7.79	118.55	108.34
1	E	24	ARG	CD-NE-CZ	-7.56	113.82	124.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	42	CYS	CA-C-N	-7.44	113.05	121.83
1	E	42	CYS	C-N-CA	-7.44	113.05	121.83
1	E	204	ASN	CA-CB-CG	-7.41	105.19	112.60
1	E	230	ARG	NE-CZ-NH2	-7.21	112.71	119.20
1	E	24	ARG	CB-CA-C	-7.15	98.53	110.68
1	E	99	VAL	CB-CA-C	-6.76	103.01	112.14
1	E	141	TRP	CA-C-N	-6.70	112.71	121.96
1	E	141	TRP	C-N-CA	-6.70	112.71	121.96
1	E	96	THR	N-CA-C	6.65	120.50	112.38
1	E	74	ASN	CA-CB-CG	-6.58	106.02	112.60
1	E	77	ASN	CA-C-N	-6.37	103.99	122.46
1	E	77	ASN	C-N-CA	-6.37	103.99	122.46
1	E	118	VAL	CB-CA-C	6.32	118.71	110.12
1	E	223	ARG	CA-C-N	-6.26	115.00	123.46
1	E	223	ARG	C-N-CA	-6.26	115.00	123.46
1	E	34	GLN	CA-C-N	-6.25	113.00	122.81
1	E	34	GLN	C-N-CA	-6.25	113.00	122.81
1	E	230	ARG	CD-NE-CZ	6.23	133.13	124.40
1	E	170(A)	SER	N-CA-C	6.14	117.97	111.28
1	E	175	THR	CA-C-N	-6.12	111.36	122.43
1	E	175	THR	C-N-CA	-6.12	111.36	122.43
1	E	218	LEU	N-CA-C	5.99	119.84	112.54
1	E	116	SER	N-CA-C	5.93	120.12	113.01
1	E	191	CYS	CA-C-N	-5.89	112.51	120.87
1	E	191	CYS	C-N-CA	-5.89	112.51	120.87
1	E	230	ARG	NE-CZ-NH1	5.83	127.33	121.50
1	E	65(A)	ARG	NE-CZ-NH1	-5.72	115.78	121.50
1	E	240	ASN	N-CA-C	5.70	117.50	111.28
1	E	133	ASN	N-CA-C	5.67	119.50	112.24
1	E	188	VAL	CB-CA-C	-5.67	106.05	111.44
1	E	67	VAL	CA-C-O	5.65	126.33	120.39
1	E	81	GLN	CB-CG-CD	-5.63	103.02	112.60
1	E	166	ALA	N-CA-C	5.51	118.00	111.33
1	E	235	ILE	N-CA-C	5.47	116.20	110.62
1	E	67	VAL	N-CA-C	5.44	115.72	108.11
1	E	59	VAL	CA-CB-CG1	5.39	119.56	110.40
1	E	65(A)	ARG	CB-CG-CD	5.27	123.42	111.30
1	E	109	ALA	CA-C-O	5.26	126.02	119.97
1	E	85	VAL	CA-C-N	-5.25	112.84	121.92
1	E	85	VAL	C-N-CA	-5.25	112.84	121.92
1	E	220	CYS	N-CA-C	5.21	116.82	111.03
1	E	91	HIS	CA-CB-CG	-5.14	108.66	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	240	ASN	CA-C-N	-5.13	113.99	120.56
1	E	240	ASN	C-N-CA	-5.13	113.99	120.56
1	E	206	GLN	CA-C-N	-5.11	114.19	121.50
1	E	206	GLN	C-N-CA	-5.11	114.19	121.50
1	E	148	ASN	CA-CB-CG	-5.08	107.52	112.60
1	E	75	GLN	CA-C-N	-5.06	114.38	121.72
1	E	75	GLN	C-N-CA	-5.06	114.38	121.72
1	E	223	ARG	N-CA-C	5.03	121.52	110.80
1	E	143	LEU	N-CA-C	5.00	117.59	110.24

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	137	TYR	Sidechain
1	E	189	SER	Peptide
1	E	24	ARG	Sidechain
1	E	60	ASP	Peptide
1	E	65(A)	ARG	Sidechain
1	E	78	GLY	Peptide
1	E	93	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1822	0	1759	31	0
2	I	69	0	55	1	0
3	E	1	0	0	0	0
4	E	5	0	0	1	0
5	E	333	0	0	13	0
5	I	4	0	0	0	0
All	All	2234	0	1814	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:221(A):VAL:HG13	1:E:224:LYS:HB2	1.74	0.69
1:E:68:VAL:HG12	1:E:81:GLN:HB2	1.77	0.66
5:E:610:HOH:O	2:I:6:PHE:HE1	1.79	0.64
1:E:59:VAL:HG12	5:E:623:HOH:O	1.99	0.62
1:E:91:HIS:CD2	1:E:93:TYR:H	2.18	0.61
1:E:216:VAL:HA	5:E:425:HOH:O	1.99	0.61
1:E:91:HIS:HD2	1:E:93:TYR:H	1.52	0.58
1:E:38:TRP:CE3	1:E:65(A):ARG:HG2	2.41	0.56
1:E:83:VAL:HG13	1:E:110:GLN:HG2	1.87	0.56
1:E:20:THR:HG23	5:E:424:HOH:O	2.08	0.54
1:E:127:GLY:HA2	4:E:401:SO4:O1	2.11	0.51
1:E:86:GLN:HG2	1:E:109:ALA:HA	1.94	0.50
1:E:112:VAL:HG13	5:E:497:HOH:O	2.10	0.49
1:E:25:ASN:HA	5:E:457:HOH:O	2.12	0.49
1:E:195:SER:HA	1:E:213:THR:HB	1.95	0.49
1:E:70:GLU:HB3	5:E:541:HOH:O	2.13	0.48
1:E:80:GLU:HG3	1:E:82:TYR:OH	2.14	0.47
1:E:118:VAL:HG22	5:E:732:HOH:O	2.15	0.46
1:E:221(A):VAL:CG1	1:E:224:LYS:HB2	2.45	0.45
1:E:237:TRP:O	1:E:241:VAL:HG13	2.17	0.45
1:E:114:LEU:HD12	5:E:732:HOH:O	2.18	0.44
1:E:52:VAL:HG23	1:E:108:LEU:HD11	1.99	0.43
1:E:227:VAL:HG12	5:E:407:HOH:O	2.17	0.43
1:E:67:VAL:HG21	5:E:463:HOH:O	2.17	0.43
1:E:46:LEU:C	1:E:46:LEU:HD13	2.44	0.43
1:E:143:LEU:HD23	1:E:151:LEU:HD23	2.01	0.42
1:E:27:TRP:HE1	1:E:157:GLN:NE2	2.16	0.42
1:E:24:ARG:HH11	1:E:24:ARG:HD2	1.60	0.41
1:E:38:TRP:CZ3	1:E:65(A):ARG:HG2	2.56	0.41
1:E:91:HIS:HD2	1:E:93:TYR:N	2.19	0.40
1:E:30:GLN:NE2	5:E:665:HOH:O	2.53	0.40
1:E:74:ASN:HB2	5:E:656:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	238/240 (99%)	229 (96%)	9 (4%)	0	100	100
2	I	2/8 (25%)	2 (100%)	0	0	100	100
All	All	240/248 (97%)	231 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	198/198 (100%)	173 (87%)	25 (13%)	4	0
2	I	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	201/201 (100%)	175 (87%)	26 (13%)	4	0

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	36	ARG
1	E	36(A)	SER
1	E	36(C)	SER
1	E	46	LEU
1	E	47	ILE
1	E	48	ARG
1	E	61	ARG
1	E	62	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	63	LEU
1	E	64	THR
1	E	67	VAL
1	E	68	VAL
1	E	70	GLU
1	E	80	GLU
1	E	99	VAL
1	E	110	GLN
1	E	118	VAL
1	E	122	VAL
1	E	163	VAL
1	E	171(B)	SER
1	E	171	TYR
1	E	174	SER
1	E	203	VAL
1	E	241	VAL
1	E	245	ASN
2	I	8	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	91	HIS
1	E	156	GLN
1	E	157	GLN
1	E	240	ASN
1	E	245	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DBU	I	4	2	3,5,6	1.68	1 (33%)	3,5,7	3.19	2 (66%)
2	TYJ	I	7	2	13,14,16	0.91	1 (7%)	17,18,22	1.85	4 (23%)
2	CIR	I	2	2	9,10,11	0.63	0	6,11,13	2.36	2 (33%)
2	GLJ	I	5	2	6,7,9	0.60	0	2,7,11	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DBU	I	4	2	-	1/1/4/6	-
2	TYJ	I	7	2	-	1/5/8/10	0/1/1/1
2	CIR	I	2	2	-	2/8/9/11	-
2	GLJ	I	5	2	-	1/5/6/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	4	DBU	C-CA	2.75	1.49	1.45
2	I	7	TYJ	OH1-CZ1	2.33	1.41	1.36

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	4	DBU	O-C-CA	-5.01	119.11	125.39
2	I	2	CIR	C5-N6-C7	-4.55	116.70	122.64
2	I	7	TYJ	CE21-CD21-CG5	-4.13	116.53	120.81
2	I	7	TYJ	CG5-CB7-CA	-2.93	109.34	113.51
2	I	7	TYJ	CD12-CG5-CD21	2.85	122.48	118.55
2	I	7	TYJ	CD12-CE11-CZ1	-2.42	118.08	120.50
2	I	2	CIR	O7-C7-N6	2.39	125.46	121.69
2	I	4	DBU	CG-CB-CA	-2.16	123.48	126.12

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	5	GLJ	O-C-CA-CB
2	I	7	TYJ	C-CA-CB7-CG5
2	I	2	CIR	C4-C3-CA-C
2	I	4	DBU	O-C-CA-CB
2	I	2	CIR	C4-C5-N6-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	E	401	-	4,4,4	0.38	0	6,6,6	0.25	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	401	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	240/240 (100%)	-0.35	0 100 100	11, 19, 32, 49	0
2	I	3/8 (37%)	0.28	0 100 100	18, 18, 21, 23	0
All	All	243/248 (97%)	-0.34	0 100 100	11, 19, 32, 49	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CIR	I	2	11/12	0.90	0.07	20,21,24,25	0
2	GLJ	I	5	8/10	0.92	0.07	16,16,17,17	0
2	TYJ	I	7	14/16	0.92	0.07	22,26,31,32	0
2	DBU	I	4	6/7	0.96	0.04	14,16,17,17	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	E	400	1/1	0.72	0.08	72,72,72,72	0
4	SO4	E	401	5/5	0.89	0.13	40,41,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.