



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:36 PM UTC

PDB ID : 1QRI / pdb_00001qri
Title : X-RAY STRUCTURE OF THE DNA-ECO RI ENDONUCLEASE COM-
PLEXES WITH AN E144D MUTATION AT 2.7 Å
Authors : Choi, J.; Kim, Y.; Greene, P.; Hager, P.; Rosenberg, J.M.
Deposited on : 1999-06-14
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	M	13	
2	A	261	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2874 atoms, of which 509 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*TP*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	M	13	Total	C	H	N	O	P	0	0	0
			292	126	29	48	77	12			

- Molecule 2 is a protein called ECO RI ENDONUCLEASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	A	261	Total	C	H	N	O	S	0	0	0
			2530	1293	480	356	394	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	ASP	GLU	engineered mutation	UNP P00642

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	M	12	Total	O	0	0
			12	12		
3	A	40	Total	O	0	0
			40	40		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 5'-D(*TP*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain M: 



- Molecule 2: ECO RI ENDONUCLEASE

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.40 Å 118.40 Å 49.70 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.60	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.60)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.164 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2874	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	M	0.76	0/294	2.51	19/452 (4.2%)
2	A	0.38	0/2083	0.77	0/2804
All	All	0.44	0/2377	1.17	19/3256 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	1

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	2	DC	O3'-P-O5'	-17.29	78.07	104.00
1	M	12	DC	OP1-P-O3'	15.94	155.82	108.00
1	M	10	DC	O3'-P-O5'	-14.19	82.72	104.00
1	M	6	DA	OP1-P-O3'	13.71	149.13	108.00
1	M	12	DC	O3'-P-O5'	-12.64	85.05	104.00
1	M	12	DC	OP2-P-O3'	-11.73	72.81	108.00
1	M	6	DA	OP2-P-O3'	-11.08	74.77	108.00
1	M	6	DA	O3'-P-O5'	-9.85	89.22	104.00
1	M	10	DC	OP1-P-O3'	9.62	136.87	108.00
1	M	2	DC	OP1-P-O3'	8.17	132.51	108.00
1	M	8	DT	O3'-P-O5'	-6.86	93.71	104.00
1	M	3	DG	O4'-C1'-C2'	-5.95	97.48	106.40
1	M	4	DC	O3'-P-O5'	5.80	112.70	104.00
1	M	13	DG	C5'-C4'-C3'	-5.80	106.20	114.90
1	M	4	DC	C3'-C2'-C1'	5.71	110.17	101.60
1	M	5	DG	O4'-C4'-C3'	-5.68	96.87	105.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	12	DC	O4'-C1'-C2'	-5.39	98.31	106.40
1	M	8	DT	OP2-P-O3'	5.12	123.35	108.00
1	M	11	DG	O3'-P-O5'	-5.01	96.48	104.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	M	5	DG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	263	29	148	1	3
2	A	2050	480	2055	29	0
3	A	40	0	0	0	0
3	M	12	0	0	0	0
All	All	2365	509	2203	30	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:35:VAL:HG23	2:A:255:MET:HE1	1.64	0.78
2:A:119:ILE:HG23	2:A:181:ILE:HG12	1.75	0.69
2:A:166:VAL:HG22	2:A:258:ILE:HD12	1.79	0.65
2:A:111:GLU:HG2	2:A:113:LYS:HE2	1.87	0.57
2:A:178:ASN:HD21	2:A:192:GLU:HG2	1.70	0.57
2:A:52:GLN:HB2	2:A:277:LYS:HB3	1.87	0.56
2:A:157:MET:HE1	2:A:164:PRO:HD2	1.87	0.55
2:A:114:HIS:CD2	2:A:170:GLU:HB3	2.42	0.54
2:A:114:HIS:HD2	2:A:170:GLU:HB3	1.75	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:12:DC:H2"	1:M:13:DG:C8	2.48	0.48
2:A:179:ILE:HG13	2:A:191:LEU:HB2	1.95	0.48
2:A:67:ASN:HD22	2:A:78:GLY:HA2	1.79	0.47
2:A:157:MET:CE	2:A:164:PRO:HD2	2.44	0.47
2:A:106:VAL:HG13	2:A:157:MET:SD	2.55	0.46
2:A:166:VAL:CG2	2:A:258:ILE:HD12	2.44	0.46
2:A:209:TYR:CE1	2:A:232:LEU:HB3	2.51	0.45
2:A:67:ASN:O	2:A:71:LYS:HG2	2.17	0.45
2:A:179:ILE:HD11	2:A:191:LEU:HD12	1.98	0.45
2:A:211:MET:HB3	2:A:212:PRO:HD2	1.98	0.44
2:A:261:THR:O	2:A:265:VAL:HG23	2.17	0.44
2:A:27:TYR:O	2:A:31:HIS:HD2	2.02	0.43
2:A:119:ILE:O	2:A:123:ARG:HB2	2.19	0.43
2:A:222:PHE:CE1	2:A:231:MET:HB2	2.54	0.42
2:A:111:GLU:HB3	2:A:167:LEU:HD12	2.00	0.42
2:A:21:ILE:HD13	2:A:21:ILE:H	1.83	0.42
2:A:130:LYS:HE2	2:A:130:LYS:HB2	1.94	0.42
2:A:38:VAL:HG21	2:A:251:MET:HE3	2.01	0.42
2:A:91:ASP:OD1	2:A:111:GLU:HG3	2.20	0.42
2:A:74:ASP:HB3	2:A:77:LEU:HD22	2.01	0.41
2:A:35:VAL:HG23	2:A:255:MET:CE	2.42	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:5:DG:O6	1:M:10:DC:H42[4_555]	1.52	0.08
1:M:6:DA:H62	1:M:9:DT:O4[4_555]	1.55	0.05
1:M:3:DG:O6	1:M:12:DC:H42[4_555]	1.56	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	259/261 (99%)	242 (93%)	14 (5%)	3 (1%)	10	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	228	LYS
2	A	128	VAL
2	A	184	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	225/225 (100%)	202 (90%)	23 (10%)	7	15

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	20	VAL
2	A	21	ILE
2	A	32	ASP
2	A	44	LYS
2	A	68	GLU
2	A	71	LYS
2	A	81	LEU
2	A	105	ARG
2	A	106	VAL
2	A	123	ARG
2	A	131	ARG
2	A	158	LEU
2	A	159	SER
2	A	166	VAL
2	A	175	LEU
2	A	193	TYR
2	A	204	LEU
2	A	217	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	240	GLN
2	A	245	GLU
2	A	263	LEU
2	A	266	LEU
2	A	273	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	67	ASN
2	A	114	HIS
2	A	124	ASN
2	A	178	ASN
2	A	220	ASN
2	A	240	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.