



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 04:05 AM UTC

PDB ID : 2QRC / pdb_00002qrc
Title : Crystal structure of the adenylate sensor from AMP-activated protein kinase
in complex with ADP and AMP
Authors : Jin, X.; Townley, R.; Shapiro, L.
Deposited on : 2007-07-28
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	137	% 58%22%15%
1	C	137	% 52%28%18%
2	B	97	9% 57%23%14%6%
2	D	97	4% 64%19%9%6%
3	E	334	% 66%25%6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	334	% 61% 31% 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SNF1-like protein kinase ssp2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	0	0
			939	608	159	163	9			
1	C	112	Total	C	N	O	S	0	1	0
			907	587	158	153	9			

- Molecule 2 is a protein called SPCC1919.03c protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	91	Total	C	N	O	S	0	0	0
			711	453	123	133	2			
2	D	91	Total	C	N	O	S	0	0	0
			714	456	122	134	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	202	MET	-	expression tag	UNP P78789
D	202	MET	-	expression tag	UNP P78789

- Molecule 3 is a protein called Protein C1556.08c.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	324	Total	C	N	O	S	0	0	0
			2543	1624	424	480	15			
3	E	324	Total	C	N	O	S	0	0	0
			2543	1623	424	482	14			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	1	ALA	-	expression tag	UNP Q10343
G	2	MET	-	expression tag	UNP Q10343

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	1	ALA	-	expression tag	UNP Q10343
E	2	MET	-	expression tag	UNP Q10343

- Molecule 4 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: $C_{10}H_{14}N_5O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	G	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	G	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	8	Total	O	0	0
			8	8		
6	B	10	Total	O	0	0
			10	10		
6	G	19	Total	O	0	0
			19	19		
6	C	11	Total	O	0	0
			11	11		
6	D	11	Total	O	0	0
			11	11		
6	E	20	Total	O	0	0
			20	20		


- Molecule 1: SNF1-like protein kinase *ssp2*



• Molecule 3: Protein C1556.08c



• Molecule 3: Protein C1556.08c



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	168.29Å 78.09Å 108.55Å 90.00° 124.13° 90.00°	Depositor
Resolution (Å)	48.28 – 2.70 48.28 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.3 (48.28-2.70) 96.3 (48.28-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.289 0.203 , 0.277	Depositor DCC
R_{free} test set	1558 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	64.9	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8540	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/963	1.05	8/1295 (0.6%)
1	C	0.71	0/933	0.98	2/1253 (0.2%)
2	B	0.72	0/728	1.21	4/995 (0.4%)
2	D	0.72	0/732	1.09	6/1002 (0.6%)
3	E	0.74	0/2585	0.98	0/3502
3	G	0.74	0/2585	0.99	4/3501 (0.1%)
All	All	0.73	0/8526	1.02	24/11548 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	4
3	E	0	3
3	G	0	4
All	All	0	11

There are no bond length outliers.

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	224	LEU	CB-CA-C	13.33	136.95	110.42
2	B	225	LYS	N-CA-CB	11.90	130.60	110.49
3	G	6	GLU	N-CA-C	-10.05	101.58	114.04
3	G	330	PHE	N-CA-CB	9.69	128.48	111.13
2	D	218	SER	N-CA-C	8.09	123.58	111.04

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	218	SER	Peptide
2	B	221	LEU	Peptide
2	B	222	GLN	Peptide
2	B	296	PHE	Peptide
3	G	262	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	939	0	936	27	0
1	C	907	0	909	24	0
2	B	711	0	717	50	0
2	D	714	0	713	39	0
3	E	2543	0	2572	65	0
3	G	2543	0	2583	81	0
4	G	23	0	12	0	0
5	E	54	0	24	1	0
5	G	27	0	12	4	0
6	A	8	0	0	0	0
6	B	10	0	0	0	0
6	C	11	0	0	1	0
6	D	11	0	0	0	0
6	E	20	0	0	0	0
6	G	19	0	0	3	0
All	All	8540	0	8478	272	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 272 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:219:ASN:CA	2:D:220:THR:HG22	1.17	1.56
2:D:219:ASN:HA	2:D:220:THR:CG2	1.18	1.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:543:SER:CB	1:A:544:HIS:HA	1.51	1.31
1:A:543:SER:HB3	1:A:544:HIS:CA	1.64	1.26
2:D:221:LEU:O	2:D:222:GLN:HB2	1.46	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/137 (82%)	104 (93%)	5 (4%)	3 (3%)	4	10
1	C	109/137 (80%)	104 (95%)	4 (4%)	1 (1%)	14	35
2	B	89/97 (92%)	72 (81%)	15 (17%)	2 (2%)	5	14
2	D	89/97 (92%)	70 (79%)	15 (17%)	4 (4%)	2	4
3	E	320/334 (96%)	302 (94%)	16 (5%)	2 (1%)	21	44
3	G	320/334 (96%)	303 (95%)	13 (4%)	4 (1%)	9	25
All	All	1039/1136 (92%)	955 (92%)	68 (6%)	16 (2%)	8	22

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	LYS
3	G	263	ASN
3	G	265	ASP
2	D	221	LEU
2	D	222	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/120 (83%)	90 (90%)	10 (10%)	7	19
1	C	95/120 (79%)	84 (88%)	11 (12%)	5	13
2	B	81/88 (92%)	71 (88%)	10 (12%)	4	12
2	D	81/88 (92%)	70 (86%)	11 (14%)	3	9
3	E	284/296 (96%)	246 (87%)	38 (13%)	4	9
3	G	284/296 (96%)	254 (89%)	30 (11%)	6	17
All	All	925/1008 (92%)	815 (88%)	110 (12%)	5	13

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	521	LEU
2	D	292	MET
3	E	310	LEU
3	E	224	SER
1	C	559	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	289	HIS
1	C	536	ASN
3	E	123	HIS
1	C	512	ASN
2	D	207	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ADP	E	1002	-	28,29,29	1.29	2 (7%)	43,45,45	1.94	10 (23%)
5	ADP	E	1004	-	28,29,29	1.44	4 (14%)	43,45,45	1.76	9 (20%)
4	AMP	G	1001	-	25,25,25	1.50	4 (16%)	37,38,38	2.20	12 (32%)
5	ADP	G	1003	-	28,29,29	1.56	5 (17%)	43,45,45	1.64	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	E	1002	-	-	6/16/32/32	0/3/3/3
5	ADP	E	1004	-	-	2/16/32/32	0/3/3/3
4	AMP	G	1001	-	-	2/10/26/26	0/3/3/3
5	ADP	G	1003	-	-	6/16/32/32	0/3/3/3

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	1003	ADP	C5-C4	5.62	1.49	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	1004	ADP	C5-C4	4.83	1.47	1.39
5	E	1002	ADP	C5-C4	4.59	1.47	1.39
4	G	1001	AMP	C5-C4	4.55	1.47	1.39
4	G	1001	AMP	C5-C6	2.79	1.48	1.41

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1001	AMP	C5-C4-N3	-6.33	118.01	126.72
5	E	1002	ADP	C5-C4-N3	-5.65	118.94	126.72
5	E	1002	ADP	N3-C4-N9	5.50	136.52	127.17
5	E	1004	ADP	C5-C4-N3	-5.15	119.62	126.72
4	G	1001	AMP	N3-C4-N9	4.77	135.28	127.17

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	1003	ADP	PA-O3A-PB-O3B
5	G	1003	ADP	C5'-O5'-PA-O1A
5	G	1003	ADP	C5'-O5'-PA-O2A
5	G	1003	ADP	C5'-O5'-PA-O3A
5	E	1002	ADP	C5'-O5'-PA-O1A


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	1004	ADP	1	0
5	G	1003	ADP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/137 (84%)	0.06	1 (0%) 81 80	40, 52, 59, 63	0
1	C	112/137 (81%)	0.31	2 (1%) 67 65	29, 52, 57, 62	1 (0%)
2	B	91/97 (93%)	0.61	9 (9%) 13 10	47, 52, 61, 65	0
2	D	91/97 (93%)	0.45	4 (4%) 39 35	45, 52, 60, 63	0
3	E	324/334 (97%)	0.14	5 (1%) 72 70	41, 52, 64, 76	0
3	G	324/334 (97%)	0.15	5 (1%) 72 70	27, 52, 61, 74	1 (0%)
All	All	1058/1136 (93%)	0.22	26 (2%) 58 55	27, 52, 61, 76	2 (0%)

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	247	TYR	6.0
3	G	119	THR	4.6
3	G	262	ALA	4.4
3	G	33	ARG	3.6
3	G	329	ASN	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ADP	G	1003	27/27	0.87	0.12	61,65,74,75	0
5	ADP	E	1004	27/27	0.87	0.12	70,72,80,81	0
5	ADP	E	1002	27/27	0.90	0.11	43,45,63,64	0
4	AMP	G	1001	23/23	0.91	0.10	41,47,48,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADP G 1003:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP E 1004:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP E 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AMP G 1001:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.