



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:43 PM UTC

PDB ID : 4QRC / pdb_00004qrc
Title : Crystal Structure of the Tyrosine Kinase Domain of FGF Receptor 4 in Complex with Ponatinib
Authors : Huang, Z.; Mohammadi, M.
Deposited on : 2014-06-30
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2475 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibroblast growth factor receptor 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	0	0
			2321	1482	406	418	15			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	431	MET	-	expression tag	UNP P22455
A	432	GLY	-	expression tag	UNP P22455
A	433	SER	-	expression tag	UNP P22455
A	434	SER	-	expression tag	UNP P22455
A	435	HIS	-	expression tag	UNP P22455
A	436	HIS	-	expression tag	UNP P22455
A	437	HIS	-	expression tag	UNP P22455
A	438	HIS	-	expression tag	UNP P22455
A	439	HIS	-	expression tag	UNP P22455
A	440	HIS	-	expression tag	UNP P22455
A	441	SER	-	expression tag	UNP P22455
A	442	GLN	-	expression tag	UNP P22455
A	443	ASP	-	expression tag	UNP P22455
A	444	PRO	-	expression tag	UNP P22455
A	477	ALA	CYS	engineered mutation	UNP P22455
A	664	GLU	ARG	engineered mutation	UNP P22455

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 3-(imidazo[1,2-b]pyridazin-3-ylethynyl)-4-methyl-N-{4-[(4-methylpiperazin-1-yl)methyl]-3-(trifluoromethyl)phenyl}benzamide (CCD ID: 0LI) (formula: C₂₉H₂₇F₃N₆O).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			39	29	3	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	95	Total	O	0	0
			95	95		

- Molecule 1: Fibroblast growth factor receptor 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.72Å 61.59Å 60.31Å 90.00° 97.94° 90.00°	Depositor
Resolution (Å)	37.02 – 1.90 37.02 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (37.02-1.90) 98.0 (37.02-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.40 (at 1.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.170 , 0.207 0.172 , 0.208	Depositor DCC
R_{free} test set	2000 reflections (5.38%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2475	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, OLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2381	0.76	0/3236

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	629	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2321	0	2313	33	0
2	A	20	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	39	0	27	11	0
4	A	95	0	0	0	0
All	All	2475	0	2340	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:520:GLU:HA	1:A:633:LEU:HD13	1.70	0.72
1:A:633:LEU:HG	3:A:802:OLI:H15	1.74	0.69
1:A:608:CYS:SG	3:A:802:OLI:H24A	2.36	0.66
1:A:633:LEU:HA	3:A:802:OLI:H22A	1.81	0.63
1:A:456:LEU:HD12	1:A:542:GLN:HG3	1.81	0.62
1:A:551:GLU:O	3:A:802:OLI:H1	2.06	0.55
1:A:611:ARG:HB2	1:A:635:ARG:HD2	1.89	0.54
1:A:481:VAL:HG22	1:A:503:LYS:HD3	1.90	0.53
1:A:683:PHE:CE2	1:A:718:LEU:HD22	2.44	0.52
1:A:630:ASP:HA	1:A:631:PHE:HB2	1.92	0.52
1:A:520:GLU:HA	1:A:633:LEU:CD1	2.40	0.50
1:A:619:LEU:HD21	1:A:631:PHE:HE2	1.77	0.49
1:A:633:LEU:CG	3:A:802:OLI:H15	2.41	0.49
1:A:482:VAL:HG12	1:A:483:ARG:O	2.13	0.48
1:A:505:LEU:HD11	1:A:514:LEU:HD13	1.95	0.48
1:A:550:VAL:HG21	3:A:802:OLI:C5	2.43	0.47
1:A:481:VAL:HG13	3:A:802:OLI:H11A	1.96	0.47
1:A:630:ASP:N	1:A:630:ASP:OD1	2.48	0.47
1:A:552:CYS:HA	3:A:802:OLI:C1	2.46	0.46
1:A:604:GLU:HG3	1:A:668:HIS:CG	2.51	0.45
1:A:552:CYS:HA	3:A:802:OLI:H1	1.98	0.45
1:A:503:LYS:HB2	3:A:802:OLI:H7	1.98	0.45
1:A:586:PHE:HB3	1:A:587:PRO:HD3	1.99	0.44
1:A:683:PHE:CZ	1:A:718:LEU:HD22	2.53	0.44
1:A:630:ASP:OD2	3:A:802:OLI:H21A	2.18	0.44
1:A:616:ARG:HD3	2:A:801:SO4:O4	2.18	0.44
1:A:630:ASP:HA	1:A:631:PHE:CB	2.49	0.43
1:A:476:GLY:O	1:A:506:LYS:HE2	2.19	0.42
1:A:520:GLU:O	1:A:524:MET:HG3	2.20	0.42
1:A:619:LEU:HD21	1:A:631:PHE:CE2	2.53	0.41
1:A:630:ASP:HB2	1:A:631:PHE:C	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:540:CYS:HB2	1:A:547:TYR:HB2	2.02	0.41
1:A:471:LYS:HA	1:A:472:PRO:HD3	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	295/323 (91%)	288 (98%)	6 (2%)	1 (0%)	36 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	633	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	249/275 (90%)	243 (98%)	6 (2%)	43 38

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	503	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	507	ASP
1	A	527	ILE
1	A	616	ARG
1	A	644	LYS
1	A	718	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	532	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	805	-	4,4,4	0.25	0	6,6,6	0.05	0
2	SO4	A	801	-	4,4,4	0.21	0	6,6,6	0.27	0
2	SO4	A	804	-	4,4,4	0.25	0	6,6,6	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	803	-	4,4,4	0.25	0	6,6,6	0.17	0
3	OLI	A	802	-	43,43,43	1.55	7 (16%)	59,62,62	2.82	22 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLI	A	802	-	-	3/21/33/33	0/5/5/5

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	OLI	C12-N2	4.51	1.48	1.35
3	A	802	OLI	C3-C2	4.32	1.49	1.42
3	A	802	OLI	C5-C4	3.36	1.50	1.43
3	A	802	OLI	C13-N2	3.02	1.47	1.41
3	A	802	OLI	C1-N1	-2.52	1.32	1.37
3	A	802	OLI	C83-N81	2.42	1.34	1.31
3	A	802	OLI	C1-C2	2.08	1.39	1.37

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	OLI	C1-C2-N82	8.97	106.79	103.61
3	A	802	OLI	C1-N1-C84	7.68	110.24	105.11
3	A	802	OLI	C84-N82-N81	-6.89	119.96	126.72
3	A	802	OLI	O1-C12-C9	-5.16	110.67	120.90
3	A	802	OLI	C83-N81-N82	4.35	120.25	113.68
3	A	802	OLI	C23-N4-C22	4.25	116.30	109.54
3	A	802	OLI	C24-N3-C21	4.20	117.89	108.84
3	A	802	OLI	C2-C1-N1	-4.18	107.88	112.35
3	A	802	OLI	C8-C9-C10	3.91	123.77	119.25
3	A	802	OLI	C82-C83-N81	-3.86	121.10	124.83
3	A	802	OLI	C3-C2-N82	3.66	128.45	122.52
3	A	802	OLI	C1-C2-C3	-3.62	124.76	130.61
3	A	802	OLI	C9-C10-C5	-3.42	117.56	120.60
3	A	802	OLI	C25-N4-C22	3.35	117.02	110.63
3	A	802	OLI	C21-C22-N4	-3.01	106.02	110.86
3	A	802	OLI	C25-N4-C23	2.91	116.17	110.63
3	A	802	OLI	C7-C6-C5	2.89	121.01	117.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	0LI	C6-C5-C4	2.67	124.00	120.46
3	A	802	0LI	C7-C8-C9	-2.59	118.03	120.80
3	A	802	0LI	C24-C23-N4	2.44	114.78	110.86
3	A	802	0LI	F1-C19-C17	-2.31	108.55	112.65
3	A	802	0LI	O1-C12-N2	2.26	129.47	123.75

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	802	0LI	C2-C3-C4-C5
3	A	802	0LI	C16-C20-N3-C24
3	A	802	0LI	C16-C20-N3-C21

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	SO4	1	0
3	A	802	0LI	11	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/323 (91%)	0.19	23 (7%) 19 21	10, 22, 58, 77	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	632	GLY	8.2
1	A	633	LEU	7.3
1	A	478	PHE	6.6
1	A	631	PHE	5.5
1	A	572	LEU	5.0
1	A	477	ALA	4.0
1	A	570	PRO	3.7
1	A	476	GLY	3.5
1	A	569	GLY	3.2
1	A	573	SER	3.1
1	A	630	ASP	3.0
1	A	571	ASP	3.0
1	A	508	ASN	2.8
1	A	509	ALA	2.6
1	A	574	PRO	2.6
1	A	507	ASP	2.5
1	A	456	LEU	2.4
1	A	750	VAL	2.4
1	A	634	ALA	2.3
1	A	527	ILE	2.2
1	A	479	GLY	2.1
1	A	511	ASP	2.1
1	A	580	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	803	5/5	0.75	0.15	67,69,73,73	0
3	OLI	A	802	39/39	0.82	0.14	25,35,40,44	0
2	SO4	A	804	5/5	0.83	0.17	54,60,62,64	0
2	SO4	A	805	5/5	0.89	0.10	69,70,71,72	0
2	SO4	A	801	5/5	0.96	0.07	32,35,38,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 0LI A 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.