



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 08:47 AM UTC

PDB ID : 7QR0 / pdb_00007qr0
Title : SpCas9 bound to TRAC off-target1 DNA substrate
Authors : Pacesa, M.; Jinek, M.
Deposited on : 2022-01-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)
RNA backbone	3983	1090 (2.60-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	84	74% 20% • •
2	B	1368	4% 86% 11% •
3	C	28	86% 14%
4	D	12	67% 25% 8%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called TRAC sgRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	82	Total	C	N	O	P	0	0	0
			1722	771	310	560	81			

- Molecule 2 is a protein called CRISPR-associated endonuclease Cas9/Csn1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1333	Total	C	N	O	S	0	0	0
			10891	6938	1891	2040	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	ALA	ASP	engineered mutation	UNP Q99ZW2
B	840	ALA	HIS	engineered mutation	UNP Q99ZW2

- Molecule 3 is a DNA chain called TRAC off-target1 target strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	28	Total	C	N	O	P	0	0	0
			570	273	105	165	27			

- Molecule 4 is a DNA chain called TRAC off-target1 non-target strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	11	Total	C	N	O	P	0	0	1
			208	99	36	63	10			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	6	Total K 6 6	0	0
6	B	9	Total K 9 9	0	0
6	C	2	Total K 2 2	0	0



- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	165	Total O 165 165	0	0
7	B	438	Total O 438 438	0	0
7	C	43	Total O 43 43	0	0
7	D	12	Total O 12 12	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRAC sgRNA



• Molecule 2: CRISPR-associated endonuclease Cas9/Csn1




• Molecule 3: TRAC off-target1 target strand

Chain C:  86% 14%



● Molecule 4: TRAC off-target1 non-target strand

Chain D:  67% 25% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	177.55Å 67.42Å 187.25Å 90.00° 111.49° 90.00°	Depositor
Resolution (Å)	47.71 – 2.30 47.71 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.71-2.30) 99.9 (47.71-2.30)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.17.1 _3660	Depositor
R, R_{free}	0.194 , 0.224 0.195 , 0.225	Depositor DCC
R_{free} test set	5626 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	49.3	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14068	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/1928	0.28	0/3004
2	B	0.09	0/11082	0.27	0/14888
3	C	0.19	0/639	0.41	0/984
4	D	0.20	0/232	0.43	0/358
All	All	0.10	0/13881	0.29	0/19234

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1722	0	862	8	0
2	B	10891	0	11064	92	0
3	C	570	0	317	3	0
4	D	208	0	115	3	0
5	A	2	0	0	0	0
6	A	6	0	0	0	0
6	B	9	0	0	0	0
6	C	2	0	0	0	0
7	A	165	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	438	0	0	7	0
7	C	43	0	0	0	0
7	D	12	0	0	0	0
All	All	14068	0	12358	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:266:LEU:HD22	2:B:294:LYS:HD3	1.67	0.75
2:B:80:CYS:SG	7:B:1880:HOH:O	2.50	0.68
2:B:1335:ARG:NH2	4:D:2:DG:O6	2.30	0.63
2:B:898:ASP:O	2:B:905:ARG:NH2	2.31	0.63
2:B:158:LEU:HD22	2:B:419:LEU:HD12	1.82	0.62
4:D:6:DT:H2''	4:D:7:DG:H5''	1.81	0.61
2:B:260:GLU:HB3	2:B:281:GLN:HE22	1.66	0.60
2:B:870:VAL:HG22	2:B:871:PRO:HD2	1.83	0.60
2:B:1151:LYS:HD2	2:B:1158:LYS:HD2	1.84	0.60
2:B:790:GLU:HG2	2:B:889:ALA:HA	1.85	0.59
2:B:654:ARG:HD2	2:B:656:TYR:CZ	2.37	0.59
2:B:1215:ALA:HB2	2:B:1221:GLN:HG3	1.86	0.56
1:A:59:U:OP1	2:B:467:ARG:NH2	2.38	0.55
2:B:1110:ILE:HG23	2:B:1122:ARG:HD2	1.87	0.55
2:B:258:LEU:HD21	2:B:282:ILE:HD11	1.89	0.55
2:B:763:MET:HE1	2:B:931:VAL:HG21	1.88	0.55
2:B:967:ARG:NH2	7:B:1509:HOH:O	2.34	0.54
2:B:179:SER:HB3	2:B:299:ALA:HB2	1.89	0.54
2:B:614:ASP:OD1	2:B:664:ARG:NH2	2.35	0.53
2:B:546:LYS:NZ	2:B:550:ASP:OD2	2.42	0.53
2:B:880:LYS:HE2	2:B:884:ARG:HH22	1.73	0.53
2:B:510:LYS:NZ	7:B:1518:HOH:O	2.41	0.53
2:B:1041:ASN:HD22	2:B:1044:ASN:HD21	1.57	0.52
1:A:5:U:OP1	2:B:515:TYR:OH	2.24	0.51
2:B:600:ILE:HG23	2:B:650:GLN:HB3	1.92	0.51
2:B:501:ASN:HB3	2:B:708:ILE:HD12	1.91	0.51
2:B:276:ASP:HB3	2:B:599:LYS:NZ	2.26	0.51
2:B:788:ILE:HG23	2:B:793:SER:HB3	1.92	0.51
2:B:400:ARG:NH2	2:B:406:ASP:OD2	2.37	0.50
1:A:33:G:N2	1:A:36:A:OP2	2.44	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1205:GLU:OE1	2:B:1359:ARG:NH2	2.44	0.50
2:B:859:ARG:NH2	7:B:1524:HOH:O	2.44	0.50
2:B:216:LEU:HD23	2:B:220:ARG:HG2	1.94	0.49
2:B:149:ALA:H	2:B:426:GLN:HE22	1.60	0.49
1:A:44:U:O2'	2:B:402:GLN:OE1	2.25	0.49
2:B:1051:THR:HA	2:B:1057:ILE:HA	1.95	0.49
1:A:49:A:N3	2:B:1122:ARG:NH2	2.56	0.48
2:B:762:GLU:HG2	2:B:990:ASN:HD21	1.78	0.48
1:A:33:G:H5'	1:A:34:A:OP2	2.14	0.48
2:B:468:LYS:N	2:B:481:VAL:O	2.40	0.47
2:B:1136:SER:HA	4:D:2:DG:O3'	2.14	0.47
2:B:1047:LYS:O	2:B:1076:LYS:NZ	2.40	0.47
2:B:455:LEU:HA	2:B:465:MET:HE2	1.97	0.46
2:B:508:LEU:HD21	2:B:664:ARG:HB2	1.96	0.46
2:B:712:GLN:NE2	7:B:1539:HOH:O	2.49	0.46
2:B:704:PHE:O	2:B:708:ILE:HG12	2.16	0.46
2:B:817:GLN:O	2:B:882:TYR:OH	2.30	0.45
2:B:780:ARG:NH1	2:B:806:LEU:O	2.47	0.45
2:B:1117:ASP:N	2:B:1117:ASP:OD1	2.49	0.45
2:B:1229:PRO:HD2	2:B:1232:TYR:HD2	1.81	0.45
2:B:468:LYS:HG3	2:B:483:ASP:HB2	1.98	0.44
2:B:214:ALA:O	2:B:221:ARG:HD3	2.18	0.44
2:B:1042:ILE:HG23	2:B:1043:MET:HG2	1.99	0.44
2:B:137:HIS:HA	2:B:322:ILE:HD11	1.99	0.44
2:B:465:MET:HE3	2:B:465:MET:HB3	1.86	0.44
2:B:822:MET:HG2	2:B:856:VAL:HG22	1.99	0.44
2:B:400:ARG:HH22	2:B:406:ASP:CG	2.24	0.44
2:B:629:ARG:HE	2:B:653:ARG:HA	1.81	0.44
2:B:497:ASN:HD21	3:C:11:DG:P	2.40	0.44
1:A:72:U:H5'	1:A:73:G:OP2	2.17	0.43
2:B:447:ARG:NH2	7:B:1545:HOH:O	2.51	0.43
2:B:489:GLN:HG3	2:B:625:LEU:HD21	2.00	0.43
2:B:725:ALA:O	2:B:734:LYS:NZ	2.51	0.43
2:B:967:ARG:NH1	2:B:986:ASP:OD1	2.52	0.43
2:B:597:LEU:O	2:B:601:ILE:HG12	2.19	0.43
2:B:841:ILE:HD13	2:B:900:LEU:HG	1.99	0.43
1:A:27:G:N2	1:A:44:U:OP2	2.51	0.43
2:B:1122:ARG:HG2	2:B:1134:PHE:CE2	2.53	0.43
2:B:557:ARG:NH1	2:B:596:ASP:OD1	2.36	0.43
2:B:1065:THR:HG22	2:B:1072:ILE:HA	2.00	0.43
2:B:1334:LYS:NZ	3:C:-5:DA:OP2	2.36	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1231:LYS:HE3	2:B:1232:TYR:CZ	2.54	0.43
2:B:853:ASP:OD1	2:B:893:THR:OG1	2.30	0.42
2:B:878:LYS:HG3	2:B:879:MET:HG2	2.01	0.42
2:B:901:THR:O	2:B:904:GLU:HG2	2.19	0.42
2:B:214:ALA:HB1	2:B:216:LEU:HD13	2.01	0.42
2:B:850:ASP:OD1	2:B:855:LYS:HD3	2.19	0.42
2:B:518:PHE:CD1	2:B:667:ILE:HD12	2.53	0.42
2:B:956:ILE:HD11	2:B:998:ILE:HD13	2.01	0.42
2:B:1002:PRO:HD3	2:B:1066:ASN:HD21	1.84	0.42
2:B:498:PHE:CZ	2:B:506:LYS:HE3	2.55	0.42
2:B:1004:LEU:HD11	2:B:1042:ILE:HD11	2.02	0.42
2:B:275:LEU:O	2:B:279:LEU:N	2.48	0.42
2:B:812:TYR:CZ	2:B:816:LEU:HD11	2.55	0.42
2:B:439:LYS:HG2	2:B:476:TRP:CD1	2.55	0.41
2:B:345:GLU:H	2:B:345:GLU:CD	2.29	0.41
2:B:973:TYR:HB3	2:B:1237:TYR:CD2	2.55	0.41
2:B:560:THR:HA	2:B:586:ARG:HA	2.02	0.41
2:B:184:LEU:HD12	2:B:299:ALA:HB2	2.02	0.41
2:B:225:LEU:HD23	2:B:242:ILE:HG21	2.04	0.41
2:B:635:ARG:NH1	7:B:1543:HOH:O	2.50	0.40
2:B:253:LYS:HD2	2:B:261:ASP:OD1	2.22	0.40
2:B:187:GLN:O	2:B:191:THR:HG22	2.22	0.40
2:B:1153:LYS:H	2:B:1153:LYS:HG3	1.67	0.40
2:B:962:LEU:HD23	2:B:962:LEU:HA	1.95	0.40
2:B:1266:LEU:HD22	2:B:1309:ILE:HD12	2.04	0.40
2:B:226:ILE:HD13	2:B:226:ILE:HA	1.93	0.40
3:C:14:DA:H2'	3:C:15:DG:C8	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	1323/1368 (97%)	1287 (97%)	36 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	1195/1225 (98%)	1182 (99%)	13 (1%)	65	81

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	75	ARG
2	B	182	ASP
2	B	228	GLN
2	B	260	GLU
2	B	272	ASP
2	B	310	THR
2	B	512	SER
2	B	679	ILE
2	B	762	GLU
2	B	790	GLU
2	B	856	VAL
2	B	870	VAL
2	B	1007	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	178	ASN
2	B	251	ASN
2	B	281	GLN
2	B	341	GLN
2	B	426	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	459	ASN
2	B	497	ASN
2	B	501	ASN
2	B	511	HIS
2	B	556	ASN
2	B	826	GLN
2	B	885	GLN
2	B	920	GLN
2	B	980	ASN
2	B	990	ASN
2	B	1041	ASN
2	B	1101	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	80/84 (95%)	13 (16%)	0

All (13) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	20	A
1	A	28	A
1	A	29	G
1	A	35	A
1	A	36	A
1	A	37	U
1	A	51	A
1	A	56	U
1	A	59	U
1	A	68	A
1	A	72	U
1	A	74	A
1	A	77	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 19 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	82/84 (97%)	-0.58	1 (1%) 76 77	38, 54, 151, 189	0
2	B	1333/1368 (97%)	0.26	50 (3%) 44 46	33, 62, 108, 146	0
3	C	28/28 (100%)	-0.38	0 100 100	44, 54, 107, 115	0
4	D	11/12 (91%)	-0.07	0 100 100	45, 68, 140, 142	0
All	All	1454/1492 (97%)	0.20	51 (3%) 47 49	33, 61, 112, 189	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	188	LEU	4.9
2	B	1052	LEU	4.3
2	B	286	TYR	4.3
2	B	713	VAL	4.1
2	B	290	PHE	4.1
2	B	293	ALA	4.1
2	B	264	LEU	3.5
2	B	1057	ILE	3.5
2	B	191	THR	3.3
2	B	278	LEU	3.2
2	B	775	LYS	3.2
2	B	259	ALA	3.1
2	B	292	ALA	3.1
2	B	291	LEU	3.1
1	A	1	A	3.0
2	B	289	LEU	3.0
2	B	804	THR	2.9
2	B	1242	TYR	2.9
2	B	1012	ASP	2.9
2	B	3	LYS	2.9
2	B	1246	LYS	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	186	ILE	2.9
2	B	1367	GLY	2.8
2	B	287	ALA	2.8
2	B	714	SER	2.8
2	B	282	ILE	2.7
2	B	279	LEU	2.7
2	B	297	SER	2.6
2	B	1306	ALA	2.6
2	B	720	LEU	2.5
2	B	862	LYS	2.5
2	B	176	PRO	2.4
2	B	296	LEU	2.3
2	B	301	LEU	2.3
2	B	272	ASP	2.3
2	B	1263	LYS	2.3
2	B	179	SER	2.3
2	B	806	LEU	2.2
2	B	294	LYS	2.2
2	B	300	ILE	2.2
2	B	298	ASP	2.2
2	B	571	LYS	2.2
2	B	312	ILE	2.1
2	B	268	LYS	2.1
2	B	717	GLY	2.1
2	B	214	ALA	2.1
2	B	267	SER	2.1
2	B	310	THR	2.0
2	B	766	GLU	2.0
2	B	178	ASN	2.0
2	B	727	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	K	B	1408	1/1	0.74	0.12	128,128,128,128	0
6	K	C	102	1/1	0.82	0.16	119,119,119,119	0
6	K	B	1407	1/1	0.86	0.08	115,115,115,115	0
6	K	B	1404	1/1	0.89	0.09	88,88,88,88	0
6	K	A	108	1/1	0.89	0.10	85,85,85,85	0
6	K	A	105	1/1	0.90	0.12	77,77,77,77	0
6	K	C	101	1/1	0.92	0.11	94,94,94,94	0
6	K	A	107	1/1	0.93	0.15	89,89,89,89	0
6	K	B	1405	1/1	0.93	0.08	97,97,97,97	0
6	K	B	1403	1/1	0.93	0.07	77,77,77,77	0
6	K	B	1402	1/1	0.95	0.07	75,75,75,75	0
6	K	B	1409	1/1	0.95	0.08	86,86,86,86	0
6	K	B	1401	1/1	0.96	0.06	62,62,62,62	0
6	K	B	1406	1/1	0.97	0.05	57,57,57,57	0
5	MG	A	101	1/1	0.98	0.05	57,57,57,57	0
6	K	A	106	1/1	0.98	0.09	54,54,54,54	0
6	K	A	104	1/1	0.98	0.06	58,58,58,58	0
5	MG	A	102	1/1	0.99	0.03	37,37,37,37	0
6	K	A	103	1/1	0.99	0.03	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.