



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 28, 2026 – 10:24 PM UTC

PDB ID : 1QSF / pdb_00001qsf
Title : STRUCTURE OF A6-TCR BOUND TO HLA-A2 COMPLEXED WITH ALTERED HTLV-1 TAX PEPTIDE Y8A
Authors : Ding, Y.H.; Baker, B.M.; Garboczi, D.N.; Biddison, W.E.; Wiley, D.C.
Deposited on : 1999-06-21
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	274	52% 41% 7%
2	B	100	57% 38% .
3	C	9	11% 78% 11%
4	D	200	40% 46% 12% .
5	E	243	36% 54% 10%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6486 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC CLASS I HLA-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	1	0	0
			2179	1364	391	415	9			

- Molecule 2 is a protein called BETA-2 MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	ALA	engineered mutation	UNP P61769

- Molecule 3 is a protein called TAX PEPTIDE Y8A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			70	50	9	11			

- Molecule 4 is a protein called HUMAN T-CELL RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	200	Total	C	N	O	S	0	0	0
			1497	938	245	308	6			

- Molecule 5 is a protein called HUMAN T-CELL RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	243	Total	C	N	O	S	0	0	0
			1882	1184	328	362	8			

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	10	Total 10	O 10	0	0
6	B	6	Total 6	O 6	0	0
6	C	1	Total 1	O 1	0	0
6	D	1	Total 1	O 1	0	0
6	E	3	Total 3	O 3	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: MHC CLASS I HLA-A



• Molecule 2: BETA-2 MICROGLOBULIN




• Molecule 3: TAX PEPTIDE Y8A



• Molecule 4: HUMAN T-CELL RECEPTOR





• Molecule 5: HUMAN T-CELL RECEPTOR

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	224.66Å 48.46Å 93.74Å 90.00° 90.46° 90.00°	Depositor
Resolution (Å)	25.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (25.00-2.80)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.245 , 0.286	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6486	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/2241	1.06	7/3043 (0.2%)
2	B	0.60	0/860	1.05	4/1162 (0.3%)
3	C	0.76	0/72	1.50	3/97 (3.1%)
4	D	0.56	0/1530	1.17	15/2084 (0.7%)
5	E	0.53	0/1935	1.10	15/2641 (0.6%)
All	All	0.56	0/6638	1.10	44/9027 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	D	0	1

There are no bond length outliers.

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	123	PHE	CA-C-N	-9.60	110.50	120.38
5	E	123	PHE	C-N-CA	-9.60	110.50	120.38
4	D	196	SER	N-CA-C	8.67	120.71	108.74
1	A	198	GLU	N-CA-C	8.42	122.35	108.96
2	B	33	SER	N-CA-C	8.38	120.42	111.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	D	88	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2179	0	2015	124	0
2	B	837	0	803	33	0
3	C	70	0	75	10	0
4	D	1497	0	1363	156	0
5	E	1882	0	1763	178	0
6	A	10	0	0	0	0
6	B	6	0	0	1	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	E	3	0	0	1	0
All	All	6486	0	6019	461	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 461 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:THR:HG22	1:A:229:GLU:H	1.21	1.04
4:D:168:MET:CE	5:E:197:ARG:HG2	1.92	0.98
2:B:17:ASN:OD1	2:B:73:THR:HA	1.66	0.96
4:D:168:MET:HE2	5:E:197:ARG:HG2	1.51	0.93
4:D:168:MET:HE2	5:E:197:ARG:CG	1.99	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/274 (99%)	242 (89%)	22 (8%)	8 (3%)	3	13
2	B	98/100 (98%)	88 (90%)	7 (7%)	3 (3%)	3	12
3	C	7/9 (78%)	7 (100%)	0	0	100	100
4	D	198/200 (99%)	166 (84%)	22 (11%)	10 (5%)	1	5
5	E	241/243 (99%)	208 (86%)	23 (10%)	10 (4%)	2	8
All	All	816/826 (99%)	711 (87%)	74 (9%)	31 (4%)	2	9

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	SER
1	A	253	GLN
4	D	131	SER
4	D	132	LYS
4	D	201	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/229 (96%)	200 (91%)	19 (9%)	9	30
2	B	95/95 (100%)	86 (90%)	9 (10%)	8	26
3	C	7/7 (100%)	7 (100%)	0	100	100
4	D	160/178 (90%)	146 (91%)	14 (9%)	9	29
5	E	200/207 (97%)	183 (92%)	17 (8%)	10	31
All	All	681/716 (95%)	622 (91%)	59 (9%)	9	30

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	11	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	174	THR
4	D	110	THR
5	E	159	LEU
5	E	80	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	30	GLN
5	E	22	GLN
5	E	235	GLN
5	E	6	GLN
5	E	37	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.