



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:17 PM UTC

PDB ID : 4QTF / pdb_00004qtf
Title : Structure and specificity of L-D-Transpeptidase from Mycobacterium tuberculosis and antibiotic resistance: Calcium binding promotes dimer formation
Authors : Gokulan, K.; Varughese, K.I.
Deposited on : 2014-07-07
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	349	6% 85% 13%

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L,d-transpeptidase LdtB.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	0	0
			2635	1651	462	514	8			

- Molecule 2 is GLCNAC(BETA1-4)-MURNAC(1,6-ANHYDRO)-L-ALA-GAMMA-D-GLU-MESO-A2PM-D-ALA (CCD ID: MLD) (formula: $C_{37}H_{59}N_7O_{20}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			46	27	6	13		

- Molecule 3 is (3S,5S)-3-({[(aminomethyl)amino]methyl}sulfanyl)-5-[(2S)-1,3-dioxobutan-2-yl]-L-proline (CCD ID: 3V5) (formula: C₁₁H₁₉N₃O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			19	11	3	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	76	Total	O	0	0
			76	76		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L,d-transpeptidase LdtB



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	57.27Å 65.88Å 207.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.35 – 2.00 42.35 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (42.35-2.00) 99.4 (42.35-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.66 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.213 , 0.258 0.222 , 0.258	Depositor DCC
R_{free} test set	1356 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.1	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2776	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MLD, 3V5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	6/2701 (0.2%)	1.13	7/3697 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	354	CYS	C-N	-23.76	1.00	1.33
1	A	353	GLY	C-N	5.51	1.41	1.33
1	A	310	VAL	CA-CB	5.39	1.59	1.53
1	A	399	ASP	C-O	-5.26	1.18	1.24
1	A	209	ARG	N-CA	5.17	1.52	1.45
1	A	135	GLY	C-O	-5.11	1.20	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	353	GLY	O-C-N	-9.51	110.33	122.70
1	A	353	GLY	CA-C-N	7.49	132.13	121.42
1	A	353	GLY	C-N-CA	7.49	132.13	121.42
1	A	354	CYS	CA-C-N	7.40	135.68	121.54
1	A	354	CYS	C-N-CA	7.40	135.68	121.54
1	A	354	CYS	O-C-N	-6.43	115.42	123.01
1	A	305	SER	CB-CA-C	-5.08	102.68	110.81

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	353	GLY	Mainchain
1	A	354	CYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2635	0	2531	30	0
2	A	46	0	37	6	0
3	A	19	0	15	4	0
4	A	76	0	0	1	0
All	All	2776	0	2583	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:ASP:OD2	1:A:111:ARG:NH1	2.00	0.93
1:A:291:ILE:H	1:A:395:ASN:HD21	1.17	0.89
1:A:93:MET:HE2	1:A:126:LEU:HD13	1.65	0.78
1:A:289:ASN:HD21	1:A:380:THR:H	1.30	0.78
1:A:302:ILE:HD11	1:A:321:ASP:OD1	1.94	0.68
2:A:501:MLD:HAJ2	2:A:501:MLD:C1	2.25	0.66
1:A:127:ASN:HD21	1:A:140:GLN:HE21	1.48	0.61
2:A:501:MLD:HA1	2:A:501:MLD:HAF2	1.84	0.60
1:A:332:GLY:O	3:A:502:3V5:H3	2.02	0.59
1:A:305:SER:HB3	1:A:310:VAL:O	2.03	0.58
2:A:501:MLD:HA1	2:A:501:MLD:CAF	2.34	0.57
1:A:308:TYR:N	1:A:308:TYR:CD1	2.72	0.57
2:A:501:MLD:HAJ2	2:A:501:MLD:C2	2.35	0.57
1:A:223:ASP:OD1	1:A:246:HIS:HD2	1.92	0.51
1:A:79:VAL:CG1	1:A:93:MET:HE1	2.41	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:PRO:HD2	1:A:340:TRP:CZ3	2.47	0.50
1:A:156:VAL:HG12	1:A:243:VAL:HG11	1.94	0.50
1:A:327:GLN:HB2	1:A:334:PHE:CE2	2.47	0.49
1:A:332:GLY:HA2	3:A:502:3V5:H4	1.94	0.48
2:A:501:MLD:HAJ2	2:A:501:MLD:O1	2.13	0.48
1:A:93:MET:CE	1:A:126:LEU:HD13	2.39	0.48
1:A:302:ILE:CD1	1:A:321:ASP:OD1	2.61	0.48
1:A:106:SER:OG	1:A:108:ASP:OD1	2.29	0.47
1:A:224:VAL:HB	1:A:245:THR:HG22	1.96	0.47
1:A:293:ILE:O	1:A:391:LEU:HD13	2.16	0.46
2:A:501:MLD:HAJ2	2:A:501:MLD:N2	2.31	0.46
1:A:132:GLY:C	1:A:134:GLY:H	2.24	0.45
1:A:99:ARG:HD3	4:A:642:HOH:O	2.18	0.44
1:A:313:ASN:C	1:A:313:ASN:HD22	2.27	0.43
1:A:79:VAL:HG11	1:A:93:MET:HE1	2.00	0.43
1:A:332:GLY:CA	3:A:502:3V5:H4	2.48	0.43
1:A:303:MET:SD	3:A:502:3V5:H13	2.59	0.42
1:A:177:ASN:ND2	1:A:206:ARG:HE	2.18	0.42
1:A:305:SER:OG	1:A:318:TYR:N	2.51	0.41
1:A:186:LYS:O	1:A:189:LYS:NZ	2.52	0.41
1:A:172:ILE:HD13	1:A:188:ILE:HD13	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/349 (99%)	333 (96%)	13 (4%)	1 (0%)	36	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	309	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	280/280 (100%)	272 (97%)	8 (3%)	37 40

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	THR
1	A	121	ASN
1	A	308	TYR
1	A	310	VAL
1	A	313	ASN
1	A	327	GLN
1	A	350	THR
1	A	393	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	A	127	ASN
1	A	177	ASN
1	A	246	HIS
1	A	289	ASN
1	A	313	ASN
1	A	316	ASN
1	A	327	GLN
1	A	363	GLN
1	A	395	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	3V5	A	502	1	15,19,19	1.80	5 (33%)	11,25,25	1.82	2 (18%)
2	MLD	A	501	-	46,47,66	1.64	9 (19%)	57,64,93	2.56	22 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3V5	A	502	1	-	8/14/31/31	0/1/1/1
2	MLD	A	501	-	-	19/50/76/114	0/3/2/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	MLD	CAF-CAE	-4.39	1.41	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	MLD	CA1-CA2	3.46	1.58	1.53
2	A	501	MLD	CA3-CA2	3.39	1.58	1.53
2	A	501	MLD	CA2-NAD	3.33	1.51	1.45
3	A	502	3V5	CAV-NAO	-3.21	1.43	1.48
2	A	501	MLD	C2-C1	3.19	1.54	1.51
3	A	502	3V5	CAW-SAQ	-3.13	1.77	1.82
3	A	502	3V5	CAX-NAO	-3.02	1.42	1.46
2	A	501	MLD	OAH-CAI	2.92	1.50	1.44
2	A	501	MLD	CAL-N	2.60	1.39	1.34
3	A	502	3V5	CAX-CAS	-2.56	1.48	1.52
3	A	502	3V5	OAH-CAS	-2.42	1.22	1.30
2	A	501	MLD	OA6-CA1	2.21	1.46	1.41
2	A	501	MLD	OAH-CA3	2.01	1.49	1.43

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	MLD	CBC-CBB-CAO	-6.59	101.00	113.16
2	A	501	MLD	OAG-CAE-CAF	-5.91	111.53	122.05
2	A	501	MLD	OAH-CAI-CAL	5.46	122.52	111.33
2	A	501	MLD	CA2-NAD-CAE	4.92	134.62	123.11
2	A	501	MLD	C1-O1-CA4	4.64	123.95	114.72
3	A	502	3V5	CAJ-NAN-CAK	4.43	122.70	113.25
2	A	501	MLD	OA6-CA1-CA2	4.29	117.58	111.08
2	A	501	MLD	CAJ-CAI-CAL	-4.18	100.68	111.11
2	A	501	MLD	CA1-CA2-NAD	3.89	117.44	110.92
2	A	501	MLD	CAP-CAO-NAN	3.83	119.45	110.57
2	A	501	MLD	CBB-CAO-NAN	-3.75	103.49	110.91
2	A	501	MLD	OA5-CA1-OA6	-3.51	103.22	105.97
2	A	501	MLD	CAI-CAL-N	3.43	129.01	118.15
2	A	501	MLD	CAF-CAE-NAD	3.32	121.63	116.12
2	A	501	MLD	OAH-CAI-CAJ	2.90	115.61	107.54
2	A	501	MLD	OAM-CAL-N	-2.90	117.77	122.96
2	A	501	MLD	OAG-CAE-NAD	2.74	126.83	121.98
2	A	501	MLD	OBE-CBD-CBC	-2.72	117.09	122.02
2	A	501	MLD	CA4-CA3-CA2	-2.57	105.75	110.55
2	A	501	MLD	OAM-CAL-CAI	-2.51	113.19	119.59
2	A	501	MLD	CBB-CBC-CBD	-2.42	107.67	113.06
2	A	501	MLD	OBA-CAP-OAQ	-2.35	118.76	124.08
3	A	502	3V5	NAB-CAJ-NAN	-2.33	110.81	113.66
2	A	501	MLD	CB-CA-N	2.29	114.59	110.38

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	MLD	O5-C1-C2-N2
2	A	501	MLD	CA1-CA2-NAD-CAE
2	A	501	MLD	OAH-CAI-CAL-OAM
2	A	501	MLD	OAH-CAI-CAL-N
2	A	501	MLD	CAJ-CAI-CAL-OAM
2	A	501	MLD	CAJ-CAI-CAL-N
2	A	501	MLD	NBW-CBS-CBT-OBV
3	A	502	3V5	OAF-CAS-CAX-NAO
3	A	502	3V5	OAH-CAS-CAX-NAO
3	A	502	3V5	CAM-CAW-SAQ-CAK
3	A	502	3V5	CAM-CAV-CAY-CAT
3	A	502	3V5	NAO-CAV-CAY-CAR
3	A	502	3V5	NAO-CAV-CAY-CAT
2	A	501	MLD	NAN-CAO-CBB-CBC
2	A	501	MLD	NBW-CBS-CBT-OBV
2	A	501	MLD	CAF-CAE-NAD-CA2
2	A	501	MLD	OAG-CAE-NAD-CA2
2	A	501	MLD	CAP-CAO-CBB-CBC
2	A	501	MLD	CA3-CA4-O1-C1
2	A	501	MLD	CA5-CA4-O1-C1
2	A	501	MLD	CBP-CBQ-CBR-CBS
3	A	502	3V5	OAE-CAR-CAY-CAT
2	A	501	MLD	CBB-CBC-CBD-NBF
2	A	501	MLD	CBB-CBC-CBD-OBE
2	A	501	MLD	CBR-CBS-CBT-OBV
2	A	501	MLD	CBR-CBS-CBT-OBV
3	A	502	3V5	CAM-CAV-CAY-CAR


There are no ring outliers.


2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	3V5	4	0
2	A	501	MLD	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	354:CYS	C	355:LEU	N	1.00

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	349/349 (100%)	0.37	22 (6%)	26 24	21, 34, 60, 78	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	TYR	5.8
1	A	305	SER	4.4
1	A	133	LEU	4.0
1	A	303	MET	3.9
1	A	310	VAL	3.7
1	A	307	THR	3.5
1	A	132	GLY	3.4
1	A	265	THR	3.4
1	A	318	TYR	3.3
1	A	312	VAL	3.1
1	A	136	ALA	3.0
1	A	244	GLN	2.9
1	A	319	ARG	2.9
1	A	311	PRO	2.9
1	A	317	GLY	2.9
1	A	59	VAL	2.8
1	A	320	THR	2.6
1	A	302	ILE	2.3
1	A	315	PRO	2.3
1	A	304	ASP	2.3
1	A	300	HIS	2.2
1	A	149	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MLD	A	501	46/64	0.64	0.22	30,71,97,105	0
3	3V5	A	502	19/19	0.67	0.20	53,79,98,106	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 3V5 A 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.