



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:52 PM UTC

PDB ID : 5QTH / pdb_00005qth
Title : T. brucei FPPS in complex with CID 2799882
Authors : Muenzker, L.; Petrick, J.K.; Schleberger, C.; Cornaciu, I.; Marquez, J.A.; Jahnke, W.
Deposited on : 2019-08-09
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PJY	A	404	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2769 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Farnesyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2623	1675	425	496	27	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q86C09
A	0	PRO	-	expression tag	UNP Q86C09

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is N,N-dimethyl-2-(methylsulfanyl)pyridine-3-carboxamide (CCD ID: PJY) (formula: $C_9H_{12}N_2OS$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			13	9	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	117	Total 117	O 117	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Farnesyl pyrophosphate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	60.56Å 60.56Å 339.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	56.64 – 1.89 56.64 – 1.89	Depositor EDS
% Data completeness (in resolution range)	69.3 (56.64-1.89) 69.2 (56.64-1.89)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	0.27	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.88Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.241 , 0.283 0.241 , 0.281	Depositor DCC
R_{free} test set	1060 reflections (3.24%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2769	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PJY, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/2673	0.31	0/3611

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2623	0	2579	16	0
2	A	10	0	0	0	0
3	A	6	0	8	0	0
4	A	13	0	0	0	0
5	A	117	0	0	1	0
All	All	2769	0	2587	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:ARG:HG2	1:A:80:ARG:HH21	1.59	0.67
1:A:239:VAL:HG13	1:A:340:LEU:HD22	1.79	0.64
1:A:160:ARG:HD2	1:A:227:ILE:HD11	1.88	0.55
1:A:337:VAL:HG11	1:A:359:TRP:CD2	2.42	0.54
1:A:52:LEU:HD23	1:A:55:ILE:HD12	1.88	0.54
1:A:148:PHE:HB2	1:A:154:LEU:HD13	1.93	0.49
1:A:10:TYR:HB2	1:A:90:TRP:CZ2	2.49	0.48
1:A:241:GLU:O	1:A:245:MET:HG2	2.13	0.47
1:A:108:ASN:ND2	5:A:513:HOH:O	2.47	0.46
1:A:61:LEU:HD21	1:A:226:LEU:HD23	1.99	0.44
1:A:277:ALA:HA	1:A:302:TYR:CE2	2.52	0.44
1:A:210:LYS:HG3	1:A:245:MET:SD	2.58	0.43
1:A:167:THR:HG23	1:A:211:TYR:CD1	2.54	0.42
1:A:27:MET:HG2	1:A:32:VAL:HG23	2.02	0.41
1:A:20:GLU:HG3	1:A:24:LYS:HD2	2.02	0.41
1:A:250:TYR:CE2	1:A:330:GLU:HB3	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/369 (87%)	313 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/322 (88%)	282 (99%)	3 (1%)	65	56

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150	ASP
1	A	179	MET
1	A	255	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	131	ASN
1	A	172	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PJY	A	404	-	13,13,13	0.68	0	16,17,17	1.80	3 (18%)
3	GOL	A	403	-	5,5,5	0.95	0	5,5,5	1.08	0
2	PO4	A	401	-	4,4,4	0.96	0	6,6,6	0.47	0
2	PO4	A	402	-	4,4,4	0.93	0	6,6,6	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	403	-	-	0/4/4/4	-
4	PJY	A	404	-	-	0/10/10/10	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	404	PJY	C1-S2-C3	-4.00	97.70	101.69
4	A	404	PJY	C8-C9-N11	3.32	121.78	118.32
4	A	404	PJY	C5-N4-C3	2.82	121.60	116.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/369 (89%)	1.60	107 (32%) 1 1	10, 38, 72, 95	4 (1%)

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	TYR	14.9
1	A	251	PHE	14.5
1	A	50	ARG	10.3
1	A	351	PHE	9.8
1	A	64	LEU	7.0
1	A	76	ALA	5.9
1	A	62	LEU	5.9
1	A	346	LEU	5.6
1	A	182	SER	5.6
1	A	232	LEU	5.4
1	A	1	MET	5.4
1	A	61	LEU	5.2
1	A	293	ALA	5.2
1	A	60	SER	5.1
1	A	344	LEU	5.1
1	A	235	VAL	5.1
1	A	122	PRO	4.9
1	A	349	PRO	4.9
1	A	231	ALA	4.7
1	A	359	TRP	4.5
1	A	350	GLY	4.5
1	A	292	SER	4.4
1	A	181	ASP	4.3
1	A	65	SER	4.2
1	A	338	LYS	4.1
1	A	121	HIS	4.1
1	A	59	GLU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	270	VAL	4.0
1	A	196	ASP	4.0
1	A	353	ALA	4.0
1	A	354	SER	3.8
1	A	75	GLY	3.8
1	A	63	SER	3.7
1	A	58	ALA	3.7
1	A	56	ASP	3.7
1	A	290	ALA	3.6
1	A	180	PHE	3.6
1	A	328	ALA	3.6
1	A	362	THR	3.6
1	A	106	MET	3.5
1	A	356	GLU	3.5
1	A	233	PRO	3.4
1	A	234	THR	3.4
1	A	272	THR	3.4
1	A	341	ILE	3.4
1	A	355	VAL	3.4
1	A	340	LEU	3.3
1	A	57	VAL	3.3
1	A	108	ASN	3.3
1	A	183	ASN	3.3
1	A	352	ALA	3.2
1	A	307	SER	3.1
1	A	345	ARG	3.1
1	A	347	CYS	3.1
1	A	331	ALA	3.1
1	A	263	PRO	3.1
1	A	337	VAL	3.0
1	A	335	GLU	3.0
1	A	77	ARG	2.9
1	A	228	VAL	2.9
1	A	2	PRO	2.9
1	A	268	GLY	2.9
1	A	47	LYS	2.8
1	A	343	LYS	2.8
1	A	342	GLU	2.8
1	A	226	LEU	2.7
1	A	152	PRO	2.7
1	A	107	ASP	2.7
1	A	348	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	179	MET	2.7
1	A	81	VAL	2.7
1	A	318	GLU	2.6
1	A	339	GLU	2.6
1	A	124	VAL	2.6
1	A	357	THR	2.6
1	A	15	MET	2.5
1	A	237	MET	2.5
1	A	104	ASP	2.5
1	A	265	GLU	2.4
1	A	295	VAL	2.4
1	A	123	ASP	2.4
1	A	153	PHE	2.4
1	A	150	ASP	2.3
1	A	250	TYR	2.3
1	A	264	PRO	2.3
1	A	230	GLU	2.3
1	A	4	GLN	2.3
1	A	336	GLN	2.3
1	A	296	ALA	2.3
1	A	236	ASP	2.3
1	A	176	VAL	2.3
1	A	308	GLU	2.2
1	A	239	VAL	2.2
1	A	82	LEU	2.2
1	A	332	ALA	2.2
1	A	80	ARG	2.2
1	A	229	SER	2.2
1	A	322	LEU	2.1
1	A	197	PHE	2.1
1	A	326	TYR	2.1
1	A	78	ARG	2.1
1	A	333	VAL	2.1
1	A	262	THR	2.1
1	A	271	GLY	2.1
1	A	327	VAL	2.0
1	A	238	GLY	2.0
1	A	314	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PJY	A	404	13/13	0.73	0.52	44,49,51,52	13
2	PO4	A	402	5/5	0.76	0.26	90,92,93,94	0
3	GOL	A	403	6/6	0.82	0.17	52,57,59,61	0
2	PO4	A	401	5/5	0.82	0.19	80,83,84,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.