



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:39 AM UTC

PDB ID : 5QTQ / pdb_00005qtq
Title : PanDDA analysis group deposition – Crystal Structure of NUDT5 in complex with FS-1169
Authors : Dubianok, Y.; Krojer, T.; Kovacs, H.; Moriaud, F.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; von Delft, F.
Deposited on : 2019-08-14
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	15% 76% 15% 7%
1	B	209	11% 76% 16% 7%
1	C	209	12% 78% 13% 8%
1	D	209	16% 76% 17% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1470	929	245	288	8			
1	B	195	Total	C	N	O	S	0	0	0
			1481	934	248	291	8			
1	C	193	Total	C	N	O	S	0	0	0
			1428	899	239	282	8			
1	D	194	Total	C	N	O	S	0	2	0
			1481	937	248	288	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 5-fluoroquinazolin-4(3H)-one (CCD ID: PWM) (formula: C₈H₅FN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			12	8	1	2	1		
3	B	1	Total	C	F	N	O	0	0
			12	8	1	2	1		
3	C	1	Total	C	F	N	O	0	0
			12	8	1	2	1		
3	D	1	Total	C	F	N	O	0	0
			12	8	1	2	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	82	Total O 82 82	0	0
6	B	75	Total O 75 75	0	0
6	C	77	Total O 77 77	0	0
6	D	53	Total O 53 53	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase

Chain A: 




- Molecule 1: ADP-sugar pyrophosphatase

Chain B: 




- Molecule 1: ADP-sugar pyrophosphatase

Chain C: 



- Molecule 1: ADP-sugar pyrophosphatase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.00Å 59.70Å 80.19Å 79.38° 81.67° 75.86°	Depositor
Resolution (Å)	47.30 – 1.82 47.30 – 1.82	Depositor EDS
% Data completeness (in resolution range)	96.2 (47.30-1.82) 96.3 (47.30-1.82)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.211 , 0.250 (Not available) , 0.253	Depositor DCC
R_{free} test set	3877 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6228	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PWM, MG, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	0/1498	1.25	2/2042 (0.1%)
1	B	1.08	1/1510 (0.1%)	1.30	1/2059 (0.0%)
1	C	1.10	1/1456 (0.1%)	1.36	1/1993 (0.1%)
1	D	1.06	1/1511 (0.1%)	1.31	0/2062
All	All	1.08	3/5975 (0.1%)	1.30	4/8156 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	70	ARG	C-O	5.52	1.30	1.24
1	D	26	GLY	N-CA	5.12	1.48	1.44
1	C	81	LYS	C-O	5.11	1.30	1.24

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	38	ASP	CA-CB-CG	6.07	118.67	112.60
1	B	38	ASP	CB-CA-C	5.85	120.05	109.45
1	A	152	ASP	CA-C-N	5.06	127.37	120.54
1	A	152	ASP	C-N-CA	5.06	127.37	120.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1470	0	1441	23	0
1	B	1481	0	1440	27	0
1	C	1428	0	1341	23	0
1	D	1481	0	1424	34	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	12	0	0	0	0
3	B	12	0	0	0	0
3	C	12	0	0	0	0
3	D	12	0	0	0	0
4	A	8	0	12	0	0
4	B	4	0	6	0	0
4	C	12	0	18	0	0
5	B	1	0	0	0	0
6	A	82	0	0	1	0
6	B	75	0	0	1	0
6	C	77	0	0	4	0
6	D	53	0	0	4	0
All	All	6228	0	5682	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:125:GLU:OE1	6:C:401:HOH:O	1.90	0.88
1:A:104:THR:HG23	1:A:107:ALA:H	1.41	0.84
1:C:206:HIS:HD2	1:D:200:TYR:OH	1.65	0.79
1:C:104:THR:HG23	1:C:107:ALA:H	1.49	0.77
1:D:104:THR:HG21	6:D:650:HOH:O	1.85	0.77
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.67	0.76
1:D:120:LYS:H	1:D:155:ASN:HD21	1.32	0.75
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.73	0.71
1:A:176:ASN:OD1	6:A:401:HOH:O	2.10	0.68
6:C:401:HOH:O	1:D:206:HIS:HE1	1.75	0.68
1:D:149:ASN:C	1:D:149:ASN:HD22	1.99	0.68
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.76	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:57:GLN:HE21	1:C:57:GLN:H	1.44	0.65
1:A:104:THR:CG2	1:A:107:ALA:H	2.11	0.62
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.36	0.60
1:D:104:THR:HG23	1:D:107:ALA:H	1.66	0.60
1:C:188:GLU:C	1:C:189:GLU:HG3	2.27	0.60
1:B:104:THR:HG23	1:B:107:ALA:H	1.66	0.59
1:B:17:ILE:HD12	1:B:34:THR:CG2	2.34	0.57
1:D:208:ASN:CG	6:D:630:HOH:O	2.47	0.56
1:A:54:ARG:HD3	1:A:60:ASP:OD1	2.05	0.56
1:B:58:THR:CB	1:B:142:HIS:NE2	2.69	0.56
1:D:149:ASN:HD21	1:D:151:ASP:HB2	1.71	0.56
1:C:194:ASP:OD2	1:C:196:ARG:NH2	2.38	0.56
1:A:206:HIS:HD2	1:B:200:TYR:OH	1.89	0.56
1:C:132:MET:HE1	1:C:196:ARG:CZ	2.35	0.56
1:C:206:HIS:CD2	1:D:200:TYR:OH	2.55	0.55
1:D:191:LEU:C	1:D:191:LEU:HD12	2.32	0.55
1:B:57:GLN:OE1	1:B:99:ILE:O	2.24	0.55
1:A:179:LEU:HD23	1:A:205:LYS:HD2	1.89	0.54
1:C:74:TYR:HB3	6:C:431:HOH:O	2.08	0.52
1:A:80:VAL:HB	1:A:168:VAL:CG1	2.40	0.51
1:D:149:ASN:C	1:D:149:ASN:ND2	2.69	0.51
1:D:64:VAL:O	1:D:94:PHE:HB3	2.10	0.51
1:B:179:LEU:CD2	1:B:205:LYS:HE2	2.42	0.50
1:A:68:LEU:HG	1:A:70:ARG:HD2	1.94	0.50
1:D:37:MET:SD	1:D:41:GLY:O	2.71	0.49
1:D:49:VAL:O	1:D:137:SER:HA	2.12	0.49
1:D:149:ASN:HD22	1:D:151:ASP:H	1.61	0.49
1:C:71:THR:OG1	1:C:151:ASP:OD2	2.29	0.48
1:C:191:LEU:HD12	1:C:191:LEU:C	2.38	0.48
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.48	0.48
1:D:104:THR:CG2	1:D:107:ALA:H	2.25	0.48
1:C:152:ASP:OD1	1:C:154:GLU:N	2.39	0.48
1:B:38:ASP:OD1	1:B:38:ASP:C	2.57	0.47
1:B:158:PRO:O	1:B:160:PRO:HD3	2.14	0.47
1:C:206:HIS:HE1	1:D:125:GLU:OE1	1.97	0.47
1:C:58:THR:CB	1:C:142:HIS:NE2	2.77	0.47
1:A:61:GLY:O	1:A:141:ILE:HA	2.15	0.47
1:B:104:THR:HG22	1:B:107:ALA:CB	2.44	0.47
1:D:17:ILE:HD12	1:D:34:THR:CG2	2.45	0.47
1:B:81:LYS:HA	1:B:91:CYS:O	2.15	0.46
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:205:LYS:O	6:D:601:HOH:O	2.21	0.45
1:B:179:LEU:HD22	1:B:205:LYS:HE2	1.97	0.45
1:B:14:LYS:HE3	1:B:39:PRO:HB2	1.99	0.45
1:C:104:THR:HG23	1:C:107:ALA:N	2.26	0.45
1:D:132:MET:HE1	1:D:196:ARG:CZ	2.46	0.45
1:A:34:THR:O	1:A:45:THR:HA	2.18	0.44
1:A:64:VAL:O	1:A:66:PRO:HD3	2.17	0.44
1:A:125:GLU:OE1	1:B:206:HIS:HE1	2.01	0.44
1:D:121:GLY:HA3	1:D:146:VAL:CG1	2.47	0.44
1:D:149:ASN:ND2	1:D:151:ASP:H	2.15	0.44
1:A:39:PRO:HG3	1:B:167:PHE:CG	2.52	0.44
1:D:208:ASN:CB	6:D:630:HOH:O	2.65	0.44
1:D:34:THR:O	1:D:45:THR:HA	2.18	0.44
1:A:206:HIS:HE1	1:B:125:GLU:OE1	2.02	0.43
1:A:85:PRO:HB2	1:A:86:PRO:HD3	2.01	0.43
1:A:188:GLU:C	1:A:189:GLU:HG3	2.43	0.43
1:D:38:ASP:OD1	1:D:40:THR:N	2.51	0.43
1:B:17:ILE:HD12	1:B:34:THR:HG21	2.01	0.43
1:B:194:ASP:CG	1:B:196:ARG:HH21	2.26	0.43
1:C:104:THR:HG22	1:C:107:ALA:HB2	2.01	0.42
1:B:82:GLN:NE2	6:B:405:HOH:O	2.51	0.42
1:C:200:TYR:OH	1:D:206:HIS:HD2	2.02	0.41
1:A:49:VAL:HG21	1:B:49:VAL:HG11	2.02	0.41
1:D:188:GLU:HG3	1:D:189:GLU:HG3	2.01	0.41
1:D:38:ASP:OD1	1:D:38:ASP:C	2.63	0.41
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.56	0.41
1:A:110:LEU:HD11	1:A:123:ILE:HG12	2.03	0.41
1:A:162:PRO:HB2	1:A:166:GLU:O	2.20	0.41
1:D:120:LYS:H	1:D:155:ASN:ND2	2.09	0.41
1:A:203:ALA:HB2	1:B:200:TYR:CD1	2.56	0.41
1:B:61:GLY:O	1:B:141:ILE:HA	2.21	0.41
1:C:49:VAL:O	1:C:137:SER:HA	2.20	0.41
1:D:72:LEU:N	1:D:72:LEU:CD1	2.84	0.41
1:D:15:GLN:NE2	1:D:39:PRO:HG3	2.35	0.40
1:B:80:VAL:HB	1:B:168:VAL:CG1	2.51	0.40
1:B:104:THR:HG22	1:B:107:ALA:HB3	2.03	0.40
1:C:196:ARG:NE	6:C:404:HOH:O	2.37	0.40
1:A:200:TYR:OH	1:B:206:HIS:HD2	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/209 (92%)	185 (96%)	6 (3%)	1 (0%)	24	14
1	B	193/209 (92%)	187 (97%)	6 (3%)	0	100	100
1	C	191/209 (91%)	186 (97%)	4 (2%)	1 (0%)	24	14
1	D	192/209 (92%)	185 (96%)	7 (4%)	0	100	100
All	All	768/836 (92%)	743 (97%)	23 (3%)	2 (0%)	36	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	66	PRO
1	A	66	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/179 (87%)	151 (97%)	5 (3%)	34	16
1	B	157/179 (88%)	150 (96%)	7 (4%)	24	8
1	C	145/179 (81%)	138 (95%)	7 (5%)	23	6
1	D	155/179 (87%)	152 (98%)	3 (2%)	50	37
All	All	613/716 (86%)	591 (96%)	22 (4%)	31	13

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ASP
1	A	43	THR
1	A	70	ARG
1	A	166	GLU
1	A	191	LEU
1	B	27	LYS
1	B	31	LEU
1	B	38	ASP
1	B	43	THR
1	B	57	GLN
1	B	205	LYS
1	B	208	ASN
1	C	17	ILE
1	C	31	LEU
1	C	45	THR
1	C	57	GLN
1	C	71	THR
1	C	164	ASP
1	C	189	GLU
1	D	43	THR
1	D	149	ASN
1	D	168	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	206	HIS
1	B	15	GLN
1	B	57	GLN
1	B	82	GLN
1	B	190	HIS
1	B	206	HIS
1	B	208	ASN
1	C	57	GLN
1	C	206	HIS
1	D	15	GLN
1	D	149	ASN
1	D	155	ASN
1	D	206	HIS
1	D	208	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 9 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	C	304	-	3,3,3	0.04	0	2,2,2	0.12	0
4	EDO	B	302	-	3,3,3	0.28	0	2,2,2	0.10	0
4	EDO	A	304	-	3,3,3	0.11	0	2,2,2	0.04	0
3	PWM	B	305	-	13,13,13	2.12	6 (46%)	15,18,18	2.98	3 (20%)
4	EDO	C	301	-	3,3,3	0.22	0	2,2,2	0.10	0
4	EDO	A	305	-	3,3,3	0.14	0	2,2,2	0.10	0
3	PWM	C	305	-	13,13,13	2.04	7 (53%)	15,18,18	3.08	4 (26%)
3	PWM	D	501	-	13,13,13	2.38	6 (46%)	15,18,18	2.95	3 (20%)
4	EDO	C	306	-	3,3,3	0.12	0	2,2,2	0.02	0
3	PWM	A	303	-	13,13,13	2.24	5 (38%)	15,18,18	3.14	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	304	-	-	0/1/1/1	-
4	EDO	A	304	-	-	0/1/1/1	-
3	PWM	B	305	-	-	-	0/2/2/2
4	EDO	C	301	-	-	0/1/1/1	-
4	EDO	A	305	-	-	0/1/1/1	-
3	PWM	C	305	-	-	-	0/2/2/2
3	PWM	D	501	-	-	-	0/2/2/2
3	PWM	A	303	-	-	-	0/2/2/2
4	EDO	C	306	-	-	0/1/1/1	-
4	EDO	B	302	-	-	0/1/1/1	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	PWM	O-C6	4.88	1.34	1.23
3	D	501	PWM	O-C6	4.73	1.34	1.23
3	C	305	PWM	O-C6	3.74	1.32	1.23
3	B	305	PWM	O-C6	3.73	1.32	1.23
3	D	501	PWM	C2-C1	3.53	1.44	1.38
3	D	501	PWM	C5-N	3.18	1.35	1.29
3	B	305	PWM	C5-N	3.15	1.35	1.29
3	A	303	PWM	C3-C4	3.12	1.44	1.40
3	C	305	PWM	C7-C6	-3.08	1.41	1.47
3	B	305	PWM	C2-C1	2.98	1.44	1.38
3	A	303	PWM	C5-N	2.91	1.34	1.29
3	C	305	PWM	C7-C	2.73	1.43	1.39
3	A	303	PWM	C7-C	2.71	1.43	1.39
3	D	501	PWM	C7-C	2.68	1.43	1.39
3	D	501	PWM	C6-N1	-2.61	1.33	1.38
3	B	305	PWM	C5-N1	2.60	1.40	1.35
3	B	305	PWM	C7-C6	-2.48	1.42	1.47
3	C	305	PWM	C5-N	2.43	1.33	1.29
3	B	305	PWM	C6-N1	-2.38	1.33	1.38
3	D	501	PWM	C5-N1	2.36	1.39	1.35
3	A	303	PWM	C2-C1	2.36	1.42	1.38
3	C	305	PWM	C2-C1	2.35	1.42	1.38
3	C	305	PWM	C5-N1	2.20	1.39	1.35
3	C	305	PWM	C3-C4	2.10	1.43	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	305	PWM	C7-C6-N1	8.36	124.98	115.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	303	PWM	C5-N1-C6	-7.55	114.35	123.42
3	A	303	PWM	C7-C6-N1	7.47	123.94	115.14
3	B	305	PWM	C7-C6-N1	7.41	123.87	115.14
3	D	501	PWM	C7-C6-N1	7.14	123.55	115.14
3	B	305	PWM	C5-N1-C6	-7.08	114.91	123.42
3	C	305	PWM	C5-N1-C6	-6.91	115.11	123.42
3	D	501	PWM	C5-N1-C6	-6.75	115.30	123.42
3	D	501	PWM	O-C6-N1	-5.30	113.98	120.44
3	A	303	PWM	O-C6-N1	-5.13	114.19	120.44
3	B	305	PWM	O-C6-N1	-4.54	114.92	120.44
3	C	305	PWM	O-C6-N1	-3.62	116.03	120.44
3	C	305	PWM	O-C6-C7	-2.38	118.95	124.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/209 (92%)	1.12	31 (15%) 5 5	16, 27, 42, 67	27 (13%)
1	B	195/209 (93%)	0.62	22 (11%) 10 11	15, 31, 59, 78	10 (5%)
1	C	193/209 (92%)	0.79	26 (13%) 7 7	19, 33, 54, 67	14 (7%)
1	D	194/209 (92%)	1.17	34 (17%) 4 4	7, 33, 58, 67	23 (11%)
All	All	776/836 (92%)	0.92	113 (14%) 6 6	7, 31, 57, 78	74 (9%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	16[A]	TYR	14.6
1	A	191	LEU	12.1
1	D	73[A]	HIS	11.7
1	A	156	ALA	10.6
1	A	14	LYS	10.6
1	A	165	GLY	10.5
1	D	207	ALA	10.5
1	A	158	PRO	10.1
1	A	153	ALA	9.6
1	A	167	PHE	9.6
1	A	157	ARG	9.4
1	A	159	LYS	8.7
1	A	155	ASN	8.5
1	A	187	ALA	8.4
1	A	189	GLU	8.2
1	B	180	GLN	7.7
1	A	166	GLU	7.2
1	A	190	HIS	7.2
1	A	188	GLU	7.1
1	A	154	GLU	7.1
1	D	208	ASN	6.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	43	THR	6.7
1	D	74	TYR	6.4
1	A	164	ASP	6.4
1	B	208	ASN	5.9
1	D	114	GLU	5.4
1	B	176	ASN	5.3
1	D	154	GLU	5.2
1	D	122	ASP	5.2
1	C	16	TYR	5.1
1	B	56	GLU	4.5
1	C	39	PRO	4.4
1	C	56	GLU	4.4
1	C	40	THR	4.2
1	B	40	THR	3.9
1	B	165	GLY	3.7
1	D	58	THR	3.7
1	A	42	LYS	3.5
1	C	23	ILE	3.5
1	C	31	LEU	3.5
1	B	191	LEU	3.4
1	C	17	ILE	3.4
1	D	23	ILE	3.4
1	C	43	THR	3.3
1	A	56	GLU	3.2
1	C	55	LYS	3.1
1	D	160	PRO	3.1
1	A	22	LEU	3.1
1	D	40	THR	3.1
1	D	152	ASP	3.1
1	D	156	ALA	3.1
1	C	74	TYR	3.1
1	C	104	THR	3.1
1	C	207	ALA	3.0
1	D	57	GLN	3.0
1	C	102	GLY	2.9
1	D	29	VAL	2.9
1	B	58	THR	2.9
1	C	28	TRP	2.9
1	A	54	ARG	2.9
1	C	22	LEU	2.9
1	A	163	GLY	2.9
1	A	162	PRO	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	162	PRO	2.8
1	A	72	LEU	2.8
1	B	102	GLY	2.8
1	B	54	ARG	2.6
1	D	167	PHE	2.6
1	D	56	GLU	2.6
1	D	101	ASP	2.6
1	B	27	LYS	2.6
1	A	40	THR	2.6
1	B	190	HIS	2.5
1	C	190	HIS	2.5
1	B	167	PHE	2.5
1	C	167	PHE	2.5
1	A	18	ILE	2.5
1	A	31	LEU	2.4
1	A	58	THR	2.4
1	B	39	PRO	2.4
1	C	57	GLN	2.4
1	D	161	LYS	2.4
1	B	160	PRO	2.4
1	D	46	TRP	2.4
1	C	53	THR	2.4
1	B	157	ARG	2.3
1	A	33	LYS	2.3
1	C	29	VAL	2.3
1	D	100	ASP	2.3
1	C	35	THR	2.3
1	D	53	THR	2.3
1	D	41	GLY	2.3
1	B	14	LYS	2.3
1	D	52	THR	2.3
1	C	165	GLY	2.2
1	C	54	ARG	2.2
1	D	39	PRO	2.2
1	D	28	TRP	2.2
1	C	58	THR	2.2
1	B	101	ASP	2.2
1	B	28	TRP	2.2
1	D	172	SER	2.2
1	D	97	GLY	2.2
1	D	54	ARG	2.1
1	D	55	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	149	ASN	2.1
1	D	49	VAL	2.1
1	A	160	PRO	2.1
1	C	168	VAL	2.0
1	B	153	ALA	2.0
1	C	153	ALA	2.0
1	D	164	ASP	2.0
1	B	166	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	C	301	4/4	0.67	0.20	47,61,61,67	0
4	EDO	B	302	4/4	0.68	0.19	46,48,50,50	0
2	MG	B	304	1/1	0.83	0.13	64,64,64,64	0
4	EDO	C	304	4/4	0.86	0.13	30,31,33,36	4
2	MG	A	302	1/1	0.87	0.10	57,57,57,57	0
5	CL	B	301	1/1	0.87	0.18	62,62,62,62	0
4	EDO	C	306	4/4	0.89	0.10	29,30,30,30	4
3	PWM	D	501	12/12	0.90	0.11	33,35,37,37	12
3	PWM	B	305	12/12	0.91	0.10	26,28,31,32	12
3	PWM	C	305	12/12	0.91	0.11	31,39,41,43	12
4	EDO	A	305	4/4	0.92	0.10	27,27,28,29	4
3	PWM	A	303	12/12	0.94	0.09	26,29,34,36	12
2	MG	C	302	1/1	0.94	0.07	42,42,42,42	0
4	EDO	A	304	4/4	0.94	0.08	21,21,22,22	4
2	MG	D	503	1/1	0.97	0.09	39,39,39,39	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	301	1/1	0.97	0.04	29,29,29,29	0
2	MG	C	303	1/1	0.98	0.03	31,31,31,31	0
2	MG	B	303	1/1	0.99	0.04	28,28,28,28	0
2	MG	D	502	1/1	0.99	0.06	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.