



Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 05:11 PM UTC

PDB ID : 7QTR / pdb_00007qtr
BMRB ID : 34700
Title : GB1 in mammalian cells, 50 uM
Authors : Gerez, J.A.; Prymaczok, N.C.; Kadavath, H.; Gosh, D.; Butikofer, M.; Guntert, P.; Riek, R.
Deposited on : 2022-01-15

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 52%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	57	88% 7% 5%

2 Ensemble composition and analysis

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:4-A:57 (54)	0.89	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 7, 8, 9, 12, 13, 14, 16, 17, 19, 20
2	3, 10, 11
3	15, 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 872 atoms, of which 427 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Immunoglobulin G-binding protein G.

Mol	Chain	Residues	Atoms						Trace
1	A	57	Total	C	H	N	O	S	0
			872	279	427	69	95	2	

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P06654
A	2	MET	-	expression tag	UNP P06654
A	3	GLN	-	expression tag	UNP P06654

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Immunoglobulin G-binding protein G




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Immunoglobulin G-binding protein G



4.2.2 Score per residue for model 2


- Molecule 1: Immunoglobulin G-binding protein G



4.2.3 Score per residue for model 3

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  82% 12% 5%



4.2.4 Score per residue for model 4

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  84% 9% 5%



4.2.5 Score per residue for model 5

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  82% 9% 5%



4.2.6 Score per residue for model 6

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  82% 12% 5%



4.2.7 Score per residue for model 7

- Molecule 1: Immunoglobulin G-binding protein G


Chain A:  88% 7% 5%



4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  86% 9% 5%



4.2.9 Score per residue for model 9

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  82% 11% 5%



4.2.10 Score per residue for model 10

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  75% 19% 5%



4.2.11 Score per residue for model 11

- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  75% 19% 5%



4.2.12 Score per residue for model 12


- Molecule 1: Immunoglobulin G-binding protein G

Chain A:  84% 11% 5%




4.2.13 Score per residue for model 13

- Molecule 1: Immunoglobulin G-binding protein G




4.2.14 Score per residue for model 14

- Molecule 1: Immunoglobulin G-binding protein G




4.2.15 Score per residue for model 15

- Molecule 1: Immunoglobulin G-binding protein G




4.2.16 Score per residue for model 16

- Molecule 1: Immunoglobulin G-binding protein G




4.2.17 Score per residue for model 17

- Molecule 1: Immunoglobulin G-binding protein G




4.2.18 Score per residue for model 18

- Molecule 1: Immunoglobulin G-binding protein G




4.2.19 Score per residue for model 19

- Molecule 1: Immunoglobulin G-binding protein G



4.2.20 Score per residue for model 20

- Molecule 1: Immunoglobulin G-binding protein G



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	3.98.13
OPALp	refinement	1.4

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	372
Number of shifts mapped to atoms	372
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	52%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.63±0.02	0±0/426 (0.0± 0.0%)	1.43±0.04	2±1/578 (0.3± 0.2%)
All	All	0.63	0/8520 (0.0%)	1.43	36/11560 (0.3%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	25	ALA	CA-C-N	7.78	130.55	120.44	4	4
1	A	25	ALA	C-N-CA	7.78	130.55	120.44	4	4
1	A	53	PHE	CA-CB-CG	7.67	121.47	113.80	6	2
1	A	55	VAL	CB-CA-C	7.18	120.97	110.77	16	2
1	A	40	VAL	CB-CA-C	6.48	117.11	110.70	16	3
1	A	47	ASP	CA-CB-CG	-6.46	106.14	112.60	17	2
1	A	47	ASP	N-CA-C	5.85	123.27	110.80	13	1
1	A	36	ASN	CA-CB-CG	5.68	118.28	112.60	16	1
1	A	37	ASP	CA-CB-CG	-5.46	107.14	112.60	2	1
1	A	46	TYR	CA-C-N	5.46	131.96	121.54	13	1
1	A	46	TYR	C-N-CA	5.46	131.96	121.54	13	1
1	A	26	THR	N-CA-CB	-5.37	102.22	110.01	15	2
1	A	26	THR	N-CA-C	-5.24	105.56	111.28	14	1
1	A	43	GLU	N-CA-C	5.22	116.66	111.07	2	1
1	A	31	PHE	CA-CB-CG	-5.16	108.64	113.80	20	1
1	A	23	ASP	CA-C-N	5.14	127.42	120.38	13	1
1	A	23	ASP	C-N-CA	5.14	127.42	120.38	13	1
1	A	19	THR	CA-CB-CG2	5.08	119.13	110.50	9	1
1	A	51	LYS	CA-C-N	5.07	127.90	120.71	10	1
1	A	51	LYS	C-N-CA	5.07	127.90	120.71	10	1
1	A	6	LEU	CA-C-N	5.04	131.04	121.97	19	1
1	A	6	LEU	C-N-CA	5.04	131.04	121.97	19	1
1	A	22	VAL	N-CA-C	5.01	115.68	110.82	20	1
1	A	8	LEU	N-CA-C	-5.00	101.59	109.25	11	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	420	401	401	1±1
All	All	8400	8020	8020	15

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:6:LEU:HD21	1:A:8:LEU:HD13	0.62	1.70	5	1
1:A:4:TYR:CD2	1:A:31:PHE:CE2	0.53	2.95	14	1
1:A:4:TYR:C	1:A:4:TYR:CD1	0.49	2.89	20	1
1:A:4:TYR:CE1	1:A:31:PHE:CD2	0.49	3.00	20	1
1:A:21:ALA:HB1	1:A:27:ALA:HB2	0.49	1.84	6	1
1:A:4:TYR:CE1	1:A:31:PHE:CE2	0.47	3.01	20	1
1:A:7:ILE:HB	1:A:54:THR:HG22	0.45	1.89	4	1
1:A:8:LEU:HA	1:A:55:VAL:HG22	0.45	1.87	1	1
1:A:42:GLY:H	1:A:55:VAL:CG2	0.45	2.25	20	1
1:A:4:TYR:CZ	1:A:19:THR:HG23	0.43	2.48	14	1
1:A:40:VAL:HG12	1:A:55:VAL:HG21	0.42	1.91	3	1
1:A:32:LYS:HA	1:A:44:TRP:CH2	0.42	2.49	14	2
1:A:47:ASP:CG	1:A:48:ASP:N	0.41	2.79	18	1
1:A:4:TYR:CE1	1:A:21:ALA:HB2	0.40	2.52	14	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	53/57 (93%)	45±3 (86±5%)	7±2 (13±4%)	1±1 (1±1%)	11	58
All	All	1060/1140 (93%)	907 (86%)	138 (13%)	15 (1%)	11	58

All 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	7	ILE	7
1	A	47	ASP	2
1	A	41	ASP	2
1	A	42	GLY	2
1	A	43	GLU	1
1	A	4	TYR	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	44/47 (94%)	38±2 (88±5%)	6±2 (12±5%)	7	48
All	All	880/940 (94%)	770 (88%)	110 (12%)	7	48

All 32 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	ASN	16
1	A	14	LYS	12
1	A	33	GLN	9
1	A	47	ASP	7
1	A	26	THR	7
1	A	55	VAL	7
1	A	48	ASP	5
1	A	19	THR	5
1	A	56	THR	3
1	A	54	THR	3
1	A	57	GLU	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	18	THR	3
1	A	40	VAL	2
1	A	34	TYR	2
1	A	7	ILE	2
1	A	6	LEU	2
1	A	8	LEU	2
1	A	20	GLU	2
1	A	17	THR	2
1	A	5	LYS	2
1	A	4	TYR	2
1	A	12	THR	2
1	A	50	THR	1
1	A	37	ASP	1
1	A	32	LYS	1
1	A	41	ASP	1
1	A	13	LEU	1
1	A	23	ASP	1
1	A	28	GLU	1
1	A	43	GLU	1
1	A	16	GLU	1
1	A	22	VAL	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 52% for the well-defined parts and 51% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch_output*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	372
Number of shifts mapped to atoms	372
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 52%, i.e. 359 atoms were assigned a chemical shift out of a possible 691. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	111/274 (41%)	111/112 (99%)	0/108 (0%)	0/54 (0%)
Sidechain	229/358 (64%)	229/230 (100%)	0/118 (0%)	0/10 (0%)
Aromatic	19/59 (32%)	19/28 (68%)	0/30 (0%)	0/1 (0%)
Overall	359/691 (52%)	359/370 (97%)	0/256 (0%)	0/65 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 51%, i.e. 372 atoms were assigned a chemical shift out of a possible 736. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	114/289 (39%)	114/118 (97%)	0/114 (0%)	0/57 (0%)
Sidechain	239/388 (62%)	239/250 (96%)	0/127 (0%)	0/11 (0%)
Aromatic	19/59 (32%)	19/28 (68%)	0/30 (0%)	0/1 (0%)
Overall	372/736 (51%)	372/396 (94%)	0/271 (0%)	0/69 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	6	LEU	HB2	-1.34	-0.07 – 3.30	-8.8
1	A	32	LYS	HE2	1.45	1.95 – 3.88	-7.6
1	A	32	LYS	HE3	1.74	1.92 – 3.89	-5.9

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

